

Arithmetic gravity and Yang-Mills theory: An approach to adelic physics via algebraic spaces

René Schmidt

*Mathematisches Institut der Westfälischen Wilhelms-Universität
Einsteinstraße 62, D-48119 Münster, Germany
e-mail¹: rene.schmidt@uni-muenster.de*

Abstract

This work is a dissertation thesis written at the WWU Münster (Germany), supervised by Prof. Dr. Raimar Wulkenhaar. We present an approach to adelic physics based on the language of algebraic spaces. Relative algebraic spaces X over a base S are considered as fundamental objects which describe space-time. This yields a formulation of general relativity which is covariant with respect to changes of the chosen domain of numbers S . With regard to adelic physics the choice of S as an excellent Dedekind scheme is of interest (because this way also the finite prime spots, i.e. the $\mathfrak{p}$ -adic degrees of freedom are taken into account). In this arithmetic case, it turns out that X is a Néron model. This enables us to make concrete statements concerning the structure of the space-time described by X . Furthermore, some solutions of the arithmetic Einstein equations are presented. In a next step, Yang-Mills gauge fields are incorporated.

¹alternative e-mail: rene.schmidt1@gmx.net

Arithmetic gravity and Yang-Mills theory: An approach to adelic physics via algebraic spaces

Inaugural-Dissertation
zur Erlangung des Doktorgrades
der Naturwissenschaften im Fachbereich
Mathematik und Informatik
der Mathematisch-Naturwissenschaftlichen Fakultät
der Westfälischen Wilhelms-Universität Münster

vorgelegt von
Dipl.-Math. René Schmidt

- 2008 -

Danksagung

Die vorliegende Dissertation ist das Ergebnis einer mehrjährigen Forschungsarbeit am Fachbereich Mathematik und Informatik der Mathematisch-Naturwissenschaftlichen Fakultät der Westfälischen Wilhelms-Universität Münster. Die Arbeit wurde ermöglicht durch die finanzielle Unterstützung des Graduiertenkollegs “Analytische Topologie und Metageometrie”.

Mein besonderer Dank richtet sich an meinen Doktorvater Herrn Prof. Dr. Raimar Wulkenhaar für die Betreuung und Förderung meiner mathematischen und physikalischen Studien. Stets hatte er ein offenes Ohr für die Fragestellungen, denen ich mich im Laufe der Promotionsphase stellen musste. Ohne die von ihm erlernten Techniken und Kenntnisse besonders im Bereich der Yang-Mills Theorie wäre das Zustandekommen dieser Dissertation nicht möglich gewesen.

Leider ist es mir an dieser Stelle nicht möglich, all jenen namentlich zu danken, die sowohl während meiner Schulzeit als auch während meiner Zeit an der Universität direkt oder indirekt am Aufbau meines mathematischen und physikalischen Fundamentes beteiligt waren.

Ganz speziell möchte ich mich jedoch bei Herrn Prof. Dr. Siegfried Bosch für die hervorragende Ausbildung bedanken, die ich unter seiner Anleitung im algebraischen Bereich erfahren durfte. In einem aufeinander aufbauenden Vorlesungszyklus, der mich vom ersten Semester bis zum Diplom begleitete, gelang es ihm meine Begeisterung für die algebraische Geometrie zu wecken. Die bei Herrn Prof. Dr. Bosch erlernten Kenntnisse in relativer algebraischer Geometrie einerseits und das von Herrn Prof. Dr. Wulkenhaar vermittelte physikalische Wissen im Bereich der Eichtheorie andererseits bilden das theoretische Fundament, das dieser Dissertation zugrunde liegt.

Nicht unerwähnt lassen möchte ich ferner den Namen von Herrn Dr. Matthias Strauch, der den Vorlesungszyklus von Herrn Prof. Dr. Bosch im Rahmen von Seminaren begleitete. Stets stand er uns Studenten für Fragen zur Verfügung. Durch gemeinsames Rechnen von Übungsaufgaben, teilweise bis spät in den Abend hinein, hat er unschätzbare Beiträge zum Aufbau auch meiner mathematischen Kenntnisse geleistet.

Ein ganz besonders großer und persönlicher Dank gilt schließlich noch meinen Eltern. Sie schufen ein Umfeld, welches mir schon während meines Studiums ermöglichte, mich vollständig auf die wissenschaftliche Arbeit zu konzentrieren.

Contents

0.1. Preface	7
0.2. Introduction	10
0.3. The continuum limit - Comparison with the adelic model	15
 I. Arithmetic theory of general relativity	 20
1. Einstein's equation in the setting of algebraic spaces	21
1.1. Smoothness in algebraic geometry	21
1.2. The arithmetic Einstein equations	29
2. Solutions of the arithmetic Einstein equations	35
2.1. The case of Zariski zero-dimensional base	35
2.1.1. Kasner solution	35
2.1.2. Schwarzschild solution	37
2.1.3. Electromagnetism in vacuum	38
2.1.4. Robertson-Walker models	41
2.2. The case of Zariski one-dimensional base	42
2.2.1. The low dimensional case	42
2.2.2. A higher dimensional solution	43
3. Some properties of the arithmetic models	49
3.1. Etale-invariance	49
3.2. Gravity	51
3.3. The structure of the archimedean component	53
3.4. Boundedness of the archimedean component	57
3.5. Lifting of structures from the archimedean to the adelic level	60
4. The dimensionality of the arithmetic models	65
4.1. The privileged character of four dimensional spaces	66
4.2. Quadratic forms	73
4.3. The metric tensor	76
4.4. Metrical fields of higher spin	86
5. Discreteness of geometry and quantization of gauge fields	89
5.1. The discrete geometry of the arithmetic models	89
5.2. Lattice gauge theory in the arithmetic setting	92

6. Appendix I	98
6.1. Néron models	98
6.2. Further statements on the archimedean component	99
 II. Arithmetic Yang-Mills theory	 102
7. Some essential techniques	103
7.1. General coordinate transformations	103
7.2. Pull-back and push-forward of tensor fields	107
7.2.1. Some formulas	115
8. Gauge theory	119
8.1. The classical, differential geometric theory	119
8.2. Torsors	120
8.3. Gauge transformations	124
8.4. Horizontality	125
8.5. Covariant derivation	129
9. Yang-Mills equation	132
9.1. The Maurer-Cartan form	132
9.2. The field strength of a gauge potential	133
9.2.1. Gauge transformations of the curvature form	134
9.2.2. The curvature form in a local trivialisation	134
9.2.3. The global field strength	138
9.3. Yang-Mills equation	138
10. Yang-Mills theory in local coordinates	141
10.1. Relations between gauge potential and field strength	142
10.2. Yang-Mills equation in local coordinates	146
11. Appendix II	150
11.1. The Zariski tangent space	150
11.2. The tangent bundle	151
11.2.1. The tangent bundle as a functor.	153
11.2.2. The tangent bundle as a scheme.	155
11.3. Differential p -forms and the exterior differential	162
11.4. Group Schemes	167
11.4.1. Definition of group schemes	167
11.4.2. The Lie-algebra as a functor	171
11.4.3. The maps Ad and ad	173
11.4.4. The Lie-algebra and invariant differential forms	174
11.5. Faithfully flat descent	178

Introduction

0.1. Preface

Unless otherwise specified, let $K \subset \mathbb{R}$ be an algebraic number field (i.e. a finite algebraic extension of $\mathbb{Q}$), and let $\mathcal{O}_K$ be the ring of integral numbers of K (i.e. the integral closure of $\mathbb{Z}$ in K). For example, think of $\mathcal{O}_K = \mathbb{Z}$ and $K = \mathbb{Q}$.

In this thesis we present a reformulation of general relativity and (pure) Yang-Mills theory within the bounds of arithmetic algebraic geometry. Relative algebraic spaces $X \rightarrow S$ will be considered as fundamental objects which describe space-time. We will see that one is reduced to the well known theories in the special case $S = \text{Spec } \mathbb{C}$. But one may also make other choices for S . We are especially interested in the case $S = \text{Spec } \mathcal{O}_K$. This choice is motivated as follows.

Since 1987, there have been many interesting applications of p -adic numbers in physics. In his influential paper [Vol1], I.V. Volovich draws the vision of number theory as the ultimate physical theory, where numbers are proposed as the fundamental entities of the universe. It is argued that the development of physics over arbitrary (number) fields might be necessary. In particular, this implies the incorporation of p -adic numbers in physical theories. Since then, many p -adic models have been constructed. A very nice overall view with respect to the various applications of p -adic numbers in mathematic physics may be taken from the book [VVZ].

At first, let us mention that applications of p -adic numbers in quantum physics are of interest ([VV1], [VV2], [VV3], see also [Meu]). Let us recall that there is a canonical way to generalize ordinary quantum mechanics to the p -adic world. As illustrated in [Wey] and [VV2], ordinary one-dimensional quantum mechanics can be given by a triple $(L_2(\mathbb{R}), W_\infty(z), U_\infty(t))$, where $L_2(\mathbb{R})$ is the Hilbert space of complex-valued, square-integrable functions on $\mathbb{R}$, z is a point of the real classical phase space, $W_\infty(z)$ is a unitary representation of the Heisenberg-Weyl group on $L_2(\mathbb{R})$, and $U_\infty(t)$ is a unitary representation of the evolution operator on $L_2(\mathbb{R})$. According to the Vladimirov-Volovich formulation, one-dimensional p -adic quantum mechanics is a triple

$$(L_2(\mathbb{Q}_p), W_p(z), U_p(t))$$

where $L_2(\mathbb{Q}_p)$ is the Hilbert space of complex-valued, square integrable functions on $\mathbb{Q}_p$ (with respect to the Haar measure), and $W_p(z)$ is an unitary representation of the Heisenberg-Weyl group on $L_2(\mathbb{Q}_p)$. Furthermore $U_p(t)$ is an unitary evolution operator

$$U_p(t)\psi_p(x) = \int_{\mathbb{Q}_p} K_p(x, t; y, 0)\psi_p(y)dy$$

on $L_2(\mathbb{Q}_p)$, whose kernel $K_p(x, t; y, 0)$ is defined in complete analogy to the real case by a path integral. The operator $U_p(t)$ and its kernel satisfy the canonical group relations $U_p(t + t') = U_p(t)U_p(t')$ and $K_p(x, t + t'; y, 0) = \int_{\mathbb{Q}_p} K_p(x, t; z, 0)K_p(z, t'; y, 0)dz$. The p -adic Feynman path integral is investigated in [DDN2], where an explicit formula for the kernel of quadratic Lagrangians is derived. This is the basis of a quantum mechanical treatment of the p -adic harmonic oscillator ([Dra1]). Like in the realm of real numbers,

this is an exactly solvable model (see also [Zel1], [Zel2]). However, these general techniques are not limited to non-relativistic quantum mechanics. For example, the free relativistic particle is considered in [DDN1].

With regard to this thesis, gravity is of special interest. In [ADFV], the p -adic version of general relativity in the setting of p -adic differential geometry is considered and cosmological models are studied (see also [Dra2], [Dra3], [DFU], [DN1] - [DN4]). The starting point are the Einstein gravitational field equations

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R + \Lambda g_{\mu\nu} = \kappa T_{\mu\nu}$$

which make sense as well over $\mathbb{R}$ as over $\mathbb{Q}_p$ (for all p) if the constants Λ and κ are assumed to be rational numbers. As an application of p -adic quantum mechanics to the p -adic cosmological models, minisuperspace cosmological models are investigated (see e.g. [DDNV]). For this purpose, Feynman's path integral method in the Hartle-Hawking approach [Haw] was exploited (see [Dra6], [Dra7]). Thereby, the wave function of the universe still takes complex values, but its argument is not necessarily real but also p -adic or even *adelic*. Adeles enable us to regard real and p -adic numbers simultaneously. More precisely, an adèle is an infinite tuple


$$x = (x_2, \dots, x_p, \dots, x_\infty),$$

where $x_\infty \in \mathbb{R}$ and $x_p \in \mathbb{Q}_p$ with the restriction that one has $x_p \in \mathbb{Z}_p$ for all but a finite set of primes. In many papers, which were cited above, not only the p -adic models are formulated, but also the adelic generalization of these models are studied. These adelic models unify in a certain way the ordinary (i.e. $\mathbb{R}$ -valued) and p -adic models. In this sense, adelic models may be considered as very canonical.

Adelic physical models are the starting point of this thesis. But, instead of working directly with adeles and the respective adelic space-time models as it is usually done, we will study a new, purely geometric approach to adelic physics based on relative algebraic spaces $X \rightarrow S$, $S = \text{Spec } \mathcal{O}_K$. However, there are close relations between these two approaches as it may be seen in the following example.

0.1 Example Let us choose $K = \mathbb{Q}$. Consequently, $\mathcal{O}_K = \mathbb{Z}$ and $S = \text{Spec } \mathbb{Z}$. Furthermore assume that the relative algebraic space X over S is representable by a smooth, separated S -scheme, i.e. let us consider a smooth, separated morphism $\pi : X \rightarrow \text{Spec } \mathbb{Z}$ of schemes. Set-theoretically, $\text{Spec } \mathbb{Z}$ consists of infinitely many closed points (one point for each prime number p) plus one generic point which we will denote by ∞ , and which corresponds to the zero ideal of $\mathbb{Z}$. Furthermore, X may be viewed as union $\bigcup_p \pi^{-1}(p) \cup \pi^{-1}(\infty)$ of the fibres of π , and at least set-theoretically we obtain the fol-

lowing picture:



In our arithmetic setting (and in analogy to complex algebraic geometry), a “physical point” x is given by an S -valued point of X , i.e. by a section $s : \text{Spec } \mathbb{Z} \hookrightarrow X$ of the structure morphism π (i.e. $\pi \circ s = \text{id}$). More precisely, x is given by the image of s (this is a closed subscheme of X). However, set-theoretically, s cuts out one closed point in each fibre. Thus, in analogy to the adelic situation, an S -valued point x may be viewed as a set of points:

$$x = \{x_2, \dots, x_p, \dots, x_\infty\}.$$

Furthermore, according to the point of view of adelic physics, each archimedean point (resp. each morphism over the archimedean prime spot at infinity) is only the archimedean component of an adelic point (resp. an adelic morphism). In short, everything in the archimedean world comes from the adelic level. If now $\varphi : Y \rightarrow \text{Spec } \mathbb{Z}$ is an arbitrary smooth S -scheme and if we denote by Y_K the pre-image $\varphi^{-1}(\infty)$ of ∞ under φ , the above extension property (from the archimedean to the adelic level) reads as follows in algebraic geometry:

For every K -morphism $f_K : Y_K \rightarrow X_K$, there is an (★)
 S -morphism $f : Y \rightarrow X$ which extends f_K .

All in all, instead of adeles, the set $X(S)$ of S -valued points of an algebraic space $X \rightarrow S$ is the set of interest in our approach. The objective of this thesis is the investigation of a new approach to general relativity and (pure) Yang-Mills theory based on algebraic spaces. The condition (★) makes clear why Néron models will be of particular interest. We will illustrate in section 0.3 that our approach is naturally settled in the realm of adelic physics. But as the starting point are algebraic spaces, powerful algebraic geometric tools yield interesting, new insight.

0.2. Introduction

According to the theory of general relativity, space-time may be described by means of a differentiable manifold. Thereby, gravity is encoded in a metrical tensor g which satisfies the Einstein equations. More precisely, our starting point are the complex gravitational field equations. Then, any solution of the Einstein equations gives rise to a complex manifold. For technical reasons, we will once and for all assume that this classical space-time manifold may be realized as a compact complex manifold $\mathfrak{X}$ which is *Moishezon*. The latter condition means that

$$\text{transdeg}_{\mathbb{C}}(K(\mathfrak{X})) = \dim_{\mathbb{C}} \mathfrak{X},$$

where $K(\mathfrak{X})$ denotes the field of meromorphic functions on $\mathfrak{X}$. For example, all algebraic manifolds fulfill this equation. Therefore, following the ideas of [ADFV], where it is among other things argued that one should restrict to algebraic manifolds in quantum cosmology, our assumption is not too restrictive. However, let us at least mention that there are Moishezon manifolds which are not algebraic.

The technical reason why we restrict attention to Moishezon manifolds is the following beautiful theorem due to Artin.

0.2 Theorem *There is an equivalence of categories*

$$\left(\text{Moishezon manifolds} \right) \longleftrightarrow \left(\text{smooth, proper algebraic spaces over } \mathbb{C} \right)$$

This theorem enables us to consider the ordinary complex space-time manifold $\mathfrak{X}$ as a complex algebraic space. Now the following observation is crucial. While, on the level of manifolds, the theory is essentially adapted to the complex numbers, the language of algebraic spaces offers the possibility to replace $\mathbb{C}$ by any commutative ring.

In 1987, I.V. Volovich suggested that a fundamental physical theory should be formulated in such a way that it is invariant under change of the underlying number field (see [Vol1]). According to the 2006 paper [Dra5] of B. Dragovich, such a number field invariant model has not yet been constructed. This motivates the following program which will be studied within the first part of this thesis:

- a) Replace the pair $(\mathfrak{X}, g)$ consisting of a (complex) manifold $\mathfrak{X}$ and a metric g by a pair

$$(X \rightarrow S, g),$$

where X is a smooth, separated algebraic space over an arbitrary base S , and where g is a metric over X (see Definition 1.27).

- b) Starting from exactly the same physical principles as in the realm of manifolds, deduce the equations of Einstein's theory of general relativity in the setting of algebraic spaces over an arbitrary base S (thus realizing a number field invariant theory). Determine the pair $(X \rightarrow S, g)$ in such a way that Einstein's equations are fulfilled.

- c) Investigate properties of hypothetical space-time models $(X \rightarrow S, g)$ depending on the choice of the base S .

0.3 Remark Principally, there are many interesting possible choices for S . For example, there is the case of positive characteristic, i.e. S might be chosen as the spectrum of a (finite) field or as function field of an algebraic curve over a finite field. However, in those models $(X \rightarrow S, g)$, which will be studied within the bounds of this thesis, we will often choose S to be representable by an excellent Dedekind-scheme with field of fractions K of characteristic zero (see Definition 0.4). Then the following two cases are of interest:

- a) S is Zariski zero-dimensional and given by the spectrum $\text{Spec } K$ of a field K of characteristic zero. Especially in the case $K = \mathbb{C}$, everything may be translated back into the language of manifolds (by Theorem 0.2).
- b) S is Zariski one-dimensional. In this case we are interested in the choice $S = \text{Spec } \mathcal{O}_K$, where $\mathcal{O}_K \subset K$ is the ring of integral numbers of an algebraic number field K (e.g. $K = \mathbb{Q}$ and $\mathcal{O}_K = \mathbb{Z}$).

But what is the physics behind the choice $S = \text{Spec } \mathcal{O}_K$? Why should we consider number fields instead of real or complex numbers? Following the ideas of B. Dragovich, V.S. Vladimirov, I.V. Volovich and many others (see also section 0.1), let us state at least two arguments at this place. The first argument concerns the process of measurement. While it is not clear at all whether transcendental numbers can be the result of a measurement, integral (or rational) numbers can. Second, we know from Einstein that gravity is encoded in deformations of space-time scales (described by means of the metrical tensor g). Looking at the energy scale that we experience, it is an empiric fact that we may assume that gravity is completely encoded in the archimedean scale and that non-archimedean, $\mathfrak{p}$ -adic scales may be neglected. Nevertheless, there is no reason why this should be true on all energy scales down to the Planck scale. It is an appealing project to study physical models where not only the ordinary, archimedean degrees of freedom are taken into consideration, but also the $\mathfrak{p}$ -adic, non-archimedean degrees of freedom. This was one starting point for the adelic models which we cited in section 0.1. Physically, the adelic approach means:

There is one degree of freedom per primespot and dimension. (*)

As already indicated in Example 0.1, the principle (*) may as well be realized by considering algebraic spaces over $\mathcal{O}_K$. This motivates the following Definition 0.4 (whose physical motivation will be illustrated in Remark 0.6). Recall that, given two relative algebraic spaces $X \rightarrow S$ and $Y \rightarrow S$, we denote by $X(Y)$ the set of S -morphisms $Y \rightarrow X$. Furthermore recall that for an algebraic space $\pi : X \rightarrow S$ we denote the fibre of π over the generic point of S by X_K (physically this generic fibre represents the archimedean component of the algebraic space).

0.4 Definition Let S be an excellent Dedekind scheme with field of fractions K of characteristic zero. Consider a pair $(X \rightarrow S, g)$ consisting of:

- a smooth, separated algebraic space $\pi : X \rightarrow S$ over S
- a metric g on X (see Definition 1.27)

such that the following conditions are fulfilled:

- (i) g satisfies the Einstein equations 1.35.
- (ii) For each smooth algebraic space $Y \rightarrow S$ and each K -morphism $u_K : Y_K \rightarrow X_K$ there is an S -morphism $u : Y \rightarrow X$ extending u_K .

Then the pair $(X \rightarrow S, g)$ is called a *model of type (GR)*.

0.5 Corollary *In the setting of Definition 0.4, let us assume that the algebraic space $\pi : X \rightarrow S$ is representable by a smooth and separated S -scheme. Then, the morphism u in Definition 0.4 b) is uniquely determined, i.e. $X \rightarrow S$ is the Néron model of its generic fibre X_K (see Definition 6.1). In particular, the following statements hold:*

- a) *If u_K is an isomorphism so is u .*
- b) *For each étale S -scheme S' with field of fractions K' the canonical map $X(S') \rightarrow X_K(K')$ is bijective.*

PROOF In order to prove the uniqueness assertion let us choose two morphisms u, v extending u_K . Using the separatedness of $X \rightarrow S$ we conclude from [Liu], Prop. 3.3.11, that u and v are equal if they coincide on a dense subset of Y . Therefore, it suffices to show that the generic fibre Y_K of Y is dense in Y . This may be done as follows: Due to smoothness, the structure morphism $f : Y \rightarrow S$ is an open map of topological spaces (use [BLR], Prop. 2.4/8 and [EGA IV₂], 2.4.6). The openness of f implies that the pre-image $f^{-1}(D)$ of any dense subset D of S is dense in Y . Therefore, we are done, because the generic point of S is dense in S . Consequently, $X \rightarrow S$ is the Néron model of its generic fibre.

The statements a) and b) follow directly from the universal property of Néron models. For example, choose $Y = S'$ in order to see b). $\square$

0.6 Remark If $S = \text{Spec } K$ is the spectrum of a field K , condition (ii) of Definition 0.4 is empty. If furthermore $K = \mathbb{R}$, any model of type (GR) induces a solution of Einstein's theory of general relativity (by evaluation at $\mathbb{R}$ -valued points). This explains the label *model of type (GR)*, because (GR) shall remind of general relativity. However, in the case $S = \text{Spec } \mathcal{O}_K$ we arrive at the following physical interpretation:

- Condition (ii) implements the “adelic” point of view.

In order to see this, let us choose $S = \text{Spec } \mathbb{Z}$ and therefore $K = \mathbb{Q}$. Recall that the generic fibre X_K of X represents the archimedean component. Then condition (ii) says that the archimedean world is only the projection from the “adelic” level to the archimedean component. In truth, everything is defined over all prime spots, and there is one degree of freedom per prime spot.

- We saw in Corollary 0.5 that condition (ii) implies a canonical bijection $X_K(K) = X(S)$. Recall that $X_K(K)$ is the set of archimedean points, and that $X(S)$ is the set of “adelic” points. In the special case $K = \mathbb{Q}$, the bijection $X_K(K) \cong X(S)$ means exactly that every archimedean point $x_\infty \in X_K(K)$ of X is in truth only the archimedean element x_∞ of an infinite set of points $x = \{x_2, \dots, x_p, \dots, x_\infty\} \in X(S)$. Finally, Corollary 0.5 a) reflects the physically crucial statement that any “deformation” of the archimedean component by means of isomorphisms extends to the “adelic” level.

Furthermore, we immediately obtain the interesting result that the pair $(X \rightarrow S, g)$ cannot be the flat Minkowski space-time if we are in the “adelic” situation $S = \text{Spec } \mathcal{O}_K$.

PROOF Let $S = \text{Spec } \mathcal{O}_K$ and assume that $(X \rightarrow S, g)$ describes the flat, topologically trivial Minkowski space-time. Then

- $g = \text{diag}(\pm 1, \pm 1, \pm 1, \pm 1)$ and
- $X = \mathbb{A}_S^n$ or $X = \mathbb{P}_S^n$ depending on whether we work projective or not.²

But if $X = \mathbb{A}_S^n$, then $K^n \cong X_K(K) \neq X(S) \cong \mathcal{O}_K^n$, and if $X = \mathbb{P}_S^n$, not every morphism u_K extends to an S -morphism u (see Example 5.4). Therefore, the flat, topologically trivial Minkowski space-time is impossible. $\square$

In chapter 3, we will study a particularly simple class of models of type (GR) which we call models of type (SR).

0.7 Definition Let $(X \rightarrow S, g)$ be a model of type (GR) in the sense of Definition 0.4, and let X_K be the generic fibre of X . Then the pair $(X \rightarrow S, g)$ is called a *model of type (SR)*, if in addition the following condition holds:

- (iii) X_K is a commutative K -group (see Definition 11.46).

More precisely, the K -group X_K should be considered as a K -torsor under X_K (see Definition 8.7). The latter means that the special choice of a zero element of the group is forgotten as it should be for physical reasons.

In order to generalize this notion slightly, one may also admit K -torsors X_K under K -groups $G_K \neq X_K$. However, we will restrict attention to the case $G_K = X_K$.

Definition 0.7 is motivated by *special relativity* with electromagnetism: The Minkowski space-time of special relativity naturally carries an additive, commutative group structure, and the gauge group of electromagnetism is commutative, too. This explains the

²Recall that the affine space $\mathbb{A}_S^n$ may be regarded as the algebraic geometric analogue of flat space. In order to see this, let $S = \text{Spec } R$ be the spectrum of a commutative ring R . Then $\mathbb{A}_S^n = \text{Spec } R[T_1, \dots, T_n]$ is the spectrum of a polynomial ring in n variables. Consequently, $\mathbb{A}_S^n(S) = \text{Hom}_R(R[T_1, \dots, T_n], R) = R^n$. In the special case $R = \mathbb{K}$, $\mathbb{K} = \mathbb{R}, \mathbb{C}$, the space-time induced by $\mathbb{A}_S^n$ is the flat manifold $\mathbb{A}_S^n(S) = \mathbb{K}^n$.

label *model of type (SR)*, because (SR) shall remind of special relativity. Due to the additional condition (iii), models of type (SR) are simpler than models of type (GR), and one can prove many properties of models of type (SR) in an abstract manner without fixing a special model. More precisely, we will see in chapter 3 that the following statements are true for *all* models of type (SR).

0.8 Properties of models of type (SR). Let $(X \rightarrow S, g)$ be a model of type (SR). Then the following statements are true:

- a) $X \rightarrow S$ is étale-invariant. More precisely, this statement means the following: Let $\varphi : X \rightarrow X$ be an étale S -morphism, and let $(X' \rightarrow S', g')$ be the pair obtained from $(X \rightarrow S, g)$ by base change with an étale morphism $S' \rightarrow S$. Then, $(X \rightarrow S, \varphi^*g)$ and $(X' \rightarrow S', g')$ are models of type (SR), too (see section 3.1).
- b) X cannot be the flat, topologically trivial Minkowski space (see section 3.2 for more details).
- c) The archimedean component $X_K(K)$ is bounded with respect to all $\mathfrak{p}$ -adic norms. In the special case $K = \mathbb{Q}$ and under the assumption that there is a closed immersion $X_K \hookrightarrow \mathbb{A}_K^n$, this is the following statement: For each prime number p , the p -adic manifold $X_K(\mathbb{Q}_p)$ is a bounded subset of some $\mathbb{Q}_p^n$ with respect to the canonical p -adic norm $|\cdot|_p$.

PROOF We know that X_K possesses a global Néron model. Consequently, the local Néron models exist (see Proposition 6.5). Therefore, due to Proposition 3.5, it is necessary that $X_K(K)$ or even the continuum $X_K(\hat{K})$ is bounded (see section 3.4 for details and the notion of bounded). $\square$

- d) The archimedean component $X_K(K)$ carries a discrete geometry. More precisely, Theorem 5.1 tells us that $X_K(K)$ is a finitely generated abelian group, i.e.

$$X_K(K) \cong \mathbb{Z}^d \oplus \mathbb{Z}/(p_1^{\nu_1}) \oplus \cdots \oplus \mathbb{Z}/(p_s^{\nu_s})$$

for some prime numbers $p_i \in \mathbb{N}$ and integers $d, s, \nu_i \in \mathbb{N}$. In the special case $d = 0$, $X_K(K)$ consists of only finitely many points. For a more detailed exposition we refer to chapter 5.

The finiteness of the set $X_K(K)$ would also be desirable in the case that $(X \rightarrow S, g)$ is a model of type (GR). However, it is unequally harder to see in how far this should be true. Nevertheless, motivated from what we already know about models of type (SR), it does not seem senseless to demand the finiteness of the set $X_K(K)$ in the case of models of type (GR).

- e) The privileged character of four-dimensional spaces: Let us consider pure gravity which is described by means of a euclidian metrical tensor $g_{\mu\nu}$. Over the field $\mathbb{R}$, the metric takes the form $\text{diag}(1, \dots, 1)$ in inertial systems. Interpreting the metric as a quadratic form $g : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^+$, we observe two fundamental properties:

- anisotropy: $g(v, v) = 0$ if and only if $v = 0$.
- surjectivity: For all positive $r \in \mathbb{R}$ there exists a $v \in \mathbb{R}^n$ such that $g(v, v) = r$.
Physically, this means that scales do not have holes.

We will see in section 4.1 that metrics over non-archimedean fields which fulfill the non-archimedean analogues of these two properties only exist in four dimensional spaces. Therefore, also in the case of number fields, we are naturally led to four dimensional spaces by means of the theorem of Hasse-Minkowski.

Additionally, in the case of number fields, we will see in section 4.1 that the archimedean tangent space $T_{X_K/K}(x_\infty)$ at an archimedean point $x_\infty \in X_K(K)$ naturally carries the structure of a quaternion algebra. This implies that we may write pairs $(\mathcal{A}, \overline{\mathcal{A}})$ consisting of a gauge field $\mathcal{A}$ and its anti-field $\overline{\mathcal{A}}$ in the form $(\mathcal{A}, \overline{\mathcal{A}}) = \sum_{\mu=0}^3 \mathcal{A}^\mu \gamma_\mu$ with gamma matrices fulfilling the relations

$$\gamma_\mu \gamma_\nu + \gamma_\nu \gamma_\mu = 2g_{\mu\nu}.$$

- f) If we do not demand the quasi-compactness of the archimedean component X_K of X , one can prove that X_K possesses a Néron model if and only if there is an exact sequence

$$0 \longrightarrow T_K \longrightarrow X_K \longrightarrow A_K \longrightarrow 0$$

over some algebraic closure of K , where T_K is an algebraic torus and A_K is an abelian variety (see section 3.3). While A_K is $\mathfrak{p}$ -adically bounded, T_K is not. This obstructs the interpretation of A_K as space part and the interpretation of the torus T_K as an internal, gauge group part which we therefore associate with electromagnetism. Thus, X_K should appear as A_K -torsor under T_K (which is the algebraic geometric analogue of the differential geometric principal bundle of gauge theory).

- g) On the “adelic” level there is some kind of entanglement of dimensions. For example, it is in general not possible to diagonalize the metric at the “adelic” points of $X(S)$ (see also Remark 3.26).

Let us remark that the statements a) and b) are also true for models of type (GR).

0.3. The continuum limit- Comparison with the adelic model

Within this section let $(X \rightarrow S, g)$ be a *model of type (GR)* (see Definition 0.4), and assume for simplicity that $S = \text{Spec } \mathbb{Z}$ and that $X \rightarrow S$ is representable by a smooth S -scheme. Then, on the fundamental level, space-time is given by the set $X(S)$. The metrical situation at a point $\alpha \in X(S)$ is described by the pull-back metric α^*g .

The purpose of this section is to answer the following questions:

- What is the archimedean, real continuum limit $\mathfrak{X}_\infty$ of $(X \rightarrow S, g)$?
- What are the non-archimedean continuum limits $\mathfrak{X}_p$ of $(X \rightarrow S, g)$?
- Does the family $\mathfrak{X}_\nu$, $\nu = 2, 3, 5, \dots, p, \dots, \infty$, give rise to an adelic continuum limit $\mathfrak{X}$ of $(X \rightarrow S, g)$ which is adelic in the sense of B. Dragovich, I.V. Volovich et. al.?

We will see that the last question has a positive answer. Therefore, our new approach via algebraic spaces presented within the bounds of this thesis fits exactly into the beautiful context of adelic, general relativistic (cosmological) models.

The archimedean continuum limit $\mathfrak{X}_\infty$

As usual, let X_K denote the generic fibre of X , and denote by g_K the pull-back of g under the canonical inclusion $X_K \hookrightarrow X$. Then, the archimedean limit is obtained by evaluation at K -valued points:

$$(X_K(K), g_K)$$

More precisely, the metric g_K at an arbitrary point $x \in X_K(K)$ is by construction the symmetric bilinear form $g_K(x) := x^* g_K : K \times K \rightarrow K$ (see the beginning of section 4.1). The continuum limit is obtained by applying the pull-back with $i : X_\mathbb{R} \rightarrow X_K$, where i is induced by the canonical map $K \hookrightarrow \mathbb{R}$. It is therefore the pair

$$(\mathfrak{X}_\infty, g_\infty) := (X_K(\mathbb{R}), i^* g_K).$$

By construction, $(\mathfrak{X}_\infty, g_\infty)$ may be endowed with the structure of a differentiable Riemannian manifold, and in addition it is a solution of Einstein's classical theory of general relativity.

As already indicated in 0.8, d), it is natural to assume that $X_K(K)$ is a finite set. Then, $X_K(K)$ is a discrete subset of the continuum $\mathfrak{X}_\infty$ and is invariant under arbitrary K -isomorphisms $\varphi_K : X_K \xrightarrow{\sim} X_K$. Furthermore, $\varphi_K^* g_K$ is a solution of the Einstein equations for all K -isomorphisms φ_K . Therefore, $(X_K(K), g_K)$ describes metrical states on smallest scales. At distances much larger than these smallest lengths, it is admissible to work with the continuum $(\mathfrak{X}_\infty, g_\infty)$ instead.

The non-archimedean continuum limits $\mathfrak{X}_p$

Let $s \in S$ be the Zariski closed point corresponding to a prime number p (i.e. more precisely corresponding to the prime ideal $(p) \subset \mathbb{Z}$). Let $\hat{R} := \hat{\mathcal{O}}_{S,s}$ denote the completion of $R := \mathcal{O}_{S,s}$ with respect to its maximal ideal s . In our special case, $\hat{R} = \mathbb{Z}_p$ is the ring of p -adic integers with field of fractions $\hat{K} = \mathbb{Q}_p$. By base change with the canonical morphism $\text{Spec } \hat{R} \rightarrow \text{Spec } R \rightarrow S$, X induces the $\hat{R}$ -scheme $X_{\hat{R}} := X \otimes_S \hat{R}$ which consists of a generic fibre and a special fibre (the latter is the fibre over the point s). In order to take into account all infinitesimal neighborhoods of the special fibre, let us

onsider the formal completion $\widehat{X}_{\widehat{R}}$ of $X_{\widehat{R}}$ along its special fibre. Let us point to the fact that $\widehat{X}_{\widehat{R}}$ coincides topologically with the special fibre of $X_{\widehat{R}}$. Furthermore, the metric g induces a metric g_p on $\widehat{X}_{\widehat{R}}$. Writing $\mathfrak{X}_p := \widehat{X}_{\widehat{R}}$ for the formal completion, we define the *non-archimedean continuum limit* of $(X \rightarrow S, g)$ over p as the pair

$$(\mathfrak{X}_p, g_p).$$

In order to illustrate the structure of the non-archimedean continuum limit, let us recall the notion of *formal completion of a scheme Y along a closed subscheme $Z \subset Y$* . So Z is defined by a quasi-coherent ideal $\mathcal{I} \subset \mathcal{O}_Y$. Then consider the sheaf $\mathcal{O}_Z$ obtained from restricting the projective limit $\varprojlim_n \mathcal{O}_Y/\mathcal{I}^n$ to Z . It follows that $(Z, \mathcal{O}_Z)$ is a locally ringed topological space, the desired formal completion $\widehat{Y}$ of Y along Z . Locally, the construction looks as follows: Let $Y = \text{Spec } A$ and assume that $\mathcal{I}$ is associated to the ideal $\mathfrak{i} \subset A$. Then, $(Z, \mathcal{O}_Z) = \text{Spf } \varprojlim_n A/\mathfrak{i}^n = \text{Spf } \hat{A}$, where $\hat{A}$ denotes the $\mathfrak{i}$ -adic completion of A .

The adelic continuum limit $\mathfrak{X}$

As illustrated above, the model of type (GR) $(X \rightarrow S, g)$ induces a family

$$\left(\begin{array}{c} \vdots \\ \vdots \end{array} \right) \quad \cdots \quad \left(\begin{array}{c} \vdots \\ \vdots \end{array} \right) \quad \cdots \quad \left(\begin{array}{c} \vdots \\ \vdots \end{array} \right)$$

$$(\mathfrak{X}_2, g_2) \quad \cdots \quad (\mathfrak{X}_p, g_p) \quad \cdots \quad (\mathfrak{X}_\infty, g_\infty)$$

where $(\mathfrak{X}_\infty, g_\infty)$ is a real manifold which is a solution of Einstein's ordinary theory of general relativity. Furthermore, each object $(\mathfrak{X}_p, g_p)$ lives (as topological space) over a finite prime spot and also solves the corresponding Einstein's equation.

We will illustrate now that this family canonically induces an adelic object $\mathfrak{X}$ which is the adelic continuum limit of $(X \rightarrow S, g)$ in the sense of B. Dragovich, I.V. Volovich et. al. For this purpose, consider the "generic fibre" $\mathfrak{X}_{p, \widehat{K}} := \widehat{X}_{\widehat{R}} := \widehat{X}_{\widehat{R}} \otimes_{\widehat{R}} \widehat{K}$ of $\mathfrak{X}_p = \widehat{X}_{\widehat{R}}$. This is a rigid-analytic space, and the metric g induces a metric $g_{p, \widehat{K}}$ on $\mathfrak{X}_{p, \widehat{K}}$ which is a solution of the p -adic Einstein equations. The pair

$$(\mathfrak{X}_{p, \widehat{K}}, g_{p, \widehat{K}}).$$

will be called the *p-adic continuum limit* of $(X \rightarrow S, g)$. Next, let us consider the direct

product

$$\mathfrak{X} := \left(\prod_p \mathfrak{X}_{p,\widehat{K}} \right) \times \mathfrak{X}_\infty.$$

We claim that $\mathfrak{X}$ is the desired adelic object. In order to illustrate this, let us fix some prime number p and let us study the structure of the p -adic continuum limits $\mathfrak{X}_{p,\widehat{K}}$ in more detail. Thereby, we make use of the notation which was already introduced in the previous subsection “The non-archimedean continuum limits $\mathfrak{X}_p$ ”.

First of all, consider the polynomial ring $A = \widehat{R}[\zeta]$ over $\widehat{R}$, where ζ denotes a set of variables $\zeta = (\zeta_1, \dots, \zeta_n)$, and where $\mathfrak{i} = (p)$ is the maximal ideal of $\widehat{R}$. So let Y be the n -dimensional affine space $\mathbb{A}_{\widehat{R}}^n$ over $\widehat{R}$, and let Z be its special fibre. Then the formal completion $\widehat{Y}$ of Y along Z yields the formal affine n -space $\mathrm{Spf} \widehat{R}[\zeta]$. Thereby, the $\widehat{R}$ -algebra $\widehat{R}\langle\zeta\rangle$ of restricted power series in the variables $\zeta_1, \dots, \zeta_n$ is defined as the subalgebra of the $\widehat{R}$ -algebra $\widehat{R}[[\zeta]]$ of formal power series, consisting of all series $\sum_{\nu \in \mathbb{N}^n} c_\nu \zeta^\nu$ with coefficients $c_\nu \in \widehat{R}$ constituting a zero sequence in $\widehat{R}$ (with respect to the p -adic topology on $\widehat{R}$). The “generic fibre” $\widehat{Y}_{\widehat{K}}$ of $\widehat{Y}$ is the affinoid $\widehat{K}$ -space $\mathrm{Sp}(\widehat{R}\langle\zeta\rangle \otimes_{\widehat{R}} \widehat{K}) = \mathrm{Sp} \widehat{K}\langle\zeta\rangle$ which (set-theoretically) coincides with the set of maximal ideals of $\widehat{K}\langle\zeta\rangle$. Consequently, $\widehat{Y}_{\widehat{K}} \subset Y_{\widehat{K}}^{\mathrm{an}} \cong Y(\widehat{K}) \cong \widehat{K}^n$ coincides with the unit ball $\mathbb{B}_{\widehat{K}}^n := \{x \in \widehat{K}^n \mid |x|_p \leq 1\}$ (where Y^{an} denote the rigid-analytification of $Y_{\widehat{K}} := Y \otimes_{\widehat{R}} \widehat{K}$). In the special case $n = 1$, $\widehat{Y}_{\widehat{K}}$ may actually be identified with the p -adic integers $\mathbb{Z}_p$.

Let us now return to our scheme $X_{\widehat{R}}$, but let us first assume that it is defined by an ideal $\mathfrak{a} \subset \widehat{R}[\zeta]$, where ζ denotes as above a set of variables $\zeta = (\zeta_1, \dots, \zeta_n)$. Then one can show that $\widehat{X}_{\widehat{R}} = \mathrm{Spf} \widehat{R}\langle\zeta\rangle / (\mathfrak{a})$. Thus, we find that $\mathfrak{X}_{p,\widehat{K}} = \mathrm{Sp} \widehat{K}\langle\zeta\rangle / (\mathfrak{a}) \subset \mathbb{B}_{\widehat{K}}^n$, i.e. set theoretically we may write:

$$\mathfrak{X}_{p,\widehat{K}} = X_K(\widehat{K}) \cap \mathbb{B}_{\widehat{K}}^n \subset \mathbb{Z}_p^n.$$

However, it is a general phenomenon that the generic fibre $\widehat{X}_{\widehat{K}} := \widehat{X}_{\widehat{R}} \otimes_{\widehat{R}} \widehat{K}$ should be viewed as an open subspace of the rigid analytification $X_{\widehat{K}}^{\mathrm{an}}$ of the scheme $X_{\widehat{K}} := X_{\widehat{R}} \otimes_{\widehat{R}} \widehat{K}$. Consequently, the direct product

$$\mathfrak{X} := \left(\prod_p \mathfrak{X}_{p,\widehat{K}} \right) \times \mathfrak{X}_\infty \subset \left(\prod_p \mathbb{Z}_p^n \right) \times \mathbb{R}^n \subset \mathbb{A}^n$$

is a subset of some power of the ring $\mathbb{A}_{\mathbb{Z}} := \widehat{\mathbb{Z}} \times \mathbb{R} \subset \mathbb{A}$ of adeles.

Finally, in the general case, we can find a covering of X by open subsets U_i , $i \in I$, such that there are S -immersions $U_i \hookrightarrow \mathbb{A}_S^n$ for all $i \in I$. Let $\mathfrak{X}_\nu$ (resp. $\mathfrak{U}_{i,\nu}$) be the respective continuum limits of X (resp. U_i), where $\nu = 2, 3, 5, \dots, p, \dots, \infty$. Let $\mathfrak{X} = (\prod_p \mathfrak{X}_{p,\widehat{K}}) \times \mathfrak{X}_\infty$ and $\mathfrak{U}_i = (\prod_p \mathfrak{U}_{i,p,\widehat{K}}) \times \mathfrak{U}_{i,\infty}$. By the above reasoning, $\mathfrak{U}_i$ is contained in an n -dimensional ring of adeles $\mathbb{A}^n$. Therefore, $\mathfrak{X}$ is obtained by gluing subsets of n -dimensional rings of adeles. Consequently, it is indeed some kind of “adelic manifold”. In particular, we find that the adelic space-time $\mathfrak{X}$ cannot be the flat, topologically trivial

adele $(\mathbb{A}_{\mathbb{Z}})^n$. This follows from the fact that on the one hand X cannot be the affine space $\mathbb{A}_S^n$ over S (see the statement below Remark 0.6), but that on the other hand $\mathfrak{X} = (\mathbb{A}_{\mathbb{Z}})^n$ if and only if $X = \mathbb{A}_S^n$.

All in all, this shows that our arithmetic geometric approach to general relativity completely fits into the well established and beautiful setting of adelic physics.

Part I.

Arithmetic theory of general relativity

1. Einstein's equation in the setting of algebraic spaces

1.1. Smoothness in algebraic geometry

In this section we will recapitulate some basic concepts of differential calculus for schemes. We introduce notions like unramified, étale and smooth morphisms and illustrate how these concepts generalize structures known from differential geometry. The aim is to perform general relativity in the setting of smooth algebraic spaces instead of smooth manifolds. The algebraic geometric Einstein equations will finally be derived in section 1.2. Using Grothendieck's language of schemes, our exposition on smoothness will follow [BLR], and we refer the reader to chapter 2 of this book for more details.

In the following let S be a base scheme and X an S -scheme (i.e. X is a scheme together with a unique morphism $X \rightarrow S$). Furthermore let R be a commutative ring with neutral element 1, and let A be an R -algebra. The reader, who is not so much familiar with the notion of global schemes, may for simplicity think of $X = \operatorname{Spec} A$ and $S = \operatorname{Spec} R$. As we are doing algebraic geometry, we have to perform differential calculus with purely algebraic methods. This is done by using so called derivations.

1.1 Definition a) An R -derivation of A into an A -module M is an R -linear map $d : A \rightarrow M$ such that

$$d(fg) = f \cdot d(g) + g \cdot d(f) \quad \text{for all } f, g \in A$$

The A -module of all R -derivations of A into an A -module M is denoted by $\operatorname{Der}_R(A, M)$.

b) The module of *relative differential forms (of degree one)* of A over R is an A -module $\Omega_{A/R}^1$ together with an R -derivation $d_{A/R} : A \rightarrow \Omega_{A/R}^1$ which has the following universal property: For each A -module M , the canonical map

$$\operatorname{Hom}_A(\Omega_{A/R}^1, M) \xrightarrow{\sim} \operatorname{Der}_R(A, M), \quad \varphi \mapsto \varphi \circ d_{A/R}$$

is bijective.

1.2 Remark $\Omega_{A/R}^1$ exists and its universal property implies:

a) $\Omega_{A/R}^1$ is unique up to canonical isomorphism.

b) Each morphism $\phi : A \rightarrow B$ of R -algebras induces a canonical B -linear map

$$\Omega_{A/R}^1 \otimes_A B \rightarrow \Omega_{B/R}^1, \quad f \cdot d_{A/R}(g) \otimes h \mapsto \phi(f) \cdot h \cdot d_{B/R}(\phi(g))$$

c) Considering A -derivations of B as R -derivations yields a B -linear map

$$\Omega_{B/R}^1 \rightarrow \Omega_{B/A}^1, \quad d_{B/R}(g) \mapsto d_{B/A}(g).$$

There is also a global notion of modules of differentials in terms of sheaves over schemes which has similar functorial properties. If X is locally of finite type over S there is a quasi-coherent $\mathcal{O}_X$ -module $\Omega_{X/S}^1$. It is called the *sheaf of relative differential forms (of degree 1)*. Furthermore we have a canonical $\mathcal{O}_X$ -linear map $d_{X/S} : \mathcal{O}_X \rightarrow \Omega_{X/S}^1$, the *exterior differential*.

Since $\Omega_{X/S}^1$ is quasi-coherent, $(\Omega_{X/S}^1, d_{X/S})$ can be described in local terms: For each open affine subset $V = \operatorname{Spec} R$ of S and for each open affine subset $U = \operatorname{Spec} A$ of X lying over V , the sheaf $\Omega_{X/S}^1|_U$ is the quasi-coherent $\mathcal{O}_X|_U$ -module associated to the A -module $\Omega_{A/R}^1$, and the map $d_{X/S}|_U$ is associated to the canonical map $d_{A/R} : A \rightarrow \Omega_{A/R}^1$.

An important operation, which we will frequently make use of, is the *pull-back* of differential forms.

1.3 Definition (pull-back of relative differential forms) Let $f : X \rightarrow Y$ be an S -morphism. Then the map in Remark 1.2, b) gives rise to a canonical $\mathcal{O}_X$ -morphism

$$f^* \Omega_{Y/S}^1 \rightarrow \Omega_{X/S}^1.$$

Each section ω of $\Omega_{Y/S}^1$ gives rise to a section ω' of $f^* \Omega_{Y/S}^1$ which is mapped to a section $f^* \omega$ of $\Omega_{X/S}^1$ under the above map. $f^* \omega$ is called the pull-back of ω .

Let us convince ourselves that this notion of pull-back reduces to the one given in differential geometry.

Physical interpretation: Let $S = \operatorname{Spec} \mathbb{R}$, $X = \mathbb{A}_{\mathbb{R}}^m$, $Y = \mathbb{A}_{\mathbb{R}}^n$ and let f be the map

$$f : \mathbb{A}_{\mathbb{R}}^m \rightarrow \mathbb{A}_{\mathbb{R}}^n, \quad T := (T_1, \dots, T_m)^t \mapsto p(T) := (p_1(T), \dots, p_n(T))^t,$$

where T denotes a set of variables, and where p_i , $i = 1, \dots, n$, are polynomials in T . Thus, on $\mathbb{R}$ -valued points $x \in X(\mathbb{R}) = \mathbb{R}^m$, f is simply the map

$$f(\mathbb{R}) : \mathbb{R}^m \rightarrow \mathbb{R}^n, \quad x := (x_1, \dots, x_m)^t \mapsto p(x) := (p_1(x), \dots, p_n(x))^t.$$

On its ring of global sections, f corresponds to the $\mathbb{R}$ -algebra homomorphism

$$\begin{aligned} f^* : \mathbb{R}[S_1, \dots, S_n] &\rightarrow \mathbb{R}[T_1, \dots, T_m] = \mathbb{R}[T]. \\ S_i &\mapsto p_i(T) \end{aligned}$$

Writing more compactly $S := (S_1, \dots, S_n)^t$ as a set of variables S_i , we can write f^* simply as:

$$f^*(S) = p(T).$$

This can be seen as follows: By Taylor expansion of p around a $\mathbb{R}$ -valued point $x \in X(\mathbb{R}) = \mathbb{R}^m$ we find a matrix $q(T) \in \text{Mat}_{\mathbb{R}[T]}(n \times m)$ such that

$$p(T) - p(x) = q(T) \cdot (T - x).$$

Therefore $(f^*)^{-1}((T - x) \cdot \mathbb{R}[T]) = (f^*)^{-1}((p(T) - p(x)) \cdot \mathbb{R}[T]) = (S - p(x)) \cdot \mathbb{R}[S]$, i.e. the $\mathbb{R}$ -valued point x is mapped by f to the $\mathbb{R}$ -valued point $p(x)$ as desired. Now going into the local description of the pull-back map, we find that we have the following canonical commutative diagram.

$$\begin{array}{ccc}
 f^*\Omega_{Y/S}^1 = f^*\Omega_{\mathbb{A}_{\mathbb{R}}^n}^1 & \xrightarrow{\quad} & \Omega_{\mathbb{A}_{\mathbb{R}}^m}^1 = \Omega_{X/S}^1 \\
 \parallel & & \parallel \\
 \Omega_{\mathbb{A}_{\mathbb{R}}^n}^1 \otimes_{\mathbb{R}[S]} \mathbb{R}[T] & \xrightarrow{\quad} & \Omega_{\mathbb{A}_{\mathbb{R}}^m}^1 \\
 \parallel & & \parallel \\
 (\oplus_{i=1}^n \mathbb{R}[S] \cdot dS_i) \otimes_{\mathbb{R}[S]} \mathbb{R}[T] & \xrightarrow{\quad} & \oplus_{i=1}^m \mathbb{R}[T] \cdot dT_i \\
 \parallel & & \parallel \\
 \oplus_{i=1}^n \mathbb{R}[T] \cdot dS_i & \xrightarrow{\quad} & \oplus_{i=1}^m \mathbb{R}[T] \cdot dT_i, \quad dS_i \longmapsto \sum_{j=1}^m \frac{\partial p_i}{\partial T_j} \cdot dT_j
 \end{array}$$

Thus the pull-back of a differential form $\omega = h(S) \cdot dS \in \Omega_{Y/S}^1 = \Omega_{\mathbb{A}_{\mathbb{R}}^n}^1$ is

$$\begin{aligned}
 f^*\omega &= h(f^*(S)) \cdot f^*(dS) = h(f^*(S)) \cdot d(f^*(S)) = h(p(T)) \cdot d(p(T)) \\
 &= h(p(T)) \cdot \left[\left(\frac{\partial p_i}{\partial T_j} \right) \cdot dT_j \right],
 \end{aligned}$$

and on $\mathbb{R}$ -valued points $x \in X(\mathbb{R}) = \mathbb{R}^m$ we get

$$(f^*(h(S)dS))(x) = (h \circ p)(x) \cdot [(Dp)(x) \cdot dx],$$

where Dp denotes the Jacobian matrix of $p = f(\mathbb{R}) : \mathbb{R}^m \rightarrow \mathbb{R}^n$. This is just the notion of pull-back of differential forms we are used to from differential geometry.

There are some exact sequences which are induced by the maps defined in Remark 1.2. As we will make use of them later let us state them here. For proofs we refer to [EGA IV₄], 16.4.

1.4 Proposition *Let $f : X \rightarrow Y$ be an S -morphism. Then the canonical sequence of $\mathcal{O}_X$ -modules*

$$f^*\Omega_{Y/S}^1 \longrightarrow \Omega_{X/S}^1 \longrightarrow \Omega_{X/Y}^1 \longrightarrow 0$$

is exact.

1.5 Proposition *Let $j : Z \hookrightarrow X$ be an immersion of S -schemes. Let $\mathcal{J}$ be the sheaf of ideals defining Z as subscheme of X . Then the canonical sequence of $\mathcal{O}_Z$ -modules*

$$\mathcal{J} / \mathcal{J}^2 \xrightarrow{\delta} j^*\Omega_{X/S}^1 \longrightarrow \Omega_{Z/S}^1 \longrightarrow 0$$

is exact, where δ is locally given by the map $\bar{a} \mapsto d(a) \otimes 1$, $\bar{a} \equiv a \pmod{\mathcal{J}^2}$.

1.6 Proposition *Let X and S' be S -schemes. Let $X' := X \times_S S'$ be the S -scheme obtained by base change, and let $p : X' \rightarrow X$ be the projection. Then the canonical map*

$$p^*\Omega_{X/S}^1 \longrightarrow \Omega_{X'/S'}^1$$

is an isomorphism.

1.7 Proposition *Let X_1 and X_2 be S -schemes. If $p_i : X_1 \times_S X_2 \rightarrow X_i$ are the projections for $i = 1, 2$, the canonical map*

$$p_1^*\Omega_{X_1/S}^1 \oplus p_2^*\Omega_{X_2/S}^1 \xrightarrow{\sim} \Omega_{X_1 \times_S X_2/S}^1$$

is an isomorphism.

Let us now explain the concepts of unramified, étale and smooth morphisms. These morphisms will provide us the analogues of typical constructions from differential geometry like submersions and immersions.

1.8 Definition A morphism of schemes $f : X \rightarrow S$ is called *unramified* at a point $x \in X$ if there exist an open neighborhood U of x and an S -immersion

$$j : U \hookrightarrow \mathbb{A}_S^n$$

of U into some linear space $\mathbb{A}_S^n$ over S such that the following conditions are satisfied:

- a) locally at $j(x)$ (i.e., in an open neighborhood of $j(x)$), the sheaf of ideals $\mathcal{J}$ defining $j(U)$ as a subscheme of $\mathbb{A}_S^n$ is generated by finitely many sections.
- b) the differential forms of type dg with sections g of $\mathcal{J}$ generate $\Omega_{\mathbb{A}_S^n/S}^1$ at $j(x)$.

The morphism $f : X \rightarrow S$ is called unramified if it is unramified at all points of X .

1.9 Proposition *Let $f : X \rightarrow S$ be locally of finite presentation, let $x \in X$ and $s := f(x)$. Then the following conditions are equivalent:*

- a) f is unramified at x .
- b) $\Omega_{X/S,x}^1 = 0$.
- c) The diagonal morphism $\Delta : X \rightarrow X \times_S X$ is a local isomorphism at x .
- d) The maximal ideal $\mathfrak{m}_x$ of $\mathcal{O}_{X,x}$ is generated by the maximal ideal $\mathfrak{m}_s$ of $\mathcal{O}_{S,s}$ and $k(x)$ is a finite separable extension of $k(s)$.

The morphism $f : X \rightarrow S$ is called *unramified* if it is unramified at all points of X .

Before defining the central notion of smooth morphisms, let us make some remarks on the notion of dimensionality in the realm of schemes. Let $f : X \rightarrow S$ be a morphism of schemes. Then the relative dimension $\dim_x f := \dim_x f^{-1}(f(x)) \equiv \dim_x X_{f(x)}$ of f at a point $x \in X$ is in general defined via lengths of chains of irreducible closed subsets of the fibre $X_{f(x)}$. This is explained in detail in [Liu], Chapter 2.5. In the case of S -immersions $X \hookrightarrow \mathbb{A}_S^n$ (in which we are interested), $X_{f(x)}$ is of finite type over the field $k(f(x))$ and one can show that $\dim_x X_{f(x)}$ coincides with the intuitively clear notion of relative dimension given in the following definition of smooth morphisms.

1.10 Definition A morphism of schemes $f : X \rightarrow S$ is called *smooth* at a point $x \in X$ (of relative dimension r) if there exist an open neighborhood U of x and an S -immersion

$$j : U \hookrightarrow \mathbb{A}_S^n$$

of U into some linear space $\mathbb{A}_S^n$ over S such that the following conditions are satisfied:

- a) locally at $y := j(x)$ (i.e., in an open neighborhood of $j(x)$), the sheaf of ideals $\mathcal{J}$ defining $j(U)$ as a subscheme of $\mathbb{A}_S^n$ is generated by $(n - r)$ sections $g_{r+1}, \dots, g_n$.
- b) the differential forms $dg_{r+1}(y), \dots, dg_n(y)$ are linearly independent in $\Omega_{\mathbb{A}_S^n/S}^1 \otimes k(y)$.

The morphism $f : X \rightarrow S$ is called *smooth* if it is smooth at all points of X .

Physical interpretation: Let $S = \text{Spec } \mathbb{R}$ and let $X = V(g_{r+1}, \dots, g_n) \subset \mathbb{A}_S^n$ be the zero set of $(n - r)$ polynomials $g_{r+1}, \dots, g_n \in \mathbb{R}[T_1, \dots, T_n]$ such that the canonical map $f : X \rightarrow S$ is smooth of relative dimension r . Therefore

$$X(\mathbb{R}) = \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid g_i(x_1, \dots, x_n) = 0 \text{ for all } r+1 \leq i \leq n\}$$

is a r -dimensional differentiable manifold. $X(\mathbb{R}) \hookrightarrow \mathbb{A}_S^n(\mathbb{R}) = \mathbb{R}^n$ is an immersion in the sense of ordinary differential geometry.

Thus the notion of smooth morphisms is a natural generalization of immersions in the sense of ordinary differential geometry and analogously unramified morphisms generalize submersions.

1.11 Definition A morphism of schemes $f : X \rightarrow S$ is called *étale* (at a point) if it is smooth (at the point) of relative dimension 0.

1.12 Corollary *An immersion $f : X \rightarrow S$ is étale if and only if f is an open immersion.*

PROOF [BLR], Lemma 2.2/4 □

1.13 Proposition *Let $f : X \rightarrow S$ be a smooth morphism of schemes. Then:*

- a) $\Omega_{X/Y}$ is locally free. Its rank at $x \in X$ is equal to the relative dimension of f at x .
- b) The canonical sequence of $\mathcal{O}_X$ -modules

$$0 \longrightarrow f^* \Omega_{Y/S}^1 \longrightarrow \Omega_{X/S}^1 \longrightarrow \Omega_{X/Y}^1 \longrightarrow 0$$

is exact and locally split.

PROOF [BLR], Prop. 2.2/5 □

1.14 Proposition *Let $f : X \rightarrow S$ be locally of finite presentation. Let $x \in X$ and set $s := f(x)$. The following conditions are equivalent:*

- a) f is smooth at x .
- b) f is flat at x and the fibre $X_s = X \times_S \operatorname{Spec} k(s)$ is smooth over $k(s)$ at x .

PROOF [BLR], Prop. 2.4/8 □

Like in differential geometry there is a Jacobi Criterion for smoothness. The analogy to the classical Jacobi Criterion is particularly evident in characterization e) of the following list of criterions for smoothness.

1.15 Jacobi Criterion Let X and Z be S -schemes and let $j : X \hookrightarrow Z$ be a closed immersion which is locally of finite presentation. Let $\mathcal{J} \subset \mathcal{O}_Z$ be the sheaf of ideals which defines X as a subscheme of Z . Let $x \in X$ and $z := j(x)$. Assume that, as an S -scheme, Z is smooth at z of relative dimension n . Then the following conditions are equivalent:

- a) As an S -scheme, X is smooth at x of relative dimension r .
- b) The canonical sequence of $\mathcal{O}_X$ -modules

$$0 \longrightarrow \mathcal{J} / \mathcal{J}^2 \longrightarrow j^* \Omega_{Z/S}^1 \longrightarrow \Omega_{X/S}^1 \longrightarrow 0$$

is split exact at x , and $r = \operatorname{rank}(\Omega_{X/S}^1 \otimes k(x))$.

- c) If $dz_1, \dots, dz_n$ is a basis of $(\Omega_{Z/S}^1)_z$, and if $g_1, \dots, g_N$ are local sections of $\mathcal{O}_Z$ generating $\mathcal{J}_z$, there exists a re-indexing of the $z_1, \dots, z_n$ and of the $g_1, \dots, g_N$ such that $g_{r+1}, \dots, g_N$ generate $\mathcal{J}$ at z and such that $dz_1, \dots, dz_r, dg_{r+1}, \dots, dg_N$ generate $(\Omega_{X/S}^1)_z$.

d) There exist local sections $g_{r+1}, \dots, g_n$ of $\mathcal{O}_Z$ such that

- (i) $g_{r+1}, \dots, g_n$ generate $\mathcal{J}_z$.
- (ii) $dg_{r+1}(z), \dots, dg_n(z)$ are linearly independent in $\Omega_{Z/S}^1 \otimes_{\mathcal{O}_Z} k(z)$, where $dg_i(z)$ is the image of $dg_i \in (\Omega_{Z/S}^1)_z$ in $\Omega_{Z/S}^1 \otimes_{\mathcal{O}_Z} k(z) = (\Omega_{Z/S}^1)_z / \mathfrak{m}_z(\Omega_{Z/S}^1)_z$.

e) There exist local sections $g_{r+1}, \dots, g_n$ of $\mathcal{O}_Z$ such that

- (i) $g_{r+1}, \dots, g_n$ generate $\mathcal{J}_z$.
- (ii) considering a representation

$$dg_j = \sum_{i=1}^n \frac{\partial g_j}{\partial z_i} dz_i$$

of the differential forms $g_{r+1}, \dots, g_n$ with respect to a basis $dz_1, \dots, dz_n$ of $(\Omega_{Z/S}^1)_z$, there is a $(n-r)$ -minor of the matrix $\left(\frac{\partial g_j}{\partial z_i}\right)$ which does not vanish at z .

PROOF [BLR], Prop. 2.2/7 □

1.16 Proposition *Let $f : X \rightarrow Y$ be an S -morphism. Let $x \in X$ and set $y := f(x)$. Assume that X is smooth over S at x and that Y is smooth over S at y . Then the following conditions are equivalent:*

- a) f is étale at x .
- b) The canonical homomorphism $(f^* \Omega_{Y/S}^1)_x \rightarrow (\Omega_{X/S}^1)_x$ is bijective.

PROOF [BLR], Cor. 2.2/10 □

1.17 Proposition *Let $f : X \rightarrow S$ be a morphism and $x \in X$. Then the following conditions are equivalent:*

- a) f is smooth of relative dimension n .
- b) There exists an open neighborhood U of x and a commutative diagram

$$\begin{array}{ccc} U & \xrightarrow{g} & \mathbb{A}_S^n \\ & \searrow f|_U & \downarrow p \\ & & S \end{array}$$

where g is étale and p is the canonical projection.

PROOF [BLR], Prop. 2.2/11 □

1.18 Proposition *If X is a smooth scheme over a field k , the set of closed points x of X such that $k(x)$ is a separable extension of k is dense in X .*

PROOF [BLR], Cor. 2.2/13 □

1.19 Proposition *Let $f : X \rightarrow S$ be a smooth morphism. Let s be a point of S , and let x be a closed point of the fibre $X_s = X \times_S \operatorname{Spec} k(s)$ such that $k(x)$ is a separable extension of $k(s)$. Then there exists an étale morphism $g : S' \rightarrow S$ and a point $s' \in S'$ above s such that the morphism $f' : X \times_S S' \rightarrow S'$ obtained from f by the base change $S' \rightarrow S$ admits a section $h : S' \rightarrow X \times_S S'$, where $h(s')$ lies above x , and where $k(s') = k(x)$.*

PROOF [BLR], Prop. 2.2/14 □

1.20 Proposition *Let R be a local henselian ring with residue field k . Let X be a smooth R -scheme. Then the canonical map $X(R) \rightarrow X(k)$ from the set of R -valued points of X to the set of k -valued points of X is surjective. In particular, if R is strictly henselian, the set of k -valued points of $X_k = X \otimes_R k$ which lift to R -valued points of X is dense in X_k .*

PROOF [BLR], Prop. 2.3/5 □

Let us finish with some remarks on flatness, because by Proposition 1.14 smooth morphisms are in particular flat.

1.21 Proposition *Let $f : X \rightarrow Y$ be locally of finite presentation. If f is flat, then f is open.*

PROOF [EGA IV₂], 2.4.6 □

1.22 Proposition *Let $f : X \rightarrow Y$ be faithfully flat (i.e. flat and surjective) and quasi-compact. Then the topology of Y is the quotient topology of X with respect to f , i.e. a subset $V \subset Y$ is open if and only if $f^{-1}(V)$ is open in X .*

PROOF [EGA IV₂], 2.3.12 □

Later, we will describe the physical models by means of smooth morphisms $f : X \rightarrow S$ of algebraic spaces, where S is an Zariski one-dimensional excellent Dedekind ring. This choice of S and the flatness of f guarantee that the considered models have got the properties that one would expect. Next to the fact that the relative dimension of the universe X is constant (see Lemma 1.25), X may be considered as a “continuous” family of fibres X_s , $s \in S$; i.e. if we remove one fibre $X_s \subset X$ over a closed point $s \in S$, we can fill up this hole uniquely from the data given on the open subscheme $X \setminus X_s \subset X$.

1.23 Proposition *Let Y be a regular, integral scheme of dimension one, let $p \in Y$ be a closed point, and let $X \hookrightarrow \mathbb{P}_{Y-\{p\}}^n$ be a closed subscheme which is flat over $Y - \{p\}$. Then there exists a unique closed subscheme $\bar{X} \hookrightarrow \mathbb{P}_Y^n$, flat over Y , whose restriction to $\mathbb{P}_{Y-\{p\}}^n$ is X .*

PROOF [Har], Chap. III, Prop. 9.8 □

1.24 Proposition *Let $f : X \rightarrow Y$ be a morphism of schemes, with Y integral and regular of dimension one. Then f is flat if and only if every associated point $x \in X$ maps to the generic point of Y . In particular, if X is reduced, this says that every irreducible component of X dominates Y .*

PROOF [Har], Chap. III, Prop. 9.7 □

Finally, let us conclude with the statements that illustrate the very nice behavior of the relative dimension of smooth morphisms. Thereby, the flatness of smooth morphisms is essential.

1.25 Lemma *Let $f : X \rightarrow S$ be locally of finite type and flat. Assume that X is irreducible and that S is locally noetherian. Then the relative dimension of f is constant on X .*

PROOF [EGA IV₂], 14.2.2 □

1.26 Lemma *Let S be locally noetherian scheme and let $f : X \rightarrow S$ be a morphism of finite type smooth at a point x . Let $s := f(x)$. Then $\Omega_{X/S}^1$ is free of rank $\dim_x X_s$ in a neighborhood of x .*

PROOF [Liu], Prop. 6.2.5 □

1.2. The arithmetic Einstein equations

Let $X \rightarrow S$ be a smooth, separated S -scheme of relative dimension n . The purpose of this section is the derivation of the fundamental equations of general relativity in our algebraic geometric setting. As the ordinary differential geometric Einstein equations are differential equation, we must expect that this holds in algebraic geometry, too. The necessary techniques concerning smoothness and differential calculus in algebraic geometry were summarized in section 1.1. Crucial are the following notions.

$\Omega_{X/S}^1$ sheaf of (relative) differential forms

$\mathcal{T}_{X/S} := \mathcal{H}om_{\mathcal{O}_X}(\Omega_{X/S}^1, \mathcal{O}_X)$ sheaf of (relative) vector fields (see Definition 11.23)

$T_{X/S} := \mathbb{V}(\Omega_{X/S}^1)$ (relative) tangent bundle (see Definition 11.24)

One can prove (Remark 11.25) that

$$\Gamma(T_{X/S}/U) := \text{Hom}_X(U, T_{X/S}) \cong \mathcal{T}_{X/S}(U)$$

for every Zariski open subset $U \subset X$. Therefore vector fields correspond to sections of the tangent bundle.

The metric tensor

Due to smoothness, the sheaves $\Omega_{X/S}^1$ and $\mathcal{T}_{X/S}$ are locally free. Let us fix a local base $\{\omega^i\}$ of $\Omega_{X/S}^1$ which is dual to the local base $\{\partial_i\}$ of $\mathcal{T}_{X/S}$.

1.27 Definition Let $g : \mathcal{T}_{X/S} \times_X \mathcal{T}_{X/S} \rightarrow \mathbb{A}_X^1$ be an X -morphism which is bilinear (see section 4.3). Equivalently, g may be interpreted as a global section of $\Omega_{X/S}^{\otimes 2}$. Locally, we may write

$$g = \sum_{1 \leq i, j \leq n} g_{ij} \omega^i \otimes \omega^j \in \Omega_{X/S}^{\otimes 2}, \quad g_{ij} \in \mathcal{O}_X.$$

Then g is called a *metric* if the following conditions hold for any sufficiently small open subset of X :

- (i) The matrix (g_{ij}) is symmetric, i.e. $g_{ij} = g_{ji}$.
- (ii) The matrix (g_{ij}) is invertible, i.e. $\det(g_{ij}) \in \mathcal{O}_X^*$.

Covariant derivation

1.28 Definition Let $\nabla : \mathcal{T}_{X/S} \times_X \mathcal{T}_{X/S} \rightarrow \mathcal{T}_{X/S}$ be an X -morphism. Interpret ∇ as a map

$$\nabla : \mathcal{T}_{X/S}(X) \times \mathcal{T}_{X/S}(X) \rightarrow \mathcal{T}_{X/S}(X), \quad (\mathbf{u}, \mathbf{v}) \mapsto \nabla_{\mathbf{u}} \mathbf{v}.$$

Let us assume that ∇ is a $\mathcal{O}_S(S)$ -bilinear map, where the $\mathcal{O}_X(X)$ -module $\mathcal{T}_{X/S}(X)$ is viewed as $\mathcal{O}_S(S)$ -module via the canonical morphism $\mathcal{O}_S(S) \rightarrow \mathcal{O}_X(X)$. Then ∇ is called a *covariant derivation* if the following conditions hold for all $f \in \mathcal{O}_X(X)$ and $\mathbf{u}, \mathbf{v} \in \mathcal{T}_{X/S}(X)$:

- (i) $\nabla_{f\mathbf{u}} \mathbf{v} = f \nabla_{\mathbf{u}} \mathbf{v}$.
- (ii) $\nabla_{\mathbf{u}}(f\mathbf{v}) = (\mathbf{u}f)\mathbf{v} + f \nabla_{\mathbf{u}} \mathbf{v}$.

Thereby, $\mathbf{u}f := \mathbf{u}(f) = (d_{X/S}f)(\mathbf{u})$ is the canonical action of vector fields on functions (the differential $d_{X/S}$ is introduced and explained directly above Definition 1.3).

1.29 Definition Let ∇ be a covariant derivation, and let $\mathbf{u}, \mathbf{v}$ and $\mathbf{w} \in \mathcal{T}_{X/S}(X)$.

- a) $T(\mathbf{u}, \mathbf{v}) := \nabla_{\mathbf{u}} \mathbf{v} - \nabla_{\mathbf{v}} \mathbf{u} - [\mathbf{u}, \mathbf{v}]$ is called the *torsion* of ∇ .
- b) ∇ is called *torsion-free* if and only if $T(\mathbf{u}, \mathbf{v}) = 0$ for all $\mathbf{u}, \mathbf{v}$.
- c) ∇ is called *metrical* if and only if $\mathbf{u}g(\mathbf{v}, \mathbf{w}) = g(\nabla_{\mathbf{u}} \mathbf{v}, \mathbf{w}) + g(\mathbf{u}, \nabla_{\mathbf{u}} \mathbf{w})$ for all $\mathbf{u}, \mathbf{v}, \mathbf{w}$.

In the same way as in differential geometry one proves that there exists a uniquely determined covariant derivation ∇ which is metrical and torsion-free, the *Levi-Civita connection*. The Levi-Civita connection is completely determined by the metrical tensor. More precisely, the Koszul formula holds.

$$2g(\nabla_{\mathbf{u}} \mathbf{v}, \mathbf{w}) = \mathbf{u}g(\mathbf{v}, \mathbf{w}) - \mathbf{w}g(\mathbf{u}, \mathbf{v}) + \mathbf{v}g(\mathbf{w}, \mathbf{u}) + g([\mathbf{u}, \mathbf{v}], \mathbf{w}) + g([\mathbf{w}, \mathbf{u}], \mathbf{v}) - g([\mathbf{v}, \mathbf{w}], \mathbf{u})$$

Curvature

From now on let ∇ be the Levi-Civita connection. Then we may introduce the curvature tensor

$$R_{uv}(\mathfrak{w}) := \nabla_u \nabla_v \mathfrak{w} - \nabla_v \nabla_u \mathfrak{w} - \nabla_{[u, v]} \mathfrak{w}.$$

Then the tensor

$$R_{\mathfrak{z}uvw} := g(R_{uv}(\mathfrak{w}), \mathfrak{z})$$

is called the *Riemannian curvature tensor*. The Riemannian curvature tensor fulfills the following identities.

1.30 Proposition *Let $u, v, w, z \in \mathcal{T}_{X/S}(X)$. Then:*

- a) $R_{uvwz} = -R_{vuwz}$
- b) $R_{uvwz} = -R_{uvz w}$
- c) $R_{uvwz} = R_{wzuv}$
- d) first Bianchi-identity: $R_{zuvw} + R_{v w u z} + R_{w u v z} = 0$
- e) second Bianchi-identity: $(\nabla_u R)_{vw} + (\nabla_v R)_{wu} + (\nabla_w R)_{uv} = 0$
Thereby, $(\nabla_u R)_{vw}(z) := \nabla_u(R_{vw}(z)) - R_{\nabla_u v, w}(z) - R_{v, \nabla_u w}(z) - R_{vw}(\nabla_u z)$.

More generally, the covariant derivation of arbitrary tensor fields S and T with respect to a vector field v may be defined inductively as follows: $\nabla_v(S \otimes T) := \nabla_v S \otimes T + \nabla_v T \otimes S$.

1.31 Definition The bi-quadratic form

$$k(u, v) := R_{uvuv}$$

is called intersection curvature.

1.32 Proposition *The Riemannian curvature tensor is completely determined by k . More precisely:*

- a) $4 \cdot R_{uvvw} = k(u + w, v) - k(u - w, v)$
- b) $6 \cdot R_{uvwz} = R_{u, v+w, v+w, z} - R_{u, v-w, v-w, z} - R_{v, u+w, u+w, z} + R_{v, u-w, u-w, z}$

1.33 Corollary *Let $X \rightarrow S$ be a smooth S -scheme with metric g . Then X is flat, i.e. $k(u, v) = 0$ for all $u, v \in \mathcal{T}_{X/S}(X)$, if and only if the Riemannian curvature tensor vanishes, i.e. $R = 0$.*

1.34 Definition Let $X \rightarrow S$ be a smooth S -scheme with metric g and consider a local base $\{\partial_i\}$ of $\mathcal{T}_{X/S}$ and a local base $\{\omega^i\}$ of $\Omega_{X/S}^1$. The curvature tensor $R_{\mathbf{u}\mathbf{v}}(\mathbf{w})$ is trilinear in $\mathbf{u}, \mathbf{v}, \mathbf{w}$ and therefore induces linear maps $R_{\bullet\mathbf{v}}(\mathbf{w})$. Taking the trace finally yields the symmetric bilinear form ric

$$\text{ric}(\mathbf{v}, \mathbf{w}) := \text{Tr}(R_{\bullet\mathbf{v}}(\mathbf{w}))$$

which is called the *Ricci-form* of $(X \rightarrow S, g)$. Now consider the uniquely determined tensor Ric which is given by $g(\text{Ric}(\mathbf{u}), \mathbf{v}) = \text{ric}(\mathbf{u}, \mathbf{v})$ for all vector-fields $\mathbf{u}, \mathbf{v}$. The *scalar curvature* sc is by definition the trace

$$\text{sc} := \text{Tr}(\text{Ric}(\bullet))$$

of the linear map $\text{Ric}(\bullet)$. Furthermore, the *divergence* $\text{div}(T)$ of any symmetric $(0, q)$ -tensor $T := \sum T_{i_1 \dots i_q} \omega^{i_1} \otimes \dots \otimes \omega^{i_q}$ is defined as follows: The covariant derivation ∇T of T is a $(0, q+1)$ -tensor $\nabla T := \sum T_{i_1 \dots i_q; j} \omega^{i_1} \otimes \dots \otimes \omega^{i_q} \otimes \omega^j$. Then $\text{div}(T)$ is the $(0, q-1)$ -tensor which is obtained by lifting the new variable and contracting it:

$$(\text{div}(T))_{i_1 \dots i_{q-1}} = g^{i_q j} T_{i_1 \dots i_q; j}.$$

Einstein's equation

Let $X \rightarrow S$ be a smooth S -scheme with metric g , and let ∇ be the Levi-Civita connection on X . Furthermore, let T denote the energy-stress tensor. This is a symmetric $(0, 2)$ -tensor on X with $\text{div}(T) = 0$. Then the equations of general relativity in our arithmetic setting are given by the following system of equations:

1.35 Einstein equations

$$\text{ric} - \frac{1}{2} \text{sc} \cdot g = \kappa T$$

where $\kappa \in \mathcal{O}_S(S)$ is a constant. Now, having written down the equations of general relativity in the setting of arithmetic algebraic geometry, one can ask for solutions. Choosing $S = \text{Spec } \mathbb{R}$ and assuming that there exists a solution of the corresponding algebraic geometric Einstein equations, it follows that this solution gives rise to a differential geometric solution of the ordinary, differential geometric Einstein equations. This follows from the purely algebraic nature of the notions metric, covariant derivation and curvature (see also section 4.3 for more details). However, we are interested in “adelic” space-time models. So, let us now make the choice $S = \text{Spec } \mathcal{O}_K$, where $\mathcal{O}_K$ denotes as usual the ring of integral numbers of an algebraic number field K . Recall that the generic fibre X_K of $X \rightarrow S$ represents the archimedean limit.

In order to solve the Einstein equations, it is most convenient to perform all computations locally and to glue the local solutions in a second step. We will see that these local computations may be performed in essentially the same way as in differential geometry. On the one hand, this is due to the fact that the local ring $\mathcal{O}_{X,x}$ at a point $x \in X_K(K)$ may be embedded into a ring of formal power series.

1.36 Proposition *Let $X \rightarrow S$ be a smooth morphism of locally Noetherian schemes. Let $s \in S$ and $x \in X_s$ be a $k(s)$ -rational point. Then there exists an isomorphism of $\widehat{\mathcal{O}}_{S,s}$ -algebras*

$$\widehat{\mathcal{O}}_{X,x} = \widehat{\mathcal{O}}_{S,s}[[x_1, \dots, x_n]]$$

where $(x_1, \dots, x_n)$ is a set of variables and $n = \dim \mathcal{O}_{X_s, x}$.

PROOF [Liu], Ex. 6.3.1 □

Furthermore, we know from Proposition 1.17 that each point $x \in X$ possesses an open environment U which is étale over some affine space $\mathbb{A}_S^n = \operatorname{Spec} \mathcal{O}_S[x_1, \dots, x_n]$. Therefore, the module $\Omega_{U/S}^1$ of differential forms over U is the free $\mathcal{O}_U$ -module generated by the differentials $dx_1, \dots, dx_n$ (see Proposition 3.1), and we may choose the base $\{\omega^i := dx_i\}$ of $\Omega_{U/S}^1$ together with the corresponding dual base $\{\partial_i\}$ of $\mathcal{T}_{U/S}$.

The functions $f \in \mathcal{O}_U$ on U are algebraic over the polynomial ring $\mathcal{O}_S[x_1, \dots, x_n]$, because we may assume that U is standard étale over $\mathbb{A}_S^n$ (see [BLR], Prop. 2.3/3). Consequently, there is a canonical differential calculus on U with respect to the coordinates $x_1, \dots, x_n$. More precisely, the vector field ∂_i acts on f by means of ordinary partial derivation with respect to the i -th coordinate x_i . This may be seen as follows: On polynomials we have clearly $\partial_i x_j^n = \delta_{ij} \cdot n x_j^{n-1}$ due to the Leibniz rule (see Definition 1.1, a)). If $f \in \mathcal{O}_U$ is arbitrary, there is an algebraic equation $\sum_{j=0}^m c_j f^j = 0$ with polynomials $c_j \in \mathcal{O}_S[x_1, \dots, x_n]$, $c_m \neq 0$. It follows that $0 = \partial_i(\sum_{j=0}^m c_j f^j) = \sum_{j=0}^m (f^j \partial_i c_j + c_j j f^{j-1} \partial_i f)$ which is a linear equation in $\partial_i f$ and thus may be solved uniquely for $\partial_i f$ on the locus where $\sum_{j=0}^m c_j j f^{j-1} \neq 0$. However, by what we already know, $\partial_i c_j$ is the ordinary partial derivation of c_j with respect to the i -th coordinate x_i , and so we are done.

Therefore, we obtain the following local formulas on U (where we make use of Einstein's summation convention):

- $g = g_{ij} \omega^i \otimes \omega^j$ with $g_{ij} = g_{ji} \in \mathcal{O}_U$
- $\nabla_{\partial_i} \partial_j = \Gamma_{ij}^k \partial_k$ where the functions $\Gamma_{ij}^k \in \mathcal{O}_U$ are called the *Christoffel-symbols*.
- $\Gamma_{ij}^k = \frac{1}{2} g^{kl} (\partial_i g_{jk} + \partial_j g_{ik} - \partial_k g_{ij})$
- $R_{\partial_i \partial_j}(\partial_k) = R_{kij}^l \partial_l$
- $R_{ijk}^l = \partial_j \Gamma_{ki}^l - \partial_k \Gamma_{ij}^l + \Gamma_{ki}^r \Gamma_{jr}^l - \Gamma_{kr}^l \Gamma_{ij}^r$
- $R_{ik} := \operatorname{ric}_{ik} = R_{ilk}^l$
- $R := \operatorname{sc} = g^{ik} R_{ik}$

Now, the Einstein equations take their well known form

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = \kappa T_{\mu\nu} \quad \text{or equivalently} \quad R_{\mu\nu} = \kappa \left(T_{\mu\nu} - \frac{1}{2}g_{\mu\nu}T \right), \quad T := g^{\mu\nu}T_{\mu\nu},$$

and it follows that Theorem 1.37 holds.

1.37 Theorem *Let g be a metric on a smooth S -scheme $X \rightarrow S$. The Einstein equations on X are universal in the following sense: For $x \in X$, let $\{\partial_i\}$ be a base of $\Omega_{X/S,x}^1$, and let $g_{ij} \in \mathcal{O}_{X,x}$ be the components of the metric tensor at x . Assume that there exists a tensor $G \neq g$ of rank two such that for all $x \in X$ the following statements hold at x :*

- a) *G is a polynomial over K in the variables g_{ij} , $\partial_k g_{ij}$ and $\partial_k(\partial_l g_{ij})$ which is linear in $\partial_k(\partial_l g_{ij})$.*
- b) *G is a symmetrical tensor.*
- c) *$\text{div}(G) = 0$.*

Then, G coincides with the Einstein tensor $\text{ric} - \frac{1}{2} \text{sc} \cdot g$.

1.38 Remark Finally, let us again point to the crucial fact that the local functions $f \in \mathcal{O}_U$ are *algebraic* functions. Therefore, the class of functions, which is available in order to solve the equations of arithmetic general relativity, is much smaller than in the differential geometric setting. In general, it is a non-trivial task to solve differential equations in the algebraic setting, and it is not clear whether there exist solutions at all.

But, as we will see in the next chapter, the equations of general relativity have the amazing property that also many classical, differential geometric solutions descent to the algebraic category. Thus, the set of models of type (GR) $(X \rightarrow S, g)$ is not the empty set.

2. Solutions of the arithmetic Einstein equations

In chapter 1, we deduced the algebraic geometric analogue of the differential geometric Einstein equations. However, it is not clear at all if there actually exist pairs $(X \rightarrow S, g)$ consisting of an algebraic space $X \rightarrow S$ and a metric g on X which solves the Einstein equations. However, we will see in this chapter that the Einstein equations already possess many physically interesting solutions in the algebraic category. This is a highly non-trivial property of the Einstein equations, because, in general, differential equations only possess solutions which are very transcendental and far from the algebraic category. A very interesting solution, which may be interpreted as the “adelic” Minkowski space, is deduced and studied in section 2.2.2.

As already announced earlier, we will consider two special choices for the base S :

- S is a Zariski zero-dimensional Dedekind scheme.
- S is a Zariski one-dimensional Dedekind scheme.

2.1. The case of Zariski zero-dimensional base

Within this section, let $S = \operatorname{Spec} K$ be the spectrum of a field K . We are looking for models of type (GR) $(X \rightarrow S, g)$ (see Definition 0.4). However, the condition (ii) of Definition 0.4 is empty in this case. Therefore, from the adelic point of view, $S = \operatorname{Spec} K$ is not the physically interesting case, but rather a toy model. The interesting “adelic” models, where all conditions of Definition 0.4 are non-trivial will be considered in section 2.2. At least there are many models of type (GR) for the choice $S = \operatorname{Spec} K$, because things are particularly easy in this case. The most easiest example is the Minkowski solution

$$(\mathbb{A}_K^n, g_0), \quad g_0 := \operatorname{diag}(\pm 1, \pm 1, \pm 1, \pm 1)$$

However, there are further, less trivial examples which correspond to certain solutions of the classical differential geometric Einstein equations.

2.1.1. Kasner solution

Again we choose $X_K := \mathbb{A}_K^n$ with coordinates $(t = x^0, x^1, \dots, x^{n-1})$, but this time we choose the following non-trivial metric g_K

- $g_{00} = 1, \quad g_{0i} = 0, \quad i \neq 0$

$$\bullet \quad g_{ij} := c\delta_{ij} \cdot t^{2k_i}, \quad i \neq 0 \neq j$$

where δ_{ij} denotes the Kronecker delta, and where $c \in K$ and $k_i \in \mathbb{Q}$ are constants. Then g_K is well defined in the category of algebraic spaces, and it remains to show that we can choose the constants in such a way that the Einstein equations are fulfilled. We may do this on stalks. Recalling the remarks below Proposition 1.36 and the formulas stated there, we may compute the Christoffel symbols corresponding to the given metric. Due to the fact that we use orthogonal coordinates we obtain:

$$\Gamma_{00}^0 = 0, \Gamma_{i0}^0 = 0, \Gamma_{ii}^i = \frac{1}{2}g^{ii}\partial_i g_{ii} = 0, \Gamma_{ii}^j = -\frac{1}{2}g^{jj}\partial_j g_{ii} = 0, \Gamma_{ij}^j = -\frac{1}{2}g^{jj}\partial_i g_{jj} = 0.$$

The only non-vanishing Christoffel symbols are

$$(*) \quad \Gamma_{ii}^0 = -\frac{1}{2}g^{00}\partial_0 g_{ii} = -ck_i t^{2k_i-1}$$

$$(**) \quad \Gamma_{0i}^i = \frac{1}{2}g^{ii}\partial_0 g_{ii} = \frac{k_i}{t}$$

We consider a vacuum solution of the Einstein equations, i.e. $T_{\mu\nu} = 0$. Therefore, the Ricci tensor $R_{\mu\nu} = R_{\mu\lambda\nu}^\lambda$ has to vanish, too. The components of the Ricci tensor are as follows:

2.1 Lemma *The Ricci tensor is diagonal in the given coordinates, i.e. $R_{\mu\nu} = \delta_{\mu\nu}R_{\mu\nu}$. For the diagonal elements one obtains:*

$$R_{00} = \frac{1}{t^2} \sum_{j \neq 0} (k_j - k_j^2), \quad R_{ii} = -ck_i t^{2(k_i-1)} \left(\sum_{j \neq 0} k_j - 1 \right), i \neq 0.$$

2.2 Corollary *Let $X_K := \mathbb{A}_K^n$ and g_K be as stated above. Furthermore, choose the constants $k_i \in \mathbb{Q}$ such that*

$$\sum_{j \neq 0} k_j = 1 = \sum_{j \neq 0} k_j^2.$$

Then (X_K, g_K) is a model of type (GR) .

It remains to prove Lemma 2.1.

PROOF First of all, $R_{00} = \partial_\mu \Gamma_{00}^\mu - \partial_0 \Gamma_{\mu 0}^\mu + \Gamma_{\mu\nu}^\mu \Gamma_{00}^\nu - \Gamma_{\nu 0}^\mu \Gamma_{\mu 0}^\nu = -\partial_0 \Gamma_{\mu 0}^\mu - \Gamma_{\nu 0}^\mu \Gamma_{\mu 0}^\nu$, because $\Gamma_{00}^\mu = 0$. Furthermore, $\Gamma_{\nu 0}^\mu \Gamma_{\mu 0}^\nu = \Gamma_{\nu 0}^0 \Gamma_{00}^\nu + \sum_{j \neq 0} \Gamma_{\nu 0}^j \Gamma_{j0}^\nu = \sum_{j \neq 0} \Gamma_{\nu 0}^j \Gamma_{j0}^\nu \delta_{\nu j} = \sum_{j \neq 0} (\Gamma_{j0}^j)^2$ where we used $\Gamma_{00}^\mu = 0$ and formula $(**)$ in order to obtain the second equality. Thus

$$R_{00} = - \sum_{j \neq 0} \left(\partial_0 \Gamma_{j0}^j + (\Gamma_{j0}^j)^2 \right)$$

and the desired formula follows from $(*)$. Let now $i \neq 0$. Then $R_{i0} = R_{i\mu 0}^\mu = \partial_\mu \Gamma_{i0}^\mu - \partial_0 \Gamma_{i\mu}^\mu + \Gamma_{\nu\mu}^\mu \Gamma_{i0}^\nu - \Gamma_{0\nu}^\mu \Gamma_{i\mu}^\nu$. Recalling that only over Greek letters is summed, we obtain

$\partial_\mu \Gamma_{i0}^\mu = \partial_i \Gamma_{i0}^i = 0$, because the Christoffel symbols are constant with respect to x^i , $i \neq 0$. Furthermore, it follows from $(**)$ that $\Gamma_{\nu\mu}^\mu \Gamma_{i0}^\nu = \Gamma_{0\mu}^\mu \Gamma_{i0}^0$ and that $\Gamma_{0\nu}^\mu \Gamma_{i\mu}^\nu = \sum_{j \neq 0} \Gamma_{0j}^j \Gamma_{ij}^j$. However, as well Γ_{ij}^j as Γ_{i0}^0 are zero and we obtain

$$R_{i0} = 0.$$

Similarly, $R_{ik} = 0$ if $i, k \neq 0$, $i \neq k$. This may be seen as follows: $R_{ik} = R_{i\mu k}^\mu = \partial_\mu \Gamma_{ik}^\mu - \partial_k \Gamma_{i\mu}^\mu + \Gamma_{\nu\mu}^\mu \Gamma_{ik}^\nu - \Gamma_{k\nu}^\mu \Gamma_{i\mu}^\nu$. Thereby, $\partial_k \Gamma_{i\mu}^\mu = 0$ because the Christoffel symbols are constant with respect to x^k , $k \neq 0$. In addition, $\Gamma_{ik}^\mu = 0$ for all μ . Thus, it remains to show that $\Gamma_{k\nu}^\mu \Gamma_{i\mu}^\nu = \Gamma_{kk}^0 \Gamma_{i0}^k + \Gamma_{k0}^i \Gamma_{ii}^0$ is zero. But this is clear, because $\Gamma_{i0}^k = 0$ due to $i, k \neq 0$, $i \neq k$. Finally, let us prove that $R_{ii} = R_{i\mu i}^\mu = \partial_\mu \Gamma_{ii}^\mu - \partial_i \Gamma_{i\mu}^\mu + \Gamma_{\nu\mu}^\mu \Gamma_{ii}^\nu - \Gamma_{i\nu}^\mu \Gamma_{i\mu}^\nu$ is as claimed. For the first summand we obtain $\partial_\mu \Gamma_{ii}^\mu = \partial_0 \Gamma_{ii}^0 = -ck_i(2k_i - 1)t^{2(k_i-1)}$. The second summand vanishes, because the Christoffel symbols are constant with respect to x^i , $i \neq 0$. The third summand reads as $\Gamma_{\nu\mu}^\mu \Gamma_{ii}^\nu = \sum_{j \neq 0} \Gamma_{0j}^j \Gamma_{ii}^0 = \sum_{j \neq 0} k_j t^{-1} (-ck_i t^{2k_i-1})$. Finally, $\Gamma_{i\nu}^\mu \Gamma_{i\mu}^\nu = \Gamma_{ii}^0 \Gamma_{i0}^i + \Gamma_{i0}^i \Gamma_{ii}^0 = -2ck_i^2 t^{2(k_i-1)}$. Everything summed up gives the desired result. $\square$

2.1.2. Schwarzschild solution

The example of the Schwarzschild metric will show very clearly the general phenomenon that the Zariski topology is too coarse for physical applications and that it is necessary to work within the context of the étale topology. However, let us again start from the affine space $\mathbb{A}_K^n$ with coordinates $(t = x^0, x^1, \dots, x^{n-1})$, but this time we consider the K -scheme $X_K := \text{Spec } K[t, x^1, \dots, x^{n-1}, r, r^{-1}] / (r^2 - \sum_{i \neq 0} (x^i)^2)$, whereby $r := \sqrt{\sum_{i \neq 0} (x^i)^2}$ should be interpreted as a spacial radius. By construction, X_K is étale over $\mathbb{A}_K^n$. In particular, the respective differential calculi “coincide”. We choose the following metric g_K on X_K :

- $g_{00} = \frac{1}{1 + \frac{2m}{r}}$, $g_{0i} = 0$, $i \neq 0$
- $g_{ij} := -\left(1 + \frac{2m}{r}\right)^2 \delta_{ij} + \frac{x^i x^j}{r^2} \left(1 + \frac{2m}{r}\right) \frac{2m}{r}$

where δ_{ij} denotes the Kronecker delta, and $m \in K$ is a constant. By means of a longer but standard calculation similar to the one in the Kasner case (see section 2.1.1), one can prove that the metric above solves the vacuum Einstein equations. Thus (X_K, g_K) is indeed a model of type (GR). But instead of deriving this result directly, we will illustrate that it corresponds to the Schwarzschild metric. This will yield the following physical interpretation: (X_K, g_K) describes the exterior of a black hole. More precisely, r scales the distance from the event horizon of the black hole, and the constant m turns out to be the Schwarzschild diameter of the black hole.

In differential geometry, one usually writes down the Schwarzschild metric in spherical coordinates. These coordinates do not make sense in algebraic geometry. But, at least in the category of Riemannian manifolds we can prove that the metric above takes the classical form in spherical coordinates. Writing

$$(x^0 = t, x^1 = r \sin(\theta) \cos(\varphi), x^2 = r \sin(\theta) \sin(\varphi), x^3 = r \cos(\theta))$$

in the four dimensional case, one obtains:

- $dR^2 = dr^2 = \sum_{i,j \neq 0} \frac{x^i x^j}{r^2} dx^i dx^j$, where $R := r + c$ for some constant $c \in K$.
- $\sum_{i \neq 0} (dx^i)^2 = dr^2 + r^2(d\theta^2 + \sin^2(\theta)d\varphi^2)$.

It follows that

$$\begin{aligned}
\frac{1}{1 - \frac{2m}{R}} dR^2 + R^2(d\theta^2 + \sin^2(\theta)d\varphi^2) &= \sum_{i,j \neq 0} \left(\frac{R}{R - 2m} \frac{x^i x^j}{r^2} + \frac{R^2}{r^2} \left(\delta_{ij} - \frac{x^i x^j}{r^2} \right) \right) dx^i dx^j \\
&= \sum_{i,j \neq 0} \left(\frac{R^2}{r^2} \delta_{ij} + \frac{x^i x^j}{r^2} \left(\frac{R}{R - 2m} - \frac{R^2}{r^2} \right) \right) dx^i dx^j \\
&= \sum_{i,j \neq 0} \left(\left(1 + \frac{c}{r} \right)^2 \delta_{ij} + \frac{x^i x^j}{r^2} \left(\frac{r + c}{r + c - 2m} - \frac{(r + c)^2}{r^2} \right) \right) dx^i dx^j \\
&= \sum_{i,j \neq 0} \left(\left(1 + \frac{c}{r} \right)^2 \delta_{ij} - \frac{x^i x^j}{r^2} (r + c) \frac{2r(c - m) + c(c - 2m)}{(r + c - 2m)r^2} \right) dx^i dx^j \\
&= \sum_{i,j \neq 0} \left(\left(1 + \frac{2m}{r} \right)^2 \delta_{ij} - \frac{x^i x^j}{r^2} \left(1 + \frac{2m}{r} \right) \frac{2m}{r} \right) dx^i dx^j \quad \text{for } c = 2m.
\end{aligned}$$

Therefore, our original metric transforms into the standard form of the Schwarzschild metric

$$ds^2 = \left(1 - \frac{2m}{R} \right) dt^2 - \frac{1}{1 - \frac{2m}{R}} dR^2 - R^2(d\theta^2 + \sin^2(\theta)d\varphi^2)$$

living on the manifold $M = \mathbb{R} \times (2m, \infty) \times S_2$ with coordinates (t, R, θ, φ) .

2.1.3. Electromagnetism in vacuum

In this subsection we will consider the electromagnetic field. This is a gauge field, and so we may apply the techniques developed in Part II of this thesis. Therefore, let us consider a smooth S -scheme X (which represents space-time) and a smooth S -group scheme G (see Definition 11.46) representing the gauge group. The fundamental structure underlying gauge theory is an X -torsor P under G_X (see Definition 8.7), where G_X denotes the X -group scheme $G \times_S X$. Then the field strength of the gauge field is given by a differential two-form $\mathcal{F}$ on X which solves the Yang-Mills equations 9.8. The differential form $\mathcal{F}$ takes values in the Lie-algebra $\mathfrak{g} := (\varepsilon^* \Omega_{G/S}^1)(S)$ of G , where $\varepsilon : S \rightarrow G$ denotes the unit section of G (see section 11.4.1).

If $S = \text{Spec } K$ is the spectrum of a field K , we will illustrate in section 3.5 that the gauge group of electromagnetism is the algebraic torus $\mathbb{G}_{m,K} := \text{Spec } K[\zeta, \zeta^{-1}]$ for some variable ζ . In absence of gravity we may choose $X = \mathbb{A}_K^n$ as space-time part, where

$n \in \mathbb{N}$ denotes the dimension. Therefore, electromagnetism in vacuum may be described by means of the trivial $\mathbb{A}_K^n$ -torsor $P = \mathbb{A}_K^n \times_{\mathbb{A}_K^n} (\mathbb{A}_K^n \times_K \mathbb{G}_{m,K})$ under $\mathbb{A}_K^n \times_K \mathbb{G}_{m,K}$, i.e.

$$P \cong \mathbb{A}_K^n \times_K \mathbb{G}_{m,K}.$$

In this setting, we will first compute the Lie-algebra $\mathfrak{g}$ of G in order to solve the Yang-Mills equations in the electrostatic case in a second step. One obtains the following two results.

2.3 Lemma *Let $S = \operatorname{Spec} R$ be the spectrum of a ring R , and let $\mathfrak{g}$ be the Lie algebra of the multiplicative group $\mathbb{G}_{m,S} = \operatorname{Spec} R[\zeta, \zeta^{-1}]$ over R . Then (with respect to the chosen coordinates of $\mathbb{G}_{m,S}$) there is a canonical isomorphism $\mathfrak{g} \cong R$ of R -modules.*

PROOF As an S -scheme, $\mathbb{G}_{m,S}$ coincides with $\operatorname{Spec} R[\zeta, \zeta^{-1}]$. The ring $R[\zeta, \zeta^{-1}]$ is the localization $R[\zeta]_{\zeta}$ of the polynomial ring $R[\zeta]$ with respect to the variable ζ . Therefore, the canonical morphism $i : \operatorname{Spec} R[\zeta, \zeta^{-1}] \hookrightarrow \operatorname{Spec} R[\zeta] = \mathbb{A}_S^1$ is an open immersion. As open immersions are étale, it follows from Proposition 3.1 that $\Omega_{\mathbb{G}_{m,S}/S}^1 \cong i^* \Omega_{\mathbb{A}_S^1/S}^1$ is an isomorphism. By definition, the Lie-algebra $\mathfrak{g}$ is the R -module $(\varepsilon^* \Omega_{\mathbb{G}_{m,S}/S}^1)(S)$, where $\varepsilon : S \rightarrow \mathbb{G}_{m,S}$ denotes the unit section corresponding to the morphism of R -algebras $\varepsilon^* : R[\zeta, \zeta^{-1}] \rightarrow R$, $\zeta \mapsto 1 \in R^*$, mapping ζ into the group of units R^* of R . In order to compute $\mathfrak{g}$, first notice that $\varepsilon^* \Omega_{\mathbb{G}_{m,S}/S}^1 \cong \varepsilon^* i^* \Omega_{\mathbb{A}_S^1/S}^1 \cong (i \circ \varepsilon)^* \Omega_{\mathbb{A}_S^1/S}^1$, whereby $i \circ \varepsilon$ corresponds to the R -algebra homomorphism $R[\zeta] \rightarrow R$, $\zeta \mapsto 1 \in R^*$. Recalling that $\Omega_{\mathbb{A}_S^1/S}^1(S)$ is the free $R[\zeta]$ -module $R[\zeta] \cdot d\zeta$ generated by the differential $d\zeta$, we obtain the following R -module isomorphisms:

$$\mathfrak{g} = \left((i \circ \varepsilon)^* \Omega_{\mathbb{A}_S^1/S}^1 \right) (S) \cong \Omega_{\mathbb{A}_S^1/S}^1(S) \otimes_{R[\zeta]} R \cong R[\zeta] \otimes_{\operatorname{id}, R[\zeta], (i \circ \varepsilon)^*} R \cong R. \quad \square$$

2.4 Corollary *Let S be an affine scheme, let X be an smooth, affine S -scheme, and let $G = \mathbb{G}_m$. Let $P = X \times_S G$ be the trivial X -torsor under $G_X = G \times_S X$. Then the field strength $\mathcal{F}$ may be viewed as differential two form on X , i.e. $\mathcal{F} \in \Omega_{X/S}^2(X)$.*

PROOF (of Corollary 2.4): By Definition 9.4, the field strength $\mathcal{F} = s^* \Omega$ is locally defined as pull back of a Lie-algebra valued global differential form $\Omega \in \Omega_{P/S}^2(P) \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$ under the canonical section s . As $\mathfrak{g} \cong \mathcal{O}_S(S)$ due to Lemma 2.3, and we may view Ω as differential form $\Omega \in \Omega_{P/S}^2(P)$ on P .

In general, the canonical section s is only defined locally in étale topology, but in our special situation, s coincides with the morphism obtained from the unit section $\varepsilon : S \rightarrow G$ by base change with $\pi : X \rightarrow S$. Therefore, s makes the following diagram commutative.

$$\begin{array}{ccc} & & X \\ & \nearrow id & \uparrow p_1 \\ X & \xrightarrow{s} & X \times_S G \\ & \searrow \varepsilon \circ \pi & \downarrow p_2 \\ & & G \end{array}$$

By Proposition 1.7, we have got $(s^*\Omega_{P/S}^1)(X) \cong ((p_1 \circ s)^*\Omega_{X/S}^1)(X) \oplus ((p_2 \circ s)^*\Omega_{G/S}^1)(X) \cong (\text{id}^*\Omega_{X/S}^1)(X) \oplus ((\varepsilon \circ \pi)^*\Omega_{G/S}^1)(X) \cong \Omega_{X/S}^1(X) \oplus (\mathfrak{g} \otimes_{\mathcal{O}_S(S)} \mathcal{O}_X(X))$, and therefore the pull-back of differential forms under s is the canonical projection

$$(s^*\Omega_{P/S}^1)(X) \cong \Omega_{X/S}^1(X) \oplus \mathcal{O}_X(X) \rightarrow \Omega_{X/S}^1(X). \quad \square$$

Now, we are prepared to compute the electromagnetic field in vacuum, i.e. we will solve the corresponding Yang-Mills equations. Let K be a field. As already stated above, we choose $S = \text{Spec } K$, $G = \mathbb{G}_{m,K} \cong \text{Spec } K[\zeta, \zeta^{-1}]$, $X = \mathbb{A}_K^4$ with metric $g = \text{diag}(-1, 1, 1, 1)$ and $P = X \times_S G$. Due to the fact that the group scheme G is commutative, the Lie-bracket on $\mathfrak{g}$ vanishes (see section 11.4.3). But then we know from Definition 10.3 that the covariant derivation D_X on X coincides with the exterior differential d introduced in Theorem 11.39. Furthermore, $\Omega_{X/S}^1(X) \cong \oplus_{i=0}^3 \mathcal{O}_X(X) dx^i$ is a free $\mathcal{O}_X(X)$ module, where $(x^0 = t, x^1, x^2, x^3)$ denote the coordinates of X . Thus we may write $\mathcal{F} = \sum_{\mu\nu} \mathcal{F}_{\mu\nu} dx^\mu \wedge dx^\nu$ with polynomials $\mathcal{F}_{\mu\nu} \in \mathcal{O}_X(X) = K[t, x^1, x^2, x^3]$. The components of the anti-symmetric tensor $\mathcal{F}$ are usually written as matrix:

$$(\mathcal{F}_{\mu\nu}) = \begin{pmatrix} 0 & E_1 & E_2 & E_3 \\ -E_1 & 0 & B_1 & B_2 \\ -E_2 & -B_1 & 0 & B_3 \\ -E_3 & -B_2 & -B_3 & 0 \end{pmatrix}$$

The vector $\vec{E} := (E_1, E_2, E_3)$ is called the electric field strength, $\vec{B} := (B_1, B_2, B_3)$ denotes the magnetic field strength. We are interested in an electrostatic solution, i.e. $\vec{B} = 0$. In this special case the Yang-Mills equations

$$\begin{aligned} *d * \mathcal{F} &= 0 \\ d\mathcal{F} &= 0 \end{aligned}$$

take the form

$$\begin{aligned} \sum_{i \neq 0} \frac{\partial}{\partial x^i} E_i &= 0 \\ \frac{\partial}{\partial x^i} E_j - \frac{\partial}{\partial x^j} E_i &= 0 \quad \text{for all } i, j \neq 0. \end{aligned}$$

This may be seen as follows: Due to the fact that $\mathcal{O}_X(X)$ is a polynomial ring, the local version of the Yang-Mills equations 10.9 is not only true at physical points, but even globally. Recalling that $g = \text{diag}(-1, 1, 1, 1)$ and that $c_{ij}^k = 0$ (due to the commutativity of the gauge group), the Yang-Mills equations take the form $\sum_{\mu, \rho, \sigma=1}^n \frac{\partial}{\partial x^\sigma} (g^{\mu\nu} g^{\rho\sigma} \mathcal{F}_{\mu\rho}) = 0$. But because of $\vec{B} = 0$, $\mathcal{F}_{\mu\nu} \neq 0$ if and only if exactly one index is zero. Thus, the only non trivial equation is: $0 = \sum_{\mu, \rho, \sigma=1}^n \frac{\partial}{\partial x^\sigma} (g^{\mu 0} g^{\rho\sigma} \mathcal{F}_{\mu\rho}) = -\sum_{\sigma=1}^n \frac{\partial}{\partial x^\sigma} \mathcal{F}_{0\sigma} = -\sum_{i \neq 0} \frac{\partial}{\partial x^i} E_i$. In order to derive the second equation, recall that we may write $\mathcal{F} =$

$\sum_{i \neq 0} E_i dt \wedge dx^i$. Thus, $d\mathcal{F} = \sum_{i,j \neq 0} \frac{\partial E_i}{\partial x^j} dx^j \wedge dt \wedge dx^i = \sum_{i,j \neq 0} \frac{\partial E_i}{\partial x^j} dt \wedge dx^i \wedge dx^j$, and if ∂_i denotes the dual of dx^i it follows from Definition 11.34, b) that

$$\begin{aligned} d\mathcal{F}(\partial_0, \partial_k, \partial_l) &= \sum_{i,j \neq 0} \frac{\partial E_i}{\partial x^j} \cdot \frac{1}{2} dt(\partial_0) \left((dx^i \wedge dx^j)(\partial_k, \partial_l) - (dx^i \wedge dx^j)(\partial_l, \partial_k) \right) \\ &= \sum_{i,j \neq 0} \frac{\partial E_i}{\partial x^j} \cdot (dx^i \wedge dx^j)(\partial_k, \partial_l) \\ &= \sum_{i,j \neq 0} \frac{\partial E_i}{\partial x^j} (\delta_{ik} \delta_{jl} - \delta_{il} \delta_{jk}) = \frac{\partial E_k}{\partial x^l} - \frac{\partial E_l}{\partial x^k} \end{aligned}$$

It is well known that $E_i := \frac{x^i}{r}$, $r^2 := \sum_{i \neq 0} (x^i)^2$ is a solution of these equations. Obviously, E_i is neither a global section of X nor of $X - \{0\} = \mathbb{A}_K^n - \{0\}$. Instead we may consider the smooth K -scheme, $U := \text{Spec } K[t, x^1, x^2, x^3, r, r^{-1}] / (r^2 - \sum_{i \neq 0} (x^i)^2)$ which is étale over $\mathbb{A}_K^4$ and contains r as global function. The étale-open “sub-scheme” U of $\mathbb{A}_K^4$ describes a space-time which contains a charged point particle located at $r = 0$ (more precisely, U is the environment of this charged particle). The electrostatic field of this point charge is described by global sections $E_i \in \mathcal{O}_U(U)$.

2.1.4. Robertson-Walker models

Last but not least, let us briefly mention the Robertson-Walker models. In differential geometry, these are defined in spherical coordinates by

$$\begin{aligned} g_{\mu\nu} &= \text{diag} \left(1, \frac{-S(t)^2}{1 - kr^2}, -S(t)^2 r^2, -S(t)^2 r^2 \sin^2(\theta) \right) \\ T_{\mu\nu} &= \text{diag} \left(\rho(t), \frac{-p(t)S(t)^2}{1 - kr^2}, -p(t)S(t)^2 r^2, -p(t)S(t)^2 r^2 \sin^2(\theta) \right) \end{aligned}$$

where $S(t)$ is a so called scale factor, and where ρ resp. p denote the density resp. pressure of energy. In the special case $p = 0$, the divergence $\text{div} T$ of the energy-stress tensor vanishes if and only if the product $C := \rho(t) \cdot S(t)^3$ is constant with respect to t . Then the Einstein equations $R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = \kappa T_{\mu\nu}$ yield the differential equation

$$\left(\frac{\partial S(t)}{\partial t} \right)^2 + k = \frac{\kappa C}{3S(t)}.$$

Whenever the solution $S(t)$ of this differential equation is an algebraic function, the corresponding Robertson-Walker metric would make sense in algebraic geometry. However, in general, $S(t)$ will not be algebraic. So, the classical Robertson-Walker models have no algebraic geometric analogues. But at least in the case $k = 0$, we find the algebraic solution

$$S(t) = t^{\frac{2}{3}} \sqrt[3]{4/3 \cdot \kappa C}.$$

Therefore, we obtain a model of type (GR) if we choose $X_K := \mathbb{A}_K^n$ with coordinates $(t = x^0, x^1, \dots, x^{n-1})$ as well as the following metric and energy-stress tensor:

$$g_{\mu\nu} = \text{diag} \left(1, -ct^{\frac{4}{3}}, -ct^{\frac{4}{3}}, -ct^{\frac{4}{3}} \right), \quad c := (4/3 \cdot \kappa C)^{\frac{2}{3}} \in K$$

$$T_{\mu\nu} = \left(\frac{4}{3t^2}, 0, 0, 0 \right).$$

2.2. The case of Zariski one-dimensional base

Within this section, let $S = \text{Spec } \mathcal{O}$ be the spectrum of a Dedekind ring which is not a field. From the “adelic” point of view, this is the physically interesting case, because the condition (ii) of Definition 0.4 is no longer empty. Consequently, it is much harder to construct models of type (GR).

In section 2.2.1, we will first consider the low dimensional case, because then the Einstein equations are trivial. But as soon as the tangent spaces exceed three dimensions, this is no longer true. Then, the conditions (i) and (ii) of Definition 0.4 are both non-trivial. This is the physically interesting situation where we are looking for models of type (GR) $(X \rightarrow S, g)$. An example, which may be interpreted as the “adelic” Minkowski space, will be studied in section 2.2.2.

2.2.1. The low dimensional case

Let $X \rightarrow S$ be a smooth S -scheme of relative dimension one or two with metric g . We will show that the Einstein equations are trivial in this case. As it suffices to show this locally, we may choose an open sub-scheme U of X such that $\Omega_{U/S}^1$ is free with base $\{\omega^i\}$. In the one dimensional case, our assertion is clear, because the curvature tensor has got only a single component, and this component vanishes due to the symmetries of the curvature tensor (see Proposition 1.30). Thus, also the Einstein tensor $\text{ric} - \frac{1}{2}g \cdot \text{sc}$ vanishes. In the two dimensional case, a small computation is necessary. Again making use of the identities of the curvature tensor several times, we obtain with respect to the given base:

$$\begin{aligned} R &:= \text{sc} = g^{\mu\nu} \text{ric}_{\mu\nu} \\ &= g^{\mu\nu} R_{\mu 1 \nu}^1 + g^{\mu\nu} R_{\mu 2 \nu}^2 = g^{\mu 2} R_{\mu 1 2}^1 + g^{\mu 1} R_{\mu 2 1}^2 = g^{\mu 2} g^{\nu 1} R_{\nu \mu 1 2} + g^{\mu 1} g^{\nu 2} R_{\nu \mu 2 1} \\ &= g^{11} g^{22} R_{1 2 1 2} + g^{21} g^{12} R_{2 1 1 2} + g^{11} g^{22} R_{2 1 2 1} + g^{21} g^{12} R_{1 2 2 1} \\ &= 2 \det g \cdot R_{2 1 2 1}. \end{aligned}$$

Therefore, one derives that

$$\begin{aligned}
R_\nu^\mu &:= \text{Ric}_\nu^\mu = g^{\mu\lambda} \text{ric}_{\lambda\nu} \\
&= g^{\mu\lambda} R_{\lambda 1\nu}^1 + g^{\mu\lambda} R_{\lambda 2\nu}^2 = g^{\mu\lambda} g^{1\iota} R_{\iota\lambda 1\nu} + g^{\mu\lambda} g^{2\iota} R_{\iota\lambda 2\nu} \\
&= g^{\mu 1} g^{12} R_{211\nu} + g^{\mu 2} g^{11} R_{121\nu} + g^{\mu 1} g^{22} R_{212\nu} + g^{\mu 2} g^{21} R_{122\nu} \\
&= R_{211\nu} (g^{\mu 1} g^{12} - g^{\mu 2} g^{11}) + R_{212\nu} (g^{\mu 1} g^{22} - g^{\mu 2} g^{21}) \\
&= \delta_{2\mu} \delta_{\nu 2} (g^{21} g^{12} - g^{22} g^{11}) R_{2112} + \delta_{\mu 1} \delta_{\nu 1} (-g^{21} g^{12} + g^{22} g^{11}) R_{2121} \\
&= R_{2121} \det g \cdot (\delta_{2\mu} \delta_{\nu 2} + \delta_{\mu 1} \delta_{\nu 1}) = \frac{1}{2} \delta_{\mu\nu} R.
\end{aligned}$$

Consequently, $R_\nu^\mu - \frac{1}{2} \delta_{\mu\nu} R = 0$ or equivalently $R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = 0$. In particular, there are many models of type (GR) in the low dimensional case.

2.5 Corollary *Let $X \rightarrow S$ be a smooth S -scheme of relative dimension one or two such that X is the Néron model of its generic fibre. Then X gives rise to a model of type (GR).*

2.2.2. A higher dimensional solution

Again, let $X \rightarrow S$ be smooth over the Zariski one-dimensional base scheme S . If the relative dimension exceeds two, the Einstein equations are no longer trivial. Therefore, it is not easy to find models of type (GR) in the “adelic” situation. However, in this section we will state at least one example, namely the fibred product $X := E_1 \times_S \dots \times_S E_n$ of smooth elliptic curves E_i over S . In a certain way, this is the “adelic” analogue of Minkowski space-time. Before we will finally prove in Theorem 2.9 that X is indeed a model of type (GR), we need some preparations. However, the idea of the proof is as follows: Locally, every smooth curve C over S may be embedded in some affine space $\mathbb{A}_S^n$. Pulling back the flat metric $\text{diag}(\pm 1, \dots, \pm 1)$ on $\mathbb{A}_S^n$ to C and pushing forward this metric by means of an étale morphism $C \rightarrow \mathbb{A}_S^1$ (which exists locally due to smoothness), we obtain the first fundamental form on C in local coordinates. Analogously, we obtain the first fundamental form on a product of curves in local coordinates. This metric is diagonal, because $\Omega_{(X_1 \times_S X_2)/S}^1 \cong \bigoplus_i p_i^* \Omega_{X_i/S}^1$, where p_i denotes the projection onto the i -th factor (see Lemma 2.6). It follows that the corresponding curvature tensor vanishes (see Corollary 2.8). Consequently, a product of elliptic curves is a vacuum solution of the Einstein equations and it is even a model of type (GR), because it is the Néron model of its generic fibre. However, let us now make the indicated steps of the proof explicit.

2.6 Lemma *Let $C_1, \dots, C_n$ be n smooth curves over S . Provide $X := C_1 \times_S \dots \times_S C_n$ with the first fundamental form g as metric (see Proposition 4.21). Then g is diagonal.*

PROOF It suffices to prove this statement locally on X . Therefore, we may assume that there are S -immersions $j_i : C_i \hookrightarrow Y_i := \mathbb{A}_S^m$ for some $m \in \mathbb{N}$ (see Definition 1.10). These S -immersions induce an S -immersion $j : X \hookrightarrow Y := \mathbb{A}_S^{nm}$. Furthermore, we know

from Proposition 1.17 that there is an open sub-scheme C'_i of C_i together with an étale morphism $g_i : C'_i \rightarrow \mathbb{A}_S^1$. Thus we obtain the following chain of maps:

$$Y_i \xleftarrow{j_i} C_i \xleftarrow{i} C'_i \xrightarrow{g_i} \mathbb{A}_S^1.$$

Eventually shrinking C_i , we may assume $C_i = C'_i$ and $i = \text{id}$. Recalling that the pull back of differential forms under étale morphisms is an isomorphism (see Proposition 3.1), these maps induce the following canonical homomorphism on the level of differential forms.

$$j_i^* \Omega_{\mathbb{A}_S^m/S}^1 \xrightarrow{j_i^*} \Omega_{C_i/S}^1 \xrightarrow[(g_i^*)^{-1}]{\sim} g_i^* \Omega_{\mathbb{A}_S^1/S}^1 \quad (*)$$

This morphism maps the generators dT_{ij} , $j = 1, \dots, m$, of the free module $j_i^* \Omega_{\mathbb{A}_S^m/S}^1$ to elements $\kappa_{ij} dT_i \in g_i^* \Omega_{\mathbb{A}_S^1/S}^1$, where dT_i denotes the generator of the free module $g_i^* \Omega_{\mathbb{A}_S^1/S}^1$.

Moreover, if $p_i : X \rightarrow C_i$ denotes the canonical projection, there is an isomorphism $\Omega_{X/S}^1 \cong \bigoplus_i p_i^* \Omega_{C_i/S}^1$ due to Proposition 1.7. Analogously, if $\pi_i : Y \rightarrow Y_i$ denotes the canonical projection making the diagram

$$\begin{array}{ccc} Y_i & \xleftarrow{j_i} & C_i \\ \uparrow \pi_i & & \uparrow p_i \\ Y & \xleftarrow{j} & X \end{array}$$

commutative, there is an isomorphism $\Omega_{\mathbb{A}_S^{nm}/S}^1 \cong \bigoplus_i \pi_i^* \Omega_{\mathbb{A}_S^m/S}^1$. Now, the pull-back $j^* \Omega_{\mathbb{A}_S^{nm}/S}^1 \rightarrow \Omega_{X/S}^1$ decomposes into a direct sum of componentwise pull-backs, i.e. the following diagram is commutative:

$$\begin{array}{ccc} j^* \Omega_{\mathbb{A}_S^{nm}/S}^1 & \xrightarrow{\sim} & \bigoplus_i j^* \pi_i^* \Omega_{\mathbb{A}_S^m/S}^1 \xrightarrow{\sim} \bigoplus_i p_i^* (j_i^* \Omega_{\mathbb{A}_S^m/S}^1) \\ \downarrow j^* & & \downarrow \bigoplus j_i^* \\ \Omega_{X/S}^1 & \xrightarrow{\sim} & \bigoplus_i p_i^* \Omega_{C_i/S}^1 \end{array}$$

The upper line of this diagram yields a morphism

$$j^* \Omega_{\mathbb{A}_S^{nm}/S}^1 \xrightarrow{\sim} \bigoplus_i p_i^* (j_i^* \Omega_{\mathbb{A}_S^m/S}^1) \xrightarrow{\text{pr}_i} p_i^* (j_i^* \Omega_{\mathbb{A}_S^m/S}^1) \quad (**)$$

where pr_i denotes the projection onto the i -th component. Pulling back the morphism $(*)$ with p_i and composing it with the above morphism $(**)$, we finally obtain the morphism

$$\begin{array}{ccccc} \phi_i : j^* \Omega_{\mathbb{A}_S^{nm}/S}^1 & \longrightarrow & (j_i \circ p_i)^* \Omega_{\mathbb{A}_S^m/S}^1 & \longrightarrow & (g_i \circ p_i)^* \Omega_{\mathbb{A}_S^1/S}^1 \\ dT_{kj} & \mapsto & \delta_{ik} dT_{kj} & \mapsto & \delta_{ik} \kappa_{kj} dT_i, \end{array}$$

where $\kappa_{ij} \in \mathcal{O}_X$ is algebraic over $\mathcal{O}_S[T_i]$ and δ_{ik} denotes the Kronecker delta. Now, we have all that we need to compute the first fundamental form. On $Y = \mathbb{A}_S^{nm}$ we choose

the trivial metric $g_0 = \sum_{\mu\nu} \varepsilon_{\mu\nu} dT_{\mu\nu} \otimes dT_{\mu\nu}$, $\varepsilon_{\mu\nu} \in \{\pm 1\}$. Then, by definition, the first fundamental form g reads as follows:

$$g = \sum_{i,j,\mu,\nu} \varepsilon_{\mu\nu} \phi_i(dT_{\mu\nu}) \otimes \phi_j(dT_{\mu\nu}) = \sum_{i,j,\mu,\nu} \varepsilon_{\mu\nu} \delta_{i\mu} \kappa_{\mu\nu} \delta_{j\mu} \kappa_{\mu\nu} dT_i \otimes dT_j$$

Therefore, $g = \sum_{ij} g_{ij} dT_i \otimes dT_j$ with $g_{ij} = \delta_{ij} \cdot \sum_{\nu} \varepsilon_{i\nu} \kappa_{i\nu}^2$. In particular, g is diagonal. $\square$

2.7 Example In order to illustrate the derivation of the first fundamental form, let us consider the example of a smooth elliptic curve E over a field K which is not of characteristic two. Let us now restrict E to the affine open subset $\text{Spec } K[X, Y, Y^{-1}] \subset \mathbb{P}_K^2$, and let us furthermore assume that E is described by an equation $P(Y, X) := Y^2 - X^3 - g_2X - g_3 = 0$. Then we have canonical morphisms of K -algebras

$$\begin{array}{ccccc} K[X, Y, Y^{-1}] & \xrightarrow{j^*} & \mathcal{O}_E := K[X, Y, Y^{-1}]/(P) & \xleftarrow{g^*} & K[X] \\ X & \mapsto & \bar{X} & \longleftarrow & X \\ Y & \mapsto & \bar{Y} & & \end{array}$$

where g^* is étale. Therefore, we obtain on the level of differential forms

$$\begin{array}{ccccc} \Omega_{K[X, Y, Y^{-1}]/K}^1 \otimes_{K[X, Y, Y^{-1}]} \mathcal{O}_E & \xrightarrow{j^*} & \Omega_{\mathcal{O}_E/K}^1 & \xrightarrow[(g^*)^{-1}]{\sim} & \Omega_{K[X]/K}^1 \otimes_{K[X]} \mathcal{O}_E \\ dX & \mapsto & d\bar{X} & \mapsto & dX \\ dY & \mapsto & d\bar{Y} & \mapsto & \frac{3X^2 + g_2}{2Y} dX \end{array}$$

where we made use of the identity $(3\bar{X}^2 + g_2)d\bar{X} = d\bar{Y}^2 = 2\bar{Y}d\bar{Y}$ in $\Omega_{\mathcal{O}_E/K}^1$. This is the map $(*)$ in the proof of Lemma 2.6. The dual of this map is the $\mathcal{O}_E$ -linear map $\partial_X \mapsto \partial_X + \frac{3X^2 + g_2}{2Y} \partial_Y$, where ∂_X (resp. ∂_Y) denotes the dual of dX (resp. dY). In particular, the tangent vectors of E in the affine open subset $\text{Spec } K[X, Y, Y^{-1}] \subset \mathbb{P}_K^2$ may be written as vectors

$$\left(\begin{array}{c} 1 \\ \frac{3X^2 + g_2}{2Y} \end{array} \right).$$

Providing $\text{Spec } K[X, Y, Y^{-1}]$ with the trivial metric $g_0 = \text{diag}(1, 1)$ and interpreting g_0 as bilinear form, we derive the first fundamental form g on E :

$$g = g_0 \left(\left(\begin{array}{c} 1 \\ \frac{3X^2 + g_2}{2Y} \end{array} \right), \left(\begin{array}{c} 1 \\ \frac{3X^2 + g_2}{2Y} \end{array} \right) \right) = 1 + \frac{(3X^2 + g_2)^2}{4Y^2}.$$

This is manifestly the same result as in differential geometry. The procedure in the case of a product of n elliptic curves is straight forward. For example, if $n = 2$, we have to compose the above homomorphism of K -algebras with the projection map $K[X, Y, Y^{-1}, Z, W] \rightarrow K[X, Y, Y^{-1}]$, $Z, W \mapsto 0$, where Z and W are the variables of the

second elliptic curve. In this case we obtain the two tangent vectors

$$\begin{pmatrix} 1 \\ \frac{3X^2+g_2}{2Y} \\ 0 \\ 0 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 0 \\ 0 \\ 1 \\ \frac{3Z^2+g'_2}{2W} \end{pmatrix}.$$

In particular, the metric is diagonal.

2.8 Corollary *Let X be as in Lemma 2.6. Then X is flat, i.e. the curvature tensor vanishes.*

PROOF As the curvature tensor is a global section of a sheaf, it suffices to prove the statement locally. Therefore, we may assume that the components g_{ij} of the metric tensor g are algebraic over some polynomial ring, say with variables $x_1, \dots, x_n$. In particular we may use the Christoffel symbols (as well as the formulas stated at the end of section 1.2) in order to compute the curvature tensor. Due to Lemma 2.6, the metric is diagonal. Consequently, the Christoffel symbols read as follows:

$$\begin{aligned} \Gamma_{ii}^i &= \frac{1}{2} g^{ii} \partial_i g_{ii}. \\ \Gamma_{ii}^k &= -\frac{1}{2} g^{kk} \partial_k g_{ii}, \quad k \neq i. \\ \Gamma_{ij}^j &= \frac{1}{2} g^{jj} \partial_i g_{jj}, \quad j \neq i. \end{aligned}$$

More precisely, we saw in the proof of Lemma 2.6 that $g_{ij} = \delta_{ij} \cdot \sum_{\nu} \varepsilon_{i\nu} \kappa_{i\nu}^2$, where $\kappa_{i\nu}^2$ is an algebraic function which only depends on the variable x_i , and where $\varepsilon_{i\nu} \in \{\pm 1\}$ is a constant. Thus, $\partial_k g_{ii}$ is zero if $k \neq i$. It follows that Γ_{ii}^i is the only non-vanishing Christoffel symbol. In terms of the Christoffel symbols, the components of the curvature tensor may be written as $R_{ijk}^l = \partial_j \Gamma_{ki}^l - \partial_k \Gamma_{ij}^l + \Gamma_{ki}^r \Gamma_{jr}^l - \Gamma_{kr}^l \Gamma_{ij}^r$. In our setting, $\partial_j \Gamma_{ki}^l = \delta_{ik} \delta_{il} \partial_j \Gamma_{ii}^i = \delta_{ij} \delta_{ik} \delta_{il} \partial_i \Gamma_{ii}^i$, where the last equality is due to the fact that g_{ii} only depends on the variable x_i and that $\Gamma_{ii}^i = \frac{1}{2g_{ii}} \partial_i g_{ii}$. Analogously, $\partial_k \Gamma_{ij}^l = \delta_{ij} \delta_{ik} \delta_{il} \partial_i \Gamma_{ii}^i$, i.e. $\partial_j \Gamma_{ki}^l - \partial_k \Gamma_{ij}^l = 0$. Therefore, it remains to prove that also $\Gamma_{ki}^r \Gamma_{jr}^l - \Gamma_{kr}^l \Gamma_{ij}^r = 0$, where over the index r is summed due to Einstein's convention. Writing this sum explicitly, we have $\sum_r \Gamma_{ki}^r \Gamma_{jr}^l = \sum_r \delta_{rk} \delta_{ir} \delta_{rj} \delta_{lr} (\Gamma_{rr}^r)^2 = \delta_{ik} \delta_{ij} \delta_{il} (\Gamma_{ii}^i)^2$. Similarly, $\sum_r \Gamma_{kr}^l \Gamma_{ij}^r = \sum_r \delta_{rk} \delta_{rl} \delta_{ir} \delta_{rj} \Gamma_{kr}^l \Gamma_{ij}^r = \delta_{ik} \delta_{ij} \delta_{il} (\Gamma_{ii}^i)^2$. This yields $R_{ijk}^l = 0$. $\square$

Let us finally prove the desired result.

2.9 Theorem *Let $X := E_0 \times_S \dots \times_S E_n$ be the fibred product of smooth elliptic curves E_i over S . Let g be the first fundamental form on X (see Proposition 4.21). Then $(X \rightarrow S, g)$ is a model of type (GR).*

PROOF By Corollary 2.8, X is flat, i.e. the curvature tensor vanishes. In particular, the Ricci tensor and the scalar curvature vanish. Therefore, (X, g) is a solution of the

vacuum Einstein equations. Due to the fact that Néron models fulfill the property (ii) of models of type (GR) (see Definition 0.4), it suffices to show that X is the Néron model of its generic fibre. Now notice that the fibred product of Néron models over S is again the Néron model of its generic fibre, because the universal property of fibred products implies the universal property of Néron models (Definition 6.1). Consequently, we are reduced to the proof that an elliptic curve over S is the Néron model of its generic fibre. Thus we are done by Proposition 3.13. $\square$

2.10 Remark Let $X := E_0 \times_S \dots \times_S E_n$ be the fibred product of n smooth elliptic curves E_i over S provided with first fundamental form g . In truth, $(X \rightarrow S, g)$ is even a model of type (SR) (see Definition 0.7). In particular, all results, which are proven in chapter 3, are true for X .

Let us now choose $S = \text{Spec } \mathcal{O}_K$, where $\mathcal{O}_K$ is the ring of integral numbers of an algebraic number field $K \subset \mathbb{R}$. Let us compare X with the Minkowski space-time $\mathbb{A}_S^n$. Both describe a space-time without gravity, because the curvature tensor vanishes identically. Furthermore, both carry a canonical, commutative group structure. In quantum field theory over Minkowski space-time, the inverse of the group law is interpreted as a simultaneous space and time reflection. Consequently, there is also a canonical notion of space and time reflection on X , too. The difference between X and Minkowski space-time is of topological nature. While Minkowski space-time (which is induced by some affine space $\mathbb{A}_S^n$) carries a trivial topology, X does not. In order to illustrate this, let us consider the time coordinates in the respective models. More precisely, let us consider the time coordinate as a complex variable (whose imaginary part represents the physical time and whose real part phenomenologically describes temperature). Then, in the Minkowski case, the time variable runs in the complex plane and therefore on a sphere if we adjoin the point at infinity. In contrast to this, the complex points of the time coordinate E_0 of X yield a torus.

This difference in the global topology, which may not be seen locally, has some interesting consequences with regard to quantum field theory: The set of archimedean points $\mathbb{A}_K^n(K) \cong K^n$ of Minkowski space-time is still a dense subset of the continuum $\mathbb{A}_K^n(\mathbb{R}) \cong \mathbb{R}^n$. In particular, $\mathbb{A}_K^n(K)$ is not a discrete subset of the continuum (it is not even a finitely generated abelian group). On the other hand, there is no discrete subgroup of $\mathbb{A}_K^n(K)$ which is invariant under arbitrary K -isomorphisms of $\mathbb{A}_K^n$. In particular, a discrete subgroup of $\mathbb{A}_K^n(K)$ cannot be Lorentz-invariant.

But, if we consider instead the model X something interesting happens: Due to a theorem of Mordell, the set $X_K(K)$ of archimedean points of X is a finitely generated abelian group. Furthermore, $X_K(K)$ is invariant under arbitrary K -isomorphisms of X_K . Consequently, it makes sense to say that $X_K(K)$ is “Lorentz-invariant”. In the case of rank zero, space-time is actually a discrete set. All in all, $X_K(K)$ may indeed be interpreted as a vacuum, and the methods of lattice gauge theory yield a well defined quantum field theory on $X_K(K)$. This will be exposed in more detail in chapter 5.2.

It would be interesting to consider a dynamical solution $X = E_t \times_S E_1(t) \times_S \dots \times_S E_n(t)$ of the Einstein equations, where E_t is a fixed elliptic curve and where $E_i(t)$ are

elliptic curves described by equations $Y^2Z = X^3 + g_{2,i}(t)XZ^2 + g_{3,i}(t)Z^3$ with coefficients $g_{j,i}(t)$ depending on E_t . For physical reasons, one should expect that the rank of each elliptic curve $E_i(t)$ stays zero for all times t if it was zero for $t = 0$. Therefore, one might conjecture some connections between the rank of elliptic curves and the Einstein equations.

3. Some properties of the arithmetic models

Unless otherwise specified, let $(X \rightarrow S, g)$ be a model of type (SR) (see Definition 0.7), and assume that S is representable by a Dedekind scheme with field of fractions K .

3.1. Etale-invariance

In its differential geometric formulation, general relativity is as well general covariant as diffeomorphism invariant. General covariance means that fundamental physical laws may not depend on the special choice of a local coordinate system and is due to the fact that the Einstein equations are tensor equations. However, diffeomorphism invariance means the following. If g is a solution of the vacuum Einstein equations and if φ is any diffeomorphisms of the space-time manifold, then φ^*g is a vacuum solution, too. Essentially this comes down to saying that $(\varphi^*\Omega)[g] = \Omega[\varphi^*g]$ for all diffeomorphisms φ . Thereby, $\Omega[g]$ denotes the functional $(\text{sc} \cdot \omega)[g] = \text{sc}[g]\omega[g]$, where $\text{sc}[g]$ resp. $\omega[g]$ denote the scalar curvature resp. the volume form corresponding to g .

In our arithmetic approach, the Riemannian space-time manifold is replaced by an appropriate algebraic space. Therefore, it is natural to study in how far general covariance and diffeomorphism invariance are realized for models of type (GR). For this purpose, we first have to find the analogue of the differential geometric diffeomorphism in the realm of schemes. In section 7.1, we will expose in a very detailed way that the class of such coordinate transformations is given by étale morphisms (see Definition 1.11). This naturally implies that the Zariski-topology has to be replaced by the so called étale-topology. This naturally leads to algebraic spaces. Central is the following characterization of étale morphisms which illustrates the formal analogy between étale morphisms and differential geometric diffeomorphisms.

3.1 Proposition *Let $f : X \rightarrow Y$ be an S -morphisms. Let $x \in X$ and set $y := f(x)$. Assume that X is smooth over S at x and that Y is smooth over S at y . Then the following conditions are equivalent:*

- a) *f is étale at x .*
- b) *The canonical homomorphism $(f^*\Omega_{Y/S}^1)_x \rightarrow (\Omega_{X/S}^1)_x$ is bijective.*

PROOF [BLR], Cor. 2.2/10

□

For physical reasons, the number of points of space-time should be invariant under coordinate transformations or “deformations” of space-time. Therefore, it is natural to

restrict attention to the class of finite, étale surjective morphisms $f : X \rightarrow Y$ which have the property that there are bijections $Y_K(K') = X_K(K')$ and $Y(S') = X(S')$ for all étale S -schemes S' with field of fractions K' . Thereby, X_K and Y_K denote as usually the respective generic fibres. However, let us point to the fact that it can happen that f changes the topology.

Now, let us return to the setting of models of type (GR). The following Proposition 3.2 shows that this “adelic”, arithmetic model for gravity is covariant and étale-invariant.

3.2 Proposition *Let $(X \rightarrow S, g)$ be a pair consisting of a smooth S -scheme $X \rightarrow S$ and a metric g on X , such that g is a solution of the Einstein equations 1.35. Let $\varphi : X \rightarrow X$ be an arbitrary étale S -morphism, let $S' \rightarrow S$ be an arbitrary étale S -scheme, and let $(X' \rightarrow S', g')$ be the pair obtained from $(X \rightarrow S, g)$ by base change with $S' \rightarrow S$. Then the following statements hold:*

- 1) *If g is a vacuum solution of the Einstein equations, so is φ^*g .*
- 2) *If $(X \rightarrow S, g)$ is a model of type (GR), so is $(X' \rightarrow S', g')$.*

PROOF It suffices to prove 1) on stalks. Therefore, we may assume that $\varphi : X \rightarrow X$ is given by functions which are algebraic over some polynomial ring such that the corresponding Jacobi matrix is invertible. Thus, we obtain an explicit expression for the components of φ^*g . Furthermore, we may make use of the explicit formulas stated directly above Remark 1.38 and write the Einstein tensor in the well-known form by means of the Christoffel symbols. Then, a very lengthy computation, which is the same as in differential geometry, finally shows that the Einstein tensor of φ^*g indeed vanishes.

Let us now consider assertion 2). Again we may prove locally that g' satisfies the Einstein equations. Due to Proposition 1.6 there is a canonical isomorphism $p^*\Omega_{X/S}^1 \cong \Omega_{X'/S'}^1$, where $p : X' \rightarrow X$ is the canonical projection. Therefore, the components of $g' := p^*g$ are obtained from the components of g by tensoring over $\mathcal{O}_S(S)$ with $\mathcal{O}_{S'}(S') \rightarrow \mathcal{O}_S(S)$, i.e. $g'_{\mu\nu} = g_{\mu\nu} \otimes 1$. Consequently, the Einstein tensor $G'_{\mu\nu}$ of $g'_{\mu\nu}$ vanishes, because $G'_{\mu\nu} = G_{\mu\nu} \otimes 1$.

In order to prove that $(X' \rightarrow S', g')$ is a model of type (GR), it suffices to show that $X' \rightarrow S'$ is a Néron model of its generic fibre, because this implies that it is already a model of type (GR) (apply the universal property of Néron models). But due to [BLR], Prop. 1.2/2 Néron models are stable under étale base change. $\square$

Proposition 3.2 states that models of type (GR) are étale-invariant: As well any étale S -morphism $X \rightarrow X$ (i.e. any deformation of the universe which leaves the shape of the underlying points invariant) as any base change by étale morphisms $S' \rightarrow S$ (i.e. any simultaneous deformation of the points) and consequently any combination of these two operations transforms models of type (GR) into models of type (GR). Therefore, the described physics is invariant as it should be.

Last but not least, let us consider the set $X(S)$ of “adelic” points and the set $X_K(K)$ of archimedean points of a model of type (GR) $(X \rightarrow S, g)$. It makes sense to choose S maximal in the following way: If $\varphi : S' \rightarrow S$ is a finite, étale surjective morphism,

then $S' = S$ and $\varphi = \text{id}$. For example, $S = \text{Spec } \mathbb{Z}$ is maximal in this sense (see [Neu], Kap. III, Thm. 2.18). However, we know from the Yoneda lemma that $X(S)$ is invariant under arbitrary S -isomorphisms $X \rightarrow X$. If S is furthermore maximal in the above sense, $X(S)$ is also invariant under étale base change by finite, étale surjective morphisms.¹ Therefore, Proposition 3.2 tells us that the pair $(X(S), g)$ which is obtained from $(X \rightarrow S, g)$ by evaluation at S -valued points is étale-invariant, too. The same is true for the pair $(X_K(K), g_K)$ which is obtained from $(X \rightarrow S, g)$ by first taking the generic fibre and then evaluating at K -valued points. Thereby, the étale-invariance of $X_K(K)$ follows from the canonical bijection $X_K(K) \cong X(S)$ and from the fact that any K -isomorphism $X_K \rightarrow X_K$ extends uniquely to an S -isomorphism $X \rightarrow X$ (recall that X is the Néron model of its generic fibre X_K). We may summarize as follows:

3.3 Corollary *Let $(X \rightarrow S, g)$ be a model of type (GR). Then the induced pairs*

$$\left(X(S), g \right) \quad \text{and} \quad \left(X_K(K), g_K \right)$$

are étale-invariant. In particular, it makes sense to consider $X(S)$ as “adelic” space-time and $X_K(K)$ as archimedean space-time.

3.2. Gravity

If we do not restrict attention to the archimedean component X_K of X , but instead consider the full “adelic” theory, something interesting may be observed: The space described by $(X \rightarrow S, g)$ cannot be the flat, topologically trivial Minkowski space.

Before, we will prove this interesting fact, let us slightly extend the notion of Néron models. Up to now, we demanded that Néron models are of finite type. In particular, they are quasi-compact, i.e. the inverse image of any affine open subset is quasi-compact as a topological space. But, we may drop the condition that they are of finite type and thus drop the compactness-condition. But due to smoothness, they are still locally of finite type and are called Néron lft-models.

3.4 Definition Let S be a Dedekind scheme with ring of fractions K . Let X_K be a smooth K -scheme. A smooth and separated S -model X is called a Néron lft-model of X_K if X satisfies the Néron mapping property (see Definition 6.1).

In the following, we denote by R a discrete valuation ring with field of fractions K . We denote by R^{sh} the strict henselization of R with field of fractions K^{sh} . Let $\widehat{R}^{\text{sh}}$ be the strict henselization of the completion $\widehat{R}$ of R , and let $\widehat{K}^{\text{sh}}$ be the field of fractions of $\widehat{R}^{\text{sh}}$.

¹Physically, it does not make sense to demand invariance of $X(S)$ under arbitrary étale base-change. For example, if $s \in S$ is a closed point, the open immersion $i : S - \{s\} \rightarrow S$ is étale, but neither finite nor surjective. Performing a base change with i simply means that we throw away the fibre of $X(S)$ over the closed point s . So, it is clear that it does not make sense to demand invariance of $X(S)$ under base change with arbitrary étale morphisms. The physically “correct” morphisms, which should leave $X(S)$ invariant, are covering maps, i.e. finite, étale, surjective maps.

Some parts make use of the notion of excellent rings, whose definition I will recall below. The reader who, is not familiar to this notion, may skip it and simply think of a Dedekind domain A of characteristic zero as an example of an excellent ring (in physical situations we will deal with Dedekind domains of characteristic zero anyway). Let us just mention that the strict henselization of an excellent ring is excellent again. Thus the extensions $\widehat{K}^{sh}/\widehat{K}$ and $\widehat{K}^{sh}/K$ are separable.

Let us first state several existing criterions. Afterwards we will give the physical interpretation.

3.5 Proposition *Let R be a discrete valuation ring with field of fractions K , and let X_K be a smooth commutative K -group scheme of finite type. Then the following conditions are equivalent:*

- a) X_K has a Néron model over R .
- b) $X_K \otimes_K \widehat{K}^{sh}$ contains no subgroup of type $\mathbb{G}_a$ or $\mathbb{G}_m$.
- c) $X_K(\widehat{K}^{sh})$ is bounded in X_K .
- d) $X_K(K^{sh})$ is bounded in X_K .

If, in addition, R is excellent, the above conditions are equivalent to

- e) $X_K \otimes_K K^{sh}$ contains no subgroup of type $\mathbb{G}_a$ or $\mathbb{G}_m$.

PROOF [BLR], Thm. 10.2/1

□

The meaning of the conditions c) and d) will be illustrated in the next section. Here b) or e) are important. If we consider Néron lft-models the existing criterion is as follows.

3.6 Proposition *Let R be a discrete valuation ring with field of fractions K , and let X_K be a smooth commutative K -group scheme of finite type. Then the following conditions are equivalent:*

- a) X_K has a Néron lft-model over R .
- b) $X_K \otimes_K \widehat{K}^{sh}$ contains no subgroup of type $\mathbb{G}_a$.

If, in addition, R is excellent, the above conditions are equivalent to

- c) X_K contains no subgroup of type $\mathbb{G}_a$.

PROOF [BLR], Thm. 10.2/2

□

In physical situation we will always consider an excellent Dedekind scheme S whose field of fractions K has characteristic zero. Then, due to Proposition 3.9, the existence of a global Néron lft-model (resp. of a global Néron model) is equivalent to the existence of the local Néron lft-models (resp. of a global Néron models).

3.7 Theorem *Let S be an excellent Dedekind scheme with field of fractions K with $\text{char}(K) = 0$. Let X_K be a smooth commutative K -group scheme of finite type. Then X_K admits a Néron lft-model over S if and only if X_K contains no subgroup of type $\mathbb{G}_a$.*

We saw that the subgroups of type $\mathbb{G}_a$ are critical. But what is the physics behind these groups? Well, due to the basic principles of general relativity, we always have to consider pairs (X, g) consisting of an object X representing space-time and a metric g on X which encodes gravity (classically X was chosen as a differentiable manifold). More precisely, g is the so called first fundamental form on X (see Proposition 4.21 for the algebraic geometric analogue of this notion). However, in our setting, we have got $X = \mathbb{G}_a$. As a scheme, $\mathbb{G}_a$ coincides with the affine space $\mathbb{A}_{\mathbb{Z}}^1$. Therefore, the first fundamental form on $\mathbb{G}_a^n$ is the trivial metric $\text{diag}(\pm 1, \dots, \pm 1)$. Moreover, the set of differential geometric points represented by $\mathbb{G}_a$ is:

$$\mathbb{G}_a(\mathbb{K}) = \mathbb{K}, \quad \mathbb{K} = \mathbb{R}, \mathbb{C}.$$

Consequently, $\mathbb{G}_a^n$ has to be interpreted as algebraic geometric analogue of the manifold $\mathbb{R}^n$ provided with the trivial metric. In particular, Minkowski-space $\mathbb{R}^4$ corresponds to the $\mathbb{Q}$ -group scheme $\mathbb{G}_{a, \mathbb{Q}}^4$ in our arithmetic approach and thus contains a subgroup $\mathbb{G}_a$. Therefore, Minkowski space is impossible due to Theorem 3.7.

Finally let us give the announced definition of excellent rings and schemes. We will not motivate this notion and refer the reader to [Liu], chapter 8.2, for more details.

3.8 Definition Let A be a noetherian ring. We say that A is *excellent* if it verifies the following three properties:

- a) $\text{Spec } A$ is universally catenary (see [Liu], Definition 8.2.1)
- b) For each $\mathfrak{p} \in \text{Spec } A$ the formal fibres of $A_{\mathfrak{p}}$ are geometrically regular, i.e. for all $x \in A_{\mathfrak{p}}$ the scheme $\text{Spec}(\widehat{A_{\mathfrak{p}}} \otimes_{A_{\mathfrak{p}}} k')$ is regular for all finitely generated field extensions $k'/k(x)$. Thereby $\widehat{A_{\mathfrak{p}}}$ denotes the $\mathfrak{p}$ -adic completion of $A_{\mathfrak{p}}$.
- c) For every finitely generated A -algebra B , the set of regular points of $\text{Spec } B$ is open in $\text{Spec } B$.

3.9 Proposition *Let S be a Dedekind scheme whose field of fractions K has characteristic zero. Then the existence of a global Néron lft-model (resp. of a global Néron model) is equivalent to the existence of the local Néron lft-models (resp. of a global Néron models).*

PROOF [BLR], page 310

□

3.3. The structure of the archimedean component

In this section we are going to analyze the structure of the archimedean limit X_K of X . We will find that there is an algebraic torus T_K and an abelian variety A_K which

is realized as a quotient $A_K = X_K/T_K$. In analogy to the situation of classical, differential geometric gauge theory, this obtrudes the interpretation that *models of type (SR)* describe gravity plus electromagnetism.

The existence of the Néron model X of X_K is equivalent to the existence of the local models $X \times_S \text{Spec } \mathcal{O}_{S,s}$ for each closed $s \in S$ by Proposition 3.9. Due to Proposition 3.6 this is equivalent to the fact that the unipotent radical of X_K is trivial. Then X_K is an extension of an Abelian variety A_K by a torus T_K , i.e. there is an exact sequence of K -group schemes.

$$0 \longrightarrow T_K \longrightarrow X_K \longrightarrow A_K \longrightarrow 0$$

Before interpreting this result physically, let us recall the notions of Abelian varieties and tori.

3.10 Definition Let K be a field. An *Abelian variety* A_K over K is defined to be a K -group scheme which is geometrically integral and proper.² One can show that an Abelian variety is always projective and commutative.

3.11 Definition Let K be a field. An *algebraic torus* T_K over K is a commutative K -group scheme of finite type over K which is isomorphic to $(\mathbb{G}_m)^r$ for some $r \in \mathbb{N}$ over an algebraic closure $\overline{K}$ of K ; i.e.

$$T_{\overline{K}} := T_K \otimes_K \overline{K} \cong (\mathbb{G}_{m,\overline{K}})^r \quad \text{for some } r \in \mathbb{N}.$$

If this isomorphism can be realized over K , i.e. $T_K \cong (\mathbb{G}_{m,K})^r$ for some $r \in \mathbb{N}$, the torus is said to split.

For simplicity, let us assume that we may write X_K as a product $X_K = T_K \times_K A_K$. Then looking at the set of physical points and using the fact that we can choose $\overline{K} = K^{\text{sep}}$ due to $\text{char}(K) = 0$, we obtain an extension of ordinary groups

$$0 \longrightarrow T_K(K^{\text{sep}}) \longrightarrow X_K(K^{\text{sep}}) \longrightarrow A_K(K^{\text{sep}}) \longrightarrow 0.$$

Then, we obtain the decomposition

$$X_K(K^{\text{sep}}) = A_K(K^{\text{sep}}) \times T_K(K^{\text{sep}}) = A_K(K^{\text{sep}}) \times ((\mathbb{G}_{m,K^{\text{sep}}})(K^{\text{sep}}))^r$$

for some $r \in \mathbb{Z}$. Thereby, the last equality is due to the fact that the torus splits over K^{sep} by definition. This isomorphism of groups can be interpreted as follows: The archimedean limit X_K decomposes into two parts. The first part is given by an Abelian variety A_K . This represents the space-time dimensions. The second part is given by a torus T_K and should be interpreted as some kind of internal space (originating from some gauge structure). We will see that T_K is p -adically unbounded.

²A scheme X over S is called *proper*, if the canonical morphism $X \rightarrow S$ is universally closed, separated and of finite type.

On the other hand one can show that each extension of an algebraic variety by a torus admits a Néron lft-model. Thus each model of type (SR) $(X \rightarrow S, g)$ is characterized by a certain structure of its archimedean limit.

3.12 Theorem *Let S be an excellent Dedekind scheme with field of fractions K with $\text{char}(K) = 0$. Let X_K be a smooth commutative K -group scheme of finite type. Then is equivalent:*

- a) X_K admits a Néron lft-model over S .
- b) X_K contains no subgroup of type $\mathbb{G}_a$.
- c) X_K is an extension of an Abelian variety A_K by a torus T_K , i.e. over an algebraic closure of K there is an exact sequence

$$0 \longrightarrow T_K \longrightarrow X_K \longrightarrow A_K \longrightarrow 0.$$

(Later, we will interpret T_K as the gauge group part and A_K as the “space-time” part of X_K (see also Definition 3.28).)

PROOF The equivalence of a) and b) is just Theorem 3.7. The implication a) $\Rightarrow$ c) was illustrated above and c) $\Rightarrow$ a) follows from the following propositions. $\square$

3.13 Proposition *Let S be a connected Dedekind scheme with field of fractions K and let A_K be an Abelian variety over K . Then A_K admits a global Néron model A over S .*

PROOF [BLR], Thm. 1.4/3 $\square$

3.14 Proposition *Let S be a Dedekind scheme with field of fractions K . Then any torus T_K over K admits a Néron lft-model over S .*

PROOF [BLR], Prop. 10.1/6 $\square$

3.15 Proposition *Let $S' \rightarrow S$ be a finite flat extension of Dedekind schemes with fields of fractions K' and K . Let G_K be a smooth K -group scheme and denote by $G_{K'}$ the K' -group scheme obtained by base change. Let H_K be a closed subgroup of G_K which is smooth. Assume that $G_{K'}$ admits a Néron lft-model G' over S' .*

Then the Néron lft-model of H_K over S exists. More precisely it can be constructed as a group smoothening of the schematic closure of H_K in the Weil restriction $\mathfrak{R}_{S'/S}(G')$.

PROOF [BLR], Prop. 10.1/4 $\square$

3.16 Proposition *Let S be a Dedekind scheme with field of fractions K . Let G_K be a smooth connected algebraic K -group which is an extension of a smooth K -group scheme H_K of finite type by a split torus T_K . Assume that $\text{Hom}(H_K, \mathbb{G}_{m,K}) = 0$; for example, the latter is the case if H_K is an extension of an Abelian variety by a unipotent group.*

Then, if H_K admits a Néron lft-model over S , the same is true for G_K .

PROOF [BLR], Prop. 10.1/7 □

Up to now we only considered the archimedean limit X_K of the model of type (SR) $(X \rightarrow S, g)$. But we want to show that also the full “adelic” object X has a very clear structure, too.

So let X_K be the archimedean limit of X . Due to Theorem 3.12 X_K is an extension of an Abelian variety A_K by a torus T_K , i.e. there is exact sequence

$$0 \longrightarrow T_K \longrightarrow X_K \longrightarrow A_K \longrightarrow 0.$$

By Definition 3.11 there exists a finite separable field extension K'/K such that $T_{K'} := T_K \otimes_K K'$ splits. Thus performing a base change with the canonical morphism $\text{Spec } K' \rightarrow \text{Spec } K$, we can replace K by K' in the exact sequence above and assume that T_K splits, say of rank r . But then the extension X_K of A_K by T_K is given by primitive line bundles $\mathcal{L}_1, \dots, \mathcal{L}_r$ on A_K by [Serre2], Chap. VII, n°15, Thm. 5. A line bundle $\mathcal{L}$ on a group scheme G is called primitive if there is an isomorphism

$$m^* \mathcal{L} \cong p_1^* \mathcal{L} \otimes p_2^* \mathcal{L},$$

where m is the group law on G and where $p_i : G \times G \rightarrow G$ are the projections, $i = 1, 2$. Since the local rings of the Néron model A of A_K are factorial, the line bundles $\mathcal{L}_i$, $i = 1, \dots, r$ extend to primitive line bundles on the identity component A^0 of A . Thus they give rise to an exact sequence

$$0 \longrightarrow T^0 \longrightarrow X^0 \longrightarrow A^0 \longrightarrow 0,$$

whose generic fibre is the exact sequence we started with. Furthermore X^0 is the identity component of the Néron lft-model X of X_K .

So the structure of X^0 is clear, if a can analyse the structure of A^0 . But A_K is an Abelian variety, and the structure of A^0 is clarified by the fundamental *Theorem on the potential semi-abelian reduction* of Abelian varieties. Before stating this theorem let us recall the notion of reduction.

Let G be a smooth group scheme of finite type over a connected Dedekind scheme S . We say that G has *abelian reduction* (resp. *semi-abelian reduction*) at a closed point $s \in S$ if the identity component G_s^0 is an Abelian variety (resp. an extension of an Abelian variety by an affine torus). In particular, if G is a Néron model of its generic fibre G_K , where K is the field of fractions of S , we will say that G_K has abelian (resp. semi-abelian) reduction at $s \in S$ if the corresponding fact is true for G . The latter amounts to the same as saying that the local Néron model $G \times_S \text{Spec } \mathcal{O}_{S,s}$ of G_K at $s \in S$ has abelian (resp. semi-abelian) reduction.

If A_K is an abelian variety over K , then A_K is said to have *potential abelian reduction* (resp. *potential semi-abelian reduction*) at a closed point $s \in S$ if there is a finite Galois extension L of K such that A_L has abelian (resp. semi-abelian reduction) at all points

over s . More precisely, we thereby mean that the Néron model A' of A_L over the normalization S' of S in L has abelian (resp. semi-abelian) reduction at all closed points $s' \in S'$ lying over s .

3.17 Theorem *Each abelian variety A_K over K has potential semi-abelian reduction at all closed points of S .*

This theorem clarifies the structure of the model of type (SR) $(X \rightarrow S, g)$. Let us conclude this chapter with a final remark.

3.18 Remark In physical applications, we will consider a pair $(X \rightarrow S, g)$ with $S = \text{Spec } \mathbb{Z}$. Now consider the space-time part A_K of the archimedean component X_K of X . We only know that A_K has potentially semi-abelian reduction, but in general A_K itself will not have semi-abelian reduction. But if we perform the classical continuum limit K (i.e. we consider $X_L := X_K \otimes_K L$ and let the field L tend towards $\overline{\mathbb{Q}}$), then, at some place, A_L happens to have semi-abelian reduction.

The property to have semi-abelian reduction is a very strong symmetry (see Proposition 3.19 or section 3.5). Thus, the “adelic” structure on smallest scales is much richer and less symmetric than the structures that we find in the continuum approximation. For example, we find some kind of entanglement of the dimensions of space-time on smallest scales. The decomposition into distinct dimensions is in general only the consequence of some approximation process (see Remark 3.26 for details).

3.19 Proposition *If an Abelian variety A_K has semi-abelian reduction, then the formation of the identity component of the Néron model of A_K is compatible with faithfully flat extensions of discrete valuation rings $R \hookrightarrow R'$.*

PROOF [BLR], Cor. 7.4/4

□

3.4. Boundedness of the archimedean component

In the previous section we saw that the archimedean limit X_K of X is an extension of an Abelian variety A_K by a torus T_K , i.e. over an algebraic closure of K there is an exact sequence

$$0 \longrightarrow T_K \longrightarrow X_K \longrightarrow A_K \longrightarrow 0$$

(by Theorem 3.12). T_K represents the gauge group part and A_K represents the space-time part of X_K .

Within this section we will prove that the archimedean limit A_K is bounded. More precisely, as well $A_K(K)$ as all non-archimedean continuum limits $A_K(\widehat{K}) = A_K(K_{\mathfrak{p}})$ are bounded with respect to the corresponding $\mathfrak{p}$ -adic norm (recall that $K_{\mathfrak{p}}$ is a finite extension of some $\mathbb{Q}_p$). In the special case $S := \text{Spec } \mathbb{Z}$, the set of all possible norms (up to equivalence of norms) is given by the unique archimedean norm $|\cdot|_{\infty}$ (also called the prime spot at infinity) and the set of non-archimedean norms $|\cdot|_p$. There is one

norm $|\cdot|_p$ for each prime number $p \in \mathbb{Z}$, namely the p -adic norm on $\mathbb{C}_p$. Then the above boundedness assertion states that the set $A_K(\mathbb{Q}_p)$ of points of the p -adic continuum limit of A_K is *necessarily* bounded (with respect to the p -adic norm $|\cdot|_p$ on $\mathbb{C}_p$) for every prime number p of $\mathbb{Z}$. The objective of this section 3.4 is as well the definition of the notion of boundedness as the illustration why this finiteness occurs.

Therefore, let S be the spectrum of the ring of integers of an algebraic number field K . As S is a Dedekind scheme one knows from number theory that the local rings $\mathcal{O}_{S,s}$ for closed points $s \in S$ are discrete valuation rings. Let $\pi \in \mathcal{O}_{S,s}$ be a generator of the maximal ideal $\mathfrak{p}_s$ of $\mathcal{O}_{S,s}$. This prime ideal is called a finite prime. π is called a uniformizing element. Any $r \in \mathcal{O}_{S,s}$ can be uniquely written in the form

$$r = \pi^n \cdot u,$$

with uniquely determined $n \in \mathbb{N}$ and $u \in \mathcal{O}_{S,s}^*$. Therefore each $r \in K := \text{Frac}(\mathcal{O}_{S,s})$ can be uniquely written in the form

$$r = \pi^n \cdot u,$$

where $n \in \mathbb{Z}$ and $u \in \mathcal{O}_{S,s}^*$. Then there is a norm $|\cdot|_{\mathfrak{p}_s}$ on $\text{Frac}(\mathcal{O}_{S,s})$ defined as follows. Let $r \in K := \text{Frac}(\mathcal{O}_{S,s})$ and decompose r as above: $r = \pi^n \cdot u$. Let $p = \text{char } k(s)$ be the characteristic of the residue class field $k(s)$ at s . Then

$$|r|_{\mathfrak{p}_s} := \mathfrak{N}(\mathfrak{p}_s)^{-n}$$

with $\mathfrak{N}(\mathfrak{p}_s) := p^{f_{\mathfrak{p}_s}}$ and $f_{\mathfrak{p}_s} := [k(s) : \mathbb{F}_p]$. This definition may look deterrent, if one is not used to it. $\mathfrak{N}(\mathfrak{p}_s)$ occurs in this definition in order to have certain nice number theoretic identities. The norm $|r|_{\mathfrak{p}_s}$ can be written more “transparently” in the form

$$|r|_{\mathfrak{p}_s} := q^{-n}$$

for some $q \in \mathbb{N}$. The exact value of q is important for number theoretic identities, but does not influence the topology induced by this norm. Thus the size of a number $r \in K$ with respect to the norm $|\cdot|_{\mathfrak{p}_s}$ is only dependent on the number of factors π that are contained in r . The norm $|\cdot|_{\mathfrak{p}_s}$ is a non-archimedian norm.

After these preparations let us introduce the notion of *boundedness*. We start with a discrete valuation ring R with field of fractions K . Furthermore, consider a faithfully flat extension of discrete valuation rings $R \hookrightarrow R'$ and let K' be the field of fractions of R' . In physical situations we will choose an étale morphism $R \hookrightarrow R'$ or we will consider the strict henselization $R' := R^{sh}$ of R . Then, by the above, R and R' give rise to absolute values on K and K' . Assuming that these absolute values coincide on K , we denote them by $|\cdot|$. This is justified by the following proposition.

3.20 Proposition *Let K be a field which is complete with respect to the absolute value $|\cdot|$. Let L/K be an algebraic extension. Then there is a unique continuation of $|\cdot|$ to L .*

If $[L : K] = n < \infty$, then for all $\alpha \in L$

$$|\alpha| = \sqrt[n]{|N_{L/K}(\alpha)|}.$$

PROOF [Neu], Chap. II, Thm. 4.8 □

Now for any K -scheme X_K , for any point $x \in X_K(K')$ corresponding to the maximal ideal $\mathfrak{m}_x$ and for any section g of $\mathcal{O}_{X_K}$ being defined at x , we may view the image $g(x)$ of g in $\mathcal{O}_{X_K, x}/(\mathfrak{m}_x)$ as an element of K' . Thus the absolute value $|g(x)|$ is well defined and it makes sense to say that g is bounded on a subset of $X_K(K')$. As an example let us consider the special case $X_K = \mathbb{A}_K^n$. Then any section g is given by a polynomial $g = g(T_1, \dots, T_n) \in K[T_1, \dots, T_n]$ with variables $T_1, \dots, T_n$. Furthermore, any point $x \in X_K(K') = (K')^n$ can be written as $x = (x_1, \dots, x_n) \in (K')^n$. Thus $g(x)$ is simply the evaluation of the polynomial g at x and $|g(x)|$ is the size of the number $g(x) \in K'$ with respect to the absolute value $|\cdot|$. Everything works like one is used to from the real or complex case, just with the archimedean norm $|\cdot|_\infty$ replaced by its non archimedean counterpart $|\cdot|$.

Applying this procedure to the coordinate functions of the affine n -space $\mathbb{A}_K^n$, we arrive at the notion of bounded subsets of $\mathbb{A}_K^n(K')$.

3.21 Definition As before, let $R \hookrightarrow R'$ be a faithfully flat extension of discrete valuation rings with fields of fractions K and K' . Furthermore, let X_K be a K -scheme of finite type and consider a subset $E \subset X_K(K')$.

- a) If X_K is affine, E is called bounded in X_K if there exists a closed immersion $X_K \hookrightarrow \mathbb{A}_K^n$ mapping E onto a bounded subset of $\mathbb{A}_K^n(K')$.
- b) In the general case, E is called bounded in X_K if there exists a covering of X_K by finitely many affine open subschemes $U_1, \dots, U_s \subset X_K$ as well as a decomposition $E = \bigcup E_i$ into subsets $E_i \subset U_i(K')$ such that, for each i , the set E_i is bounded in U_i in the sense of a).

One can show that condition b) of Definition 3.21 is independent of the particular affine open covering U_i of X_K (see [BLR], Lemma 1.1/3). Furthermore the image of a bounded set is bounded again as one would expect intuitively.

3.22 Proposition *Let $R \hookrightarrow R'$ be a faithfully flat extension of discrete valuation rings with fields of fractions K and K' . Consider a K -morphism $f : X_K \rightarrow Y_K$ between K -schemes of finite type. Then, for any bounded subset $E \subset X_K(K')$, its image under $X_K(K') \rightarrow Y_K(K')$ is bounded in Y_K .*

PROOF [BLR], Prop. 1.1/4 □

3.23 Proposition *Let $R \hookrightarrow R'$ be a faithfully flat extension of discrete valuation rings with fields of fractions K and K' . Consider a proper K -scheme X_K . Then any subset $E \subset X_K(K')$ is bounded in X_K .*

PROOF [BLR], Prop. 1.1/6 □

As A_K is an Abelian variety, it is proper and thus bounded due to Proposition 3.23. Due to our exposition in the previous section 3.3, X^0 is given by the extension

$$0 \longrightarrow T^0 \longrightarrow X^0 \longrightarrow A^0 \longrightarrow 0 ,$$

where A^0 is the identity component of the Néron model A of A_K . Recalling conditions *c*) and *d*) of Proposition 3.5 we see that the boundedness of the extended space-time dimensions is not by chance, but a fundamental property of models of type (SR). The boundedness of the extended space-time dimensions is necessary. In particular, one obtains an effective $\mathfrak{p}$ -adic infrared cutoff.

Last but not least let us consider the “gauge part” of X^0 . Due to the equality $\mathbb{G}_{m,K}(K') = (K')^*$, this is *not* a bounded set. If $\alpha \in (K')^*$ with $|\alpha| > 1$, then two points with coordinates α and α^n are arbitrary far separated from each other with respect to $|\cdot|$ if n is big enough.

3.5. Lifting of structures from the archimedean to the adelic level

Fibre-bundles are an essential tool in real-valued physics (based on differential geometry). In particular, they allow a global, coordinate independent description of physics, e.g. of fields mediating forces. In the framework of vector-bundles, physical fields occur as local sections of vector-bundles with space-time as base manifold M . Let

$$f : N \rightarrow M$$

be the corresponding projection map from the vector-bundle onto the base, whose sections are therefore physical fields.

The notion of principal bundles is furthermore a principal item of gauge theory which we are interested in. Let P be a G -principal bundle over M , where M is a Riemannian manifold and G is a Lie-group. In particular, a principal bundle gives rise to a canonical exact sequence

$$0 \longrightarrow G \longrightarrow P \longrightarrow M \longrightarrow 0$$

and is equipped with a free and transitive G -action ψ on P

$$\psi : P \times G \rightarrow P$$

(see section 8.1). Let us consider the analogues of these structures in our algebraic geometric setting which is exposed in detail in section 8.2. Then the above morphism f corresponds to a *smooth* K -morphism $f_K : X_K \rightarrow Y_K$. The notion of a field (e.g. the electromagnetic field) as section of this morphism makes sense, because locally (with

respect to étale topology) at physical points smooth morphisms admit sections due to Proposition 1.19.

Furthermore, in algebraic geometry, the differential geometric G -principal bundles of gauge theory is the so called torsor under an algebraic K -group G_K (see Definition 8.7). Again there is a canonical exact sequence

$$0 \longrightarrow G_K \longrightarrow P_K \longrightarrow X_K \longrightarrow 0$$

and P_K is equipped with a free and transitive G_K -action ψ_K which is given by an isomorphism

$$\psi_K : P_K \times_{X_K} G_{X_K} \xrightarrow{\sim} P_K \times_{X_K} P_K$$

with $G_{X_K} := G_K \times_K X_K$. But we want to do physics not only in the archimedean limit, but also in the “adelic” situation. In particular, we intend to do gauge theory in this setting. Thus, the above structures like the morphism f_K or the X_K -torsor P_K under G_K (given by the above exact sequence) should lift to the “adelic” level in certain situations.

We will see within this section that this lifting of structures from the archimedean to the “adelic” level in fact happens under quite general assumptions.

Lifting of physical fields Let us consider an archimedean, physical field given by a section $s_K : Y_K \hookrightarrow X_K$ of a smooth K -morphism $f_K : X_K \rightarrow Y_K$ and let X, Y the corresponding “adelic” models of type (SR). Then due to the Néron mapping property there exist unique morphisms $f : X \rightarrow Y$ and $s : Y \rightarrow X$ extending f_K and s_K . By functoriality, we have $f \circ s = \text{id}|_Y$, i.e. s is a section of f and thus a physical field in the “adelic” world.

Lifting of exact sequences There are some quite general situations where exact sequences lift to the “adelic” level.

3.24 Proposition *Let R be a discrete valuation ring with field of fractions K , and let $0 \rightarrow A'_K \rightarrow A_K \rightarrow A''_K \rightarrow 0$ be an exact sequence of Abelian varieties. Consider the associated sequence of Néron models $0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0$. Assume that the following condition is satisfied:*

R has mixed characteristic and the ramification index $e = \nu(p)$ satisfies $e < p - 1$, where p is the residue characteristic of R and where ν is the valuation on R which is normalized by the condition that ν assumes the value 1 at uniformizing elements of R .

Then the following assertions hold:

- a) *If A' has semi-abelian reduction, $A' \rightarrow A$ is a closed immersion.*
- b) *If A has semi-abelian reduction, the sequence $0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0$ is exact.*
- c) *If A has abelian reduction, the sequence $0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0$ is exact and consists of Abelian R -schemes.*

PROOF [BLR], Thm. 7.5/4 □

3.25 Lemma *Let $0 \rightarrow A'_K \rightarrow A_K \rightarrow A''_K \rightarrow 0$ be an exact sequence of Abelian varieties over K . Then A_K has semi-abelian (resp. abelian) reduction if and only if A'_K and A''_K have semi-abelian (resp. abelian) reduction.*

PROOF [BLR], Lemma 7.4/2 □

3.26 Remark Let K be an algebraic number field, and let $S = \text{Spec } \mathcal{O}_K$ be the spectrum of the ring of integral numbers of K . If the archimedean world X_K has no gauge part, X_K is already an Abelian variety. Let us assume that there is an inductive decomposition of X_K into lower dimensional subspaces $A_K^{(i)}$, $i = 1, \dots, n$ which are Abelian varieties, too. More precisely, assume that there are exact sequences of Abelian varieties

$$0 \longrightarrow B_K^{(i)} \longrightarrow B_K^{(i+1)} \longrightarrow A_K^{(i)} \longrightarrow 0, \quad B_K^{(1)} := A_K^{(1)},$$

such that $X_K = B_K^{(n+1)}$. For example, one may assume that each Abelian variety $A_K^{(i)}$ is an elliptic curve. From the physical point of view, the latter yields a decomposition of space-time X_K into the different dimensions. We know that each variety $A_K^{(i)}$ (resp. $B_K^{(i)}$) possesses a Néron model $A^{(i)}$ (resp. $B^{(i)}$). Physically, $A^{(i)}$ (resp. $B^{(i)}$) describes an “adelic” world whose archimedean component is given by $A_K^{(i)}$ (resp. $B_K^{(i)}$). However, in general, the Abelian varieties $A_K^{(i)}$, $B_K^{(i)}$ will not have semi-abelian reduction. Consequently, the above decomposition of X_K will in general not lift to the “adelic” level, i.e. the induced sequences

$$0 \longrightarrow B^{(i)} \longrightarrow B^{(i+1)} \longrightarrow A^{(i)} \longrightarrow 0$$

possibly fail to be exact (see Proposition 3.24). Physically, this defect of exactness may be interpreted as some kind of entanglement of the dimensions on the smallest scales. This is clearly an “adelic” effect. Also with respect to the metric we meet the same phenomenon: At S -valued points, the metric takes values in the ring $\mathcal{O}_K$. But in general, even if $\mathcal{O}_K$ is by chance a principal ideal ring, the metric cannot be diagonalized. Only upon the archimedean component X_K , the metric may be diagonalized, and we find the decomposition of X_K into distinct dimensions (see Proposition 4.13).

However, there is still another phenomenon. If we perform the continuum limit (i.e. we consider $X_L := X_K \otimes_K L$ and the field L tend towards $\overline{\mathbb{Q}}$), then, as soon as L becomes big enough, $A_L^{(i)} := A_K^{(i)} \otimes_K L$ happens to have semi-abelian reduction (Theorem 3.17). If $A'^{(i)}$ (resp. $B'^{(i)}$) denotes the Néron model of $A_L^{(i)}$ (resp. $B_L^{(i)}$) over the normalization S' of S in L , it follows that induced exact sequences

$$0 \longrightarrow B_L^{(i)} \longrightarrow B_L^{(i+1)} \longrightarrow A_L^{(i)} \longrightarrow 0$$

actually lifts to the “adelic” level, because each $A_L^{(i)}$ has semi-abelian reduction (apply Proposition 3.24). More precisely, we obtain exact sequences

$$0 \longrightarrow B'^{(i)} \longrightarrow B'^{(i+1)} \longrightarrow A'^{(i)} \longrightarrow 0$$

on the “adelic” level, too. Therefore, the entanglement of the dimensions on the “adelic” level is lost if we perform the continuum limit. Furthermore, there is suddenly not only compatibility of $X' = A'^{(n)}$ with unramified, but also with *ramified* base change (see Proposition 3.19).

Let us now incorporate gauge theory. Therefore, consider a X -torsor P under a gauge group $G_X := G \times_S X$. We know that the model of type (SR) X is the Néron model of its generic fibre X_K . But in general, G need not be the Néron model of its generic fibre. Physically the latter means that G is the gauge group of a gauge field which is limited to the quantum level. Exactly those gauge fields, which appear as classical fields with unlimited range (like the electromagnetic field), must possess a gauge group which is the Néron model of its generic fibre. Let us now determine all gauge groups with this property. For this purpose assume that G_K is the (commutative) gauge group of a classical gauge field with infinite range. However, there is a canonical gauge theory associated to the trivial X_K -torsor $P_K := X_K \times_{X_K} G_{X_K} = X_K \times_K G_K$ under $G_{X_K} := X_K \times_K G_K$ which fits into an exact sequence $0 \rightarrow G_K \rightarrow P_K \rightarrow X_K \rightarrow 0$ of group schemes. This gauge theory realizes the most simplest vacuum structure which is possible and is therefore usually chosen in physical applications. By assumption, the considered gauge field is classical with infinite range. Therefore, its gauge group possesses a Néron model. Now there is the following result.

3.27 Lemma *Let S be a Dedekind scheme with field of fractions K , and let*

$$0 \longrightarrow G'_K \longrightarrow G_K \longrightarrow G''_K \longrightarrow 0 .$$

be an exact sequence of smooth K -group schemes of finite type (which are not necessarily commutative).

- a) If G_K admits a Néron model over S , the same is true for G'_K .*
- b) If G'_K and G''_K admit a Néron model over S' , the same is true for G_K .*

PROOF [BLR], Prop. 7.5/1

□

Therefore, also P_K possesses a Néron model P . It follows from Theorem 3.12 that P_K also fits into an exact sequence $0 \rightarrow T_K \rightarrow P_K \rightarrow A_K \rightarrow 0$, where T_K is a torus, and where A_K is an Abelian variety. Comparing the two exact sequences it is therefore natural to interpret T_K as gauge group part of P_K . As higher-dimensional tori are products of one-dimensional tori, only the one-dimensional torus is indecomposable, i.e. describing gauge bosons which are elementary particles. Therefore, we make the following definition.

3.28 Definition Let $\mathbb{G}_{m,K}$ be the one-dimensional torus. Then the gauge field associated to the (commutative) gauge group

$$G_K := \mathbb{G}_{m,K}$$

is called the arithmetic *electromagnetic field*.

4. The dimensionality of the arithmetic models

Within this section let us consider a *model of type (GR)* $(X \rightarrow S, g)$, and let X_K be the generic fibre of X (see Definition 0.4). Recall that, unless otherwise specified, $K \subset \mathbb{R}$ is an algebraic number field (i.e. a finite algebraic extension of $\mathbb{Q}$), and that $\mathcal{O}_K$ is the ring of integral numbers of K (i.e. the integral closure of $\mathbb{Z}$ in K). For example, think of $\mathcal{O}_K = \mathbb{Z}$ and $K = \mathbb{Q}$. As already illustrated, we consider a smooth algebraic space X_K over K as the fundamental object underlying the archimedean space-time limit (see Definition 1.10). Then, in the archimedean continuum limit, we obtain a classical differentiable manifold M which is realized as the set of $\mathbb{R}$ -valued points of X_K , i.e. $M = X_K(\mathbb{R})$.

In this chapter we will draw some conclusions from the presence of gravity concerning the dimensionality of space-time. Let us once for all denote the relative dimension of X_K by n . In contrast to relativistic theories over the infinite prime spots $\mathbb{R}$ or $\mathbb{C}$, which may be formulated for arbitrary values of n , it will turn out that the finite prime spots of the algebraic number field K exert a wide influence on the admissible number of dimensions. Already B. Dragovich, I.V. Volovich et al. mentioned some remarkable coincidences between p -adics and dimensionality of space-time (see [Dra], Chap. 5, concluding remarks and [ADFV], end of section 2.1). We will meet this phenomenon in our approach, too.

But first, let us recall that there are a priori two different ways of describing the $\mathbb{R}$ -valued Minkowski-space-time $\mathbb{R}^n$ which underlies the theory of special relativity:

- a) Both, space $\vec{x}$ and time t , are $\mathbb{R}$ -valued variables, i.e. $\vec{x} \in \mathbb{R}^{n-1}$, $t \in \mathbb{R}$. The metric of space-time is given by the Minkowski-metric $g_{\mu\nu} = \text{diag}(1, -1, \dots, -1)$.
- b) Space $\vec{x}$ is a $\mathbb{R}$ -valued variable, but time t is purely imaginary, i.e. $t \in i\mathbb{R}$ (as it is proposed by the principles of quantum mechanics). The metric of space-time is given by the euclidian-metric $g_{\mu\nu} = \text{diag}(1, \dots, 1)$.

However, within the bounds of this chapter, only the second point of view (that time is a purely imaginary variable) will be presented. Assuming that gravity is a field of spin 2, one may prove the following statement:

- In the case b), $n = 4$ and each tangent space is equipped with the structure of a quaternion algebra with “Pauli matrices” as generators.

Let us at least mention another issue. If one admits furthermore that next to the spin-2 field of gravity there is also a metrical spin-3 field, one can show that this implies $n \geq 10$.

In order to be able to talk about gravity in our algebraic geometric setting, we have to introduce the notion of a metric. For details we refer to section 4.3.

4.1. The privileged character of four dimensional spaces

Let us consider our archimedean limit $X_K \rightarrow \text{Spec } K$ of relative dimension n . We assume that gravity is a field of spin-2 metric. Therefore, we may choose a metric

$$g_K : T_{X_K/K} \times_{X_K} T_{X_K/K} \rightarrow \mathbb{A}_{X_K}^1$$

(see Definition 4.20). Thereby, $T_{X_K/K}$ denotes the tangent bundle over X_K , and $\mathbb{A}_{X_K}^1$ denotes the one-dimensional affine space over X_K . For those readers, who are not used to these notions, we may give an equivalent description of the metric in terms of matrices. For simplicity, let us assume that the module of differential forms $\Omega_{X_K/K}^1$ is globally free. If $(\omega^1, \dots, \omega^n)$ is a global base of the differential forms $\Omega_{X_K/K}^1$, then g_K corresponds to a non-degenerate symmetric bilinear form

$$g_K = \sum_{1 \leq \mu, \nu \leq n} g_{\mu\nu} \omega^\mu \otimes \omega^\nu \in \Omega_{X_K/K}^{\otimes 2}(X_K), \quad g_{\mu\nu} \in \mathcal{O}_{X_K}(X_K).$$

Thus, the metric may be interpreted as a matrix $(g_{\mu\nu})$. At this place we will not derive the concrete structure of g_K , we simply work with a given structure. The reader may think that $g_{\mu\nu}$ is stipulated by the equations of general relativity. We interpret the metric as an instrument which enables us to measure space-time distances.

Let us study the metric in more detail. Choose a physical point $x \in X_K$ (i.e. a closed point of X_K). Then x takes values in a finite separable extension $k(x) \subset \overline{\mathbb{Q}}$ of K , where $\overline{\mathbb{Q}} \subset \mathbb{C}$ denotes an algebraic closure of $\mathbb{Q}$. Using the physical interpretation given in section 4.3 we see that the evaluation of g_K at the physical point $x \in X_K$ induces a bilinear form

$$g_K(x) : T_{X_K/K}(x) \times T_{X_K/K}(x) \rightarrow k(x).$$

Due to the equality

$$T_{X_K/K}(x) = \text{Hom}_{k(x)\text{-lin}}(\Omega_{X_K/K}^1 \otimes k(x), k(x)) = \text{Hom}_{k(x)\text{-lin}}(k(x)^n, k(x)) \cong k(x)^n$$

we may write $g_K(x)$ as a bilinear form on the $k(x)$ -vector space $k(x)^n$

$$g_K(x) : k(x)^n \times k(x)^n \rightarrow k(x). \quad (4.1)$$

Let us by some abuse of notation denote the matrix representing the bilinear form $g_K(x)$ by $g_{\mu\nu}(x)$ or simply by $g_{\mu\nu}$. As $\text{char}(k(x)) = 0 \neq 2$, we can equivalently write $g_K(x)$ as a quadratic form

$$g_K(x)(T_1, \dots, T_n) = \sum_{\mu, \nu} g_{\mu\nu}(x) T^\mu T^\nu, \quad g_{\mu\nu}(x) \in k(x).$$

Taking into regard that the metric $g_K(x)$ is an instrument to perform measurements in space-time, we can make a further statement on the structure of $g_K(x)$: As the

building stones of a classical space-time are infinite small zero-dimensional point particles, arbitrary small distances in space-time are realized. Thus tangent-vectors of arbitrary lengths may be constructed in the classical world. In particular, if we submit a measurable length, i.e. a number of $k(x)$, then, at least in a classical world, one can construct a tangent-vector which takes exactly this length. This amounts to saying that $g_K(x)$ is surjective.

Consider the infinite prime spot $\mathbb{R}$ of the rational numbers $\mathbb{Q}$. Performing a base change with $K \hookrightarrow \mathbb{R}$, the archimedean limit X_K with its metric g_K induces a differentiable manifold

$$(X_{\mathbb{R}}(\mathbb{R}), g_{\mathbb{R}}),$$

the space-time manifold of the archimedean continuum limit.

Before we proceed, let us introduce some notation which we will need below. The field K possesses valuations which enable us to measure the “size” of its elements. These valuations are called the prime spots of K (see Definition 4.12). In particular, each prime spot induces a topology on K . However, the topological field K need not be complete with respect to a chosen prime spot $\mathfrak{p}$. As usual, let $K_{\mathfrak{p}}$ denote the completion of K with respect to a prime spot $\mathfrak{p}$. One can prove that either $K_{\mathfrak{p}} = \mathbb{R}$, $K_{\mathfrak{p}} = \mathbb{C}$ or that there exists a prime number $p \in \mathbb{Z}$ such that $K_{\mathfrak{p}}$ is a finite extension of $\mathbb{Q}_p$. The complete fields $K_{\mathfrak{p}}$ are called *local* fields, whereas K is a *global* field. The field $\mathbb{R}$ of real numbers is an ordered field (i.e. given two elements $x, y \in \mathbb{R}$, then either $x \leq y$ or $y \leq x$). In contrast to this, the p -adic fields are not ordered. Therefore, in the case $K_{\mathfrak{p}} \neq \mathbb{R}$, the field $K_{\mathfrak{p}}$ is not ordered: There does not exist a split into positive and negative numbers (i.e. the arrow of time and the split into future and past disappears. In a certain way, there exists only future). This interpretation motivates the following definition.

4.1 Definition Let $\alpha \in K_{\mathfrak{p}}$.

- a) If $K_{\mathfrak{p}} = \mathbb{R}$, α is called *strictly positive* if and only if $\alpha > 0$.
- b) If $K_{\mathfrak{p}} \neq \mathbb{R}$, α is called *strictly positive* if and only if $\alpha \neq 0$.

Finally, let us consider a quadratic form $g_K : K^n \times K^n \rightarrow K$ over K . Evaluation of g_K at pairs $(x, y) \in K_{\mathfrak{p}}^n \times K_{\mathfrak{p}}^n$ induces a quadratic form over $K_{\mathfrak{p}}$ which we denote by $g_{K_{\mathfrak{p}}}$.

The case of euclidian gravity.

Let us now illustrate the privileged character of four-dimensional relativistic models. We will assume that $(X_{\mathbb{R}}(\mathbb{R}), g_{\mathbb{R}})$ describes euclidian gravity. Therefore, in inertial systems, the metric takes the form

$$g_{\mu\nu} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

over the prime spots at infinity. This quadratic form is anisotropic over $\mathbb{R}$ and represents all strictly positive real numbers (see Definition 4.15). These two properties of the relativistic metric $g_{\mu\nu}$ over $\mathbb{R}$ motivate the following definition (where we make use of the notation which was introduced above in the paragraph containing Definition 4.1).

4.2 Definition Let K be either a global or a local field. Let $g_K : K^n \times K^n \rightarrow K$ be a quadratic form over K . Then g_K is called *of type (R)* if and only if there exists a finite prime spot $\mathfrak{p}$ of K , such that:

- a) $g_{K_{\mathfrak{p}}}$ is anisotropic.
- b) $g_{K_{\mathfrak{p}}}$ represents all strictly positive numbers $\alpha \in K_{\mathfrak{p}}$.

If K does not possess a finite prime spot, g_K is called of type (R) if a) and b) are true with $K_{\mathfrak{p}} := K$.

4.3 Remark Let g_K be a quadratic form of type (R).

- a) More intuitively, the two conditions a) and b) of Definition 4.2 may be stated as follows:

- $\|x\|_{\mathfrak{p}}^2 := |g_{K_{\mathfrak{p}}}(x, x)|_{\mathfrak{p}} = 0 \iff x = 0$.
- Classical scales do not have holes!

- b) By the Hasse-Minkowski theorem 4.18, g_K does not represent zero over K .
- c) If $n \geq 4$, then the second condition in Definition 4.2 is actually true for *all* finite prime spots of K (as we will see soon). Especially, if $K \subset \mathbb{R}$, we deduce from the Hasse-Minkowski theorem 4.18 that either g_K or $-g_K$ represents all strictly positive numbers $0 < \alpha \in K$.

This fact justifies the “global” statement: Classical scales do not have holes!

- d) If $n = 4$ and $K = \mathbb{R}$, then $g_K = \text{diag}(1, 1, 1, 1)$, i.e. g_K describes a space-time which is relativistic in the sense of euclidian gravity. Therefore, Definition 4.2 generalizes the notion of a relativistic metric (in the sense of the theory of special relativity) from $\mathbb{R}$ to arbitrary local and global fields.

Now we are prepared to incorporate gravity.

4.4 Definition Let $X_K \rightarrow \text{Spec } K$ be of dimension n . Let g_K be a spin-2 metric on X_K describing gravity.

Then X_K is called *of type (R)* if and only if $g_K(x)$ is of type (R) in the sense of Definition 4.2 for all physical (i.e. closed) points $x \in X_K$.

4.5 Theorem Let $X_K \rightarrow \text{Spec } K$ be of type (R) in the sense of Definition 4.4 and of dimension n . Then the following statements are true:

- $n = 4$.
- In inertial systems, the metric may be written as

$$g_{\mu\nu} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -\varepsilon & 0 & 0 \\ 0 & 0 & -\pi & 0 \\ 0 & 0 & 0 & \varepsilon \cdot \pi \end{pmatrix}$$

with numbers $\varepsilon, \pi \in K^*$ which have the following property:

There exists a finite prime spot $\mathfrak{p}$ of K such that

- (i) π is a uniformizing element of $K_{\mathfrak{p}}$.
- (ii) The reduction of $\varepsilon \bmod \pi$ is not a square (i.e. $\bar{\varepsilon} \notin k_{\mathfrak{p}}^{*2}$).

More precisely, $g_{\mu\nu}$ is the norm form of the non-split quaternion algebra (ε, π) over $K_{\mathfrak{p}}$ (see Definition 4.16).

PROOF In order to perform the proof in an elegant manner, let us introduce some notation. For this purpose let us consider the field $\mathbb{Q}_p$ of p -adic numbers (p being a prime number). Let $a, b \in \mathbb{Q}_p^*$ and put

$$(a, b) = \begin{cases} 1 & \text{if } z^2 - ax^2 - by^2 = 0 \text{ has a solution } (z, x, y) \neq (0, 0, 0) \text{ in } \mathbb{Q}_p^3. \\ -1 & \text{otherwise.} \end{cases}$$

The number $(a, b) \in \{\pm 1\}$ is called the *Hilbert symbol* of a and b relative to $\mathbb{Q}_p$. The number (a, b) does not change when a and b are multiplied by squares; thus we may consider the Hilbert symbol as a map

$$(\cdot, \cdot) : \mathbb{Q}_p^* / \mathbb{Q}_p^{*2} \times \mathbb{Q}_p^* / \mathbb{Q}_p^{*2} \rightarrow \{\pm 1\}.$$

Due to [Serre1], Chap. III, §1, Thm. 1, this map has an interesting property which will be important later: It is bilinear and nondegenerate. The assertion “nondegenerate” has to be understood as follows: Each number $b \in \mathbb{Q}_p^*$ which has the property that $(a, b) = 1$ for all $a \in \mathbb{Q}_p^*$ is a square, i.e. $b \in \mathbb{Q}_p^{*2}$.

If $Q = \sum_{i,j=1}^n a_{ij} X_i X_j$ is a quadratic form in n variables over $\mathbb{Q}_p$, we know from Proposition 4.13 that Q may be diagonalized:

$$Q \sim a_1 X_1^2 + \dots + a_n X_n^2.$$

Then the two elements

$$\begin{aligned} \det(Q) &:= a_1 \cdot \dots \cdot a_n \in \mathbb{Q}_p^* / \mathbb{Q}_p^{*2} \\ s(Q) &:= \prod_{1 \leq i < j \leq n} (a_i, a_j) \in \{\pm 1\} \end{aligned}$$

are important invariants of the quadratic form Q .

After this preparation, let us begin with the proof of Theorem 4.5. Let $x \in X_K$ be a physical point. We may assume that $K = k(x) = \mathbb{Q}$. By assumption, X_K is of type (R), i.e. there exists a prime number p such that on the one hand $g_{\mathbb{Q}_p}(x)$ does not represent zero, but on the other hand represents all non-zero numbers $a \in \mathbb{Q}_p^*$. Let us prove that this already implies $n = 4$. We will do so by showing that dimensions $n \neq 4$ are impossible. As we are free to perform coordinate transformations (i.e. it suffices to consider similarity classes of quadratic forms), all numbers may be viewed as elements of $\mathbb{Q}_p^* / \mathbb{Q}_p^{*2}$. Furthermore, let us use the abbreviation $Q := g_{\mathbb{Q}_p}(x)$.

n=1: By Proposition 4.9 b), Q represents $a \in \mathbb{Q}_p^*$ if and only if $a = \det(Q)$. But $\mathbb{Q}_p^* / \mathbb{Q}_p^{*2} \neq \{1\}$ by Proposition 4.14. In particular, Q does not represent all $a \in \mathbb{Q}_p^*$. Therefore, $n = 1$ is not possible.

n=2: On the one hand, we know from Proposition 4.9 b) that $(a, -\det(Q)) = s(Q)$ for all $a \in \mathbb{Q}_p^* / \mathbb{Q}_p^{*2}$. On the other hand, it follows from Proposition 4.9 a) that $-\det(Q) \neq 1$. But, we saw above that the Hilbert symbol $(\cdot, \cdot)$ is a *nondegenerate* bilinear form. Therefore, it is possible to choose a number $b \in \mathbb{Q}_p^* / \mathbb{Q}_p^{*2}$ such that $(b, -\det(Q)) \neq s(Q)$, and we arrive at a contradiction.

Thus, $n = 2$ is not possible.

n=3: For each $a \in \mathbb{Q}_p^* / \mathbb{Q}_p^{*2}$, we must have either

$$a \neq -\det(Q) \quad \text{or} \quad \left(a = -\det(Q) \text{ and } s(Q) = (-1, -\det(Q)) \right).$$

But on the other hand, we already know that $s(Q) \neq (-1, -\det(Q))$, because otherwise Q would represent zero by Proposition 4.9 a). This implies the contradiction $a \neq -\det(Q)$ for all $a \in \mathbb{Q}_p^* / \mathbb{Q}_p^{*2}$.

Therefore, $n = 3$ is not possible.

$n \geq 5$: By Proposition 4.9 a), the quadratic form Q is isotropic.

Therefore, $n \geq 5$ is not possible either, and it remains the case $n = 4$. Now the theorem follows from Proposition 4.17. $\square$

4.6 Corollary *K -schemes $X_K \rightarrow \text{Spec } K$ of type (R) exist.*

PROOF As in the proof of Theorem 4.5, we may assume that $K = k(x) = \mathbb{Q}$. By Theorem 4.5, we know that X_K has to be four-dimensional. Therefore, Proposition 4.19 shows that K -schemes $X_K \rightarrow \text{Spec } K$ of type (R) actually exist. $\square$

4.7 Corollary *Let $X \rightarrow S$ be a smooth S -scheme of relative dimension n with metric g such that the archimedean limit X_K is of type (R). Then*

$$n = 4.$$

PROOF Due to Theorem 4.5, the relative dimension n_K of X_K is four. But n_K coincides with n by Lemma 4.8. $\square$

Evoking Lemma 1.25 and Lemma 1.26, we finally obtain the following result.

4.8 Lemma *Let S be a Dedekind scheme with field of fractions K . Let $f : X \rightarrow S$ be a smooth morphism of relative dimension n , and assume that X is connected. Let n_K be the relative dimension of the generic fibre $f_K : X_K \rightarrow \text{Spec } K$ of f . Then*

$$n = n_K.$$

PROOF As already stated, this is a direct consequence from Lemma 1.25 and Lemma 1.26. $\square$

4.9 Proposition *Let p be a prime number, and let Q be a quadratic form of rank n on $\mathbb{Q}_p$. Then the following statements are true.*

a) *For Q to represent zero it is necessary and sufficient that*

- (i) $n = 2$ and $\det(Q) = -1$ (in $\mathbb{Q}_p^* / \mathbb{Q}_p^{*2}$).
- (ii) $n = 3$ and $s(Q) = (-1, -\det(Q))$.
- (iii) $n = 4$ and either $\det(Q) \neq 1$ or $(\det(Q) = 1 \text{ and } s(Q) = (-1, -1))$.
- (iv) $n \geq 5$.

b) *Let $a \in \mathbb{Q}_p^*$. In order that Q represents a it is necessary and sufficient that*

- (i) $n = 1$ and $a = \det(Q)$.
- (ii) $n = 2$ and $s(Q) = (a, -\det(Q))$.
- (iii) $n = 3$ and either $a \neq -\det(Q)$ or $(a = -\det(Q) \text{ and } s(Q) = (-1, -\det(Q)))$.
- (iv) $n \geq 4$.

Note that in this statement a and $\det(Q)$ are viewed as elements of $\mathbb{Q}_p^ / \mathbb{Q}_p^{*2}$. Also all equations have to be read in $\mathbb{Q}_p^* / \mathbb{Q}_p^{*2}$; e.g. the inequality $a \neq -\det(Q)$ means that a is not equal to the product of $-\det(Q)$ and a square.*

PROOF [Serre1], Chap. IV, §2, Thm. 6 and the corollary to this theorem. $\square$

4.10 Remark In particular, we see that the tangent space $T_{X_K/K}(x)$ of X_K at a physical point $x \in X_K$ is not only a vector space, but even a quaternion algebra if X_K is of type (R). We may choose a basis of $T_{X_K/K}(x)$ which is actually the basis of a non-split quaternion algebra

$$\begin{aligned} T_{X_K/K}(x) &\cong Ke_0 \oplus Ke_1 \oplus Ke_2 \oplus Ke_3. \\ e_1e_2 &= e_3, \quad e_2e_1 = -e_3, \quad e_0 = 1 \\ e_1^2 &= \varepsilon \cdot 1 = \varepsilon, \quad e_2^2 = \pi \cdot 1 = \pi \end{aligned}$$

Thus, we may write a gauge field $\mathcal{A}(x)$ at x in the general form

$$\mathcal{A}(x) = \mathcal{A}^0 e_0 + \sum_{i=1}^3 \mathcal{A}^i e_i.$$

The elements e_1 , e_2 and e_3 are analogues of the Pauli spin matrices. By means of conjugation in the quaternion algebra (see Definition 4.16), the gauge field $\mathcal{A}(x)$ gives rise to the field $\overline{\mathcal{A}}(x)$,

$$\overline{\mathcal{A}}(x) = \mathcal{A}^0 e_0 - \sum_{i=1}^3 \mathcal{A}^i e_i.$$

We call $\overline{\mathcal{A}}(x)$ the anti-field of $\mathcal{A}(x)$. Considering pairs $(\mathcal{A}, \overline{\mathcal{A}})$, we may write

$$(\mathcal{A}, \overline{\mathcal{A}}) = \mathcal{A}_0 \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + \mathcal{A}_1 \begin{pmatrix} 0 & e_1 \\ -e_1 & 0 \end{pmatrix} + \mathcal{A}_2 \begin{pmatrix} 0 & e_2 \\ -e_2 & 0 \end{pmatrix} + \mathcal{A}_3 \begin{pmatrix} 0 & e_3 \\ -e_3 & 0 \end{pmatrix}.$$

Introducing the gamma matrices

$$\gamma_0 := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \gamma_i := \begin{pmatrix} 0 & e_i \\ -e_i & 0 \end{pmatrix}, \quad \text{for } i = 1, 2, 3,$$

we finally arrive at the expression

$$(\mathcal{A}, \overline{\mathcal{A}}) = \sum_{\mu=0}^3 \mathcal{A}^\mu \gamma_\mu.$$

We claim that the gamma matrices fulfill the relations

$$\gamma_\mu \gamma_\nu + \gamma_\nu \gamma_\mu = 2g_{\mu\nu} \cdot \mathbf{1}$$

Thereby, $\mathbf{1}$ denotes the unit matrix, and $g_{\mu\nu}$ denotes the metric at the physical point $x \in X_K$ (whose concrete form is determined by Theorem 4.5).

PROOF Let $i, j = 1, 2, 3$. Then we derive that

- $\gamma_0 \gamma_0 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = g_{00} \cdot \mathbf{1}$
- $\gamma_0 \gamma_i = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & e_i \\ -e_i & 0 \end{pmatrix} = \begin{pmatrix} -e_i & 0 \\ 0 & e_i \end{pmatrix}$ and $\gamma_i \gamma_0 = \begin{pmatrix} 0 & e_i \\ -e_i & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} e_i & 0 \\ 0 & -e_i \end{pmatrix}.$

Thus, $\gamma_0 \gamma_i + \gamma_i \gamma_0 = 0 = 2g_{0i} \cdot \mathbf{1}$.

- $\gamma_i \gamma_j = \begin{pmatrix} 0 & e_i \\ -e_i & 0 \end{pmatrix} \begin{pmatrix} 0 & e_j \\ -e_j & 0 \end{pmatrix} = \begin{pmatrix} -e_i e_j & 0 \\ 0 & -e_i e_j \end{pmatrix}.$

Thus, $\gamma_i \gamma_j + \gamma_j \gamma_i = -(e_i e_j + e_j e_i) \cdot \mathbf{1}$. But the generators of the quaternion algebra fulfill the following identities:

- $e_1e_2 = -e_2e_1$, $e_1e_3 = -e_1e_1e_2 = -e_1e_2e_1 = -e_3e_1$, $e_2e_3 = e_2e_1e_2 = -e_1e_2e_2 = -e_3e_2$.
- $e_1^2 = \varepsilon = -g_{11}$, $e_2^2 = \pi = -g_{22}$, $e_3^2 = e_1e_2e_1e_2 = -e_1^2e_2^2 = -\varepsilon\pi = -g_{33}$.

This shows that $e_ie_j + e_je_i = -2g_{ij}$. With this, everything is proven. $\square$

Therefore, pairs $(\mathcal{A}, \overline{\mathcal{A}})$ are elements of a Clifford algebra. In particular, we see that the archimedean limit X_K of X has got the property that gauge fields act canonically on spinors, if X_K is of type (R) (in the sense of Definition 4.4). The coupling of gauge fields to spinors via gamma matrices occurs naturally.

4.2. Quadratic forms

Consider the standard situation of a metric g_K living on a smooth K -scheme X_K (see Definition 1.10). In physical applications, the field K will be an algebraic number field, i.e. a finite algebraic extension of the rational numbers $\mathbb{Q}$. As a consequence, the evaluation of g_K at physical (i.e. closed) points of X_K yields quadratic forms over algebraic number fields (see section 4.3). A good understanding of these quadratic forms is therefore a prerequisite for a successful treatment of a theory of gravity. In particular, some properties of algebraic number fields have to be collected which we are going to present now. An important tool for the analysis of algebraic number fields are valuations.

4.11 Definition Let K be a field, and let K^* be the group of units of K . A *valuation* of K is a map $v : K \rightarrow \mathbb{R} \cup \{\infty\}$ which has the following properties:

- a) $v(a) = \infty \Leftrightarrow a = 0$.
- b) $v(ab) = v(a) + v(b)$ for all $a, b \in K^*$.
- c) $v(a + b) \geq \min\{v(a), v(b)\}$ for all $a, b \in K^*$.

4.12 Definition A prime spot $\mathfrak{p}$ of an algebraic number field K is an equivalence class of valuations of K . The non-archimedean equivalence classes are called finite primes, the archimedean ones are called infinite primes. We write $\mathfrak{p} \nmid \infty$ (resp. $\mathfrak{p} \mid \infty$) if $\mathfrak{p}$ is finite (resp. infinite). In the case $\mathfrak{p} \nmid \infty$, we write $\mathfrak{p} \mid p$ if the residue field $\kappa(\mathfrak{p})$ corresponding to $\mathfrak{p}$, is of characteristic p . Recall that

$$\kappa(\mathfrak{p}) := \mathcal{O}_{\mathfrak{p}} / \mathfrak{m}_{\mathfrak{p}},$$

where $\mathcal{O}_{\mathfrak{p}} := \{\alpha \in K \mid \mathfrak{p}(\alpha) \geq 0\}$ and where $\mathfrak{m}_{\mathfrak{p}} \subset \mathcal{O}_{\mathfrak{p}}$ is the maximal ideal.

In analogy to the case of finite prime spots exposed in Lemma 4.27, the infinite prime spots are given by embeddings $\tau : K \hookrightarrow \mathbb{C}$. An infinite prime spot $\mathfrak{p}$ is called real or complex depending on whether the completion $K_{\mathfrak{p}}$ is isomorphic to $\mathbb{R}$ or $\mathbb{C}$, and we define

$$\kappa(\mathfrak{p}) := K_{\mathfrak{p}}.$$

We may associate to each prime spot $\mathfrak{p}$ of K a canonical homomorphism

$$v_{\mathfrak{p}} : K^* \rightarrow \mathbb{R}.$$

If $\mathfrak{p}$ is finite, we define $v_{\mathfrak{p}}$ to be the normed $\mathfrak{p}$ -adic valuation on K given by $v_{\mathfrak{p}}(K^*) = \mathbb{Z}$. If $\mathfrak{p}$ is infinite, we define $v_{\mathfrak{p}}(a) := -\log |\tau a|$, where $\tau : K \rightarrow \mathbb{C}$ is the embedding defining $\mathfrak{p}$.

Let us now return to quadratic forms. A central result is the fact that quadratic forms over fields of characteristic zero may be diagonalized.

4.13 Proposition *Every quadratic module (V, Q) over a field k (i.e. V is a k -vector space with a quadratic form Q on V) has an orthogonal basis.*

PROOF [Serre1], Chap. IV, Thm. 1 □

Therefore let $(g_{\mu\nu}) = \text{diag}(g_{11}, \dots, g_{nn}) =: \langle g_{11}, \dots, g_{nn} \rangle$ be diagonal. As we are working with spin-2 gravity (i.e. with quadratic forms), and as we are only looking up to coordinate transformation (i.e. up to isometry of quadratic modules), we are free to multiply the g_{ii} with elements of $K^{*2} := \{\alpha^2 | \alpha \in K^*\}$. Therefore, it is of interest to understand fields modulo squares.

4.14 Proposition *Let $K_{\mathfrak{p}}$ be a $\mathfrak{p}$ -adic field with residue class field $k_{\mathfrak{p}}$ and prime π .*

a) *If $\text{char}(k_{\mathfrak{p}}) \neq 2$, then $K_{\mathfrak{p}}$ has exactly four square classes*

$$K_{\mathfrak{p}}^*/K_{\mathfrak{p}}^{*2} = \{1, \varepsilon, \pi, \varepsilon\pi\}$$

*where ε is a unit whose reduction mod π is not a square, i.e. $\bar{\varepsilon} \notin k_{\mathfrak{p}}^{*2}$.*

b) *If $\text{char}(k_{\mathfrak{p}}) = 2$ that is $K_{\mathfrak{p}}$ is a finite extension of $\mathbb{Q}_2$, then $K_{\mathfrak{p}}$ has exactly 2^{n+2} square classes for $n = [K_{\mathfrak{p}} : \mathbb{Q}_2]$.*

c) *Up to isomorphism $K_{\mathfrak{p}}$ has exactly one non-split¹ quaternion algebra. If we define the “Hasse-Minkowski symbol” by*

$$s(\alpha, \beta) := \begin{cases} 1 & \text{if } (\alpha, \beta) \text{ splits} \\ -1 & \text{if } (\alpha, \beta) \text{ does not splits} \end{cases}$$

then the map $s : K_{\mathfrak{p}}^/K_{\mathfrak{p}}^{*2} \times K_{\mathfrak{p}}^*/K_{\mathfrak{p}}^{*2} \rightarrow \{\pm 1\}$ is nonsingular. If $\text{char}(k_{\mathfrak{p}}) \neq 2$, then (ε, π) is a non-split quaternion algebra.*

PROOF [Schar], Chap. 6, Fact 4.1 □

¹Let (α, β) be a quaternion algebra (see Definition 4.16). Then the following statements are equivalent:

- (α, β) splits
- (α, β) is not a skew-field.
- The norm form N of (α, β) is isotropic.

Before we state the important Hasse-Minkowski theorem, let us introduce the following notion.

4.15 Definition Let K be a field and let $Q(T_1, \dots, T_n) := \sum_{i,j} a_{ij} T^i T^j$, $a_{ij} \in K$ be a quadratic form in n variables over K .

- a) We say that Q represents an element $a \in K$ if there exists $t \in K^n$, $t \neq 0$, such that $Q(t) = a$.
- b) Q is called *isotropic* if it represents $0 \in K$.
- c) Q is called *anisotropic* if it does not represent $0 \in K$.

4.16 Definition For $a, b \in K^*$ define a 4-dimensional K -algebra with basis $1, e_1, e_2, e_3$ by the following multiplication table:

$$e_1 e_2 = e_3, \quad e_2 e_1 = -e_3, \quad e_1^2 = a \cdot 1 = a, \quad e_2^2 = b \cdot 1 = b$$

where 1 denotes the unit element which is sometimes also denoted by e_0 . The elements of this algebra are linear combinations

$$\alpha_0 e_0 + \alpha_1 e_1 + \alpha_2 e_2 + \alpha_3 e_3, \quad \alpha_i \in K.$$

This algebra is associative. It is denoted by $(a, b) = (a, b)_K$ and is called a quaternion algebra over K . There is a canonical involution

$$- : (a, b) \rightarrow (a, b), \quad x = x^0 e_0 + \sum_{i=1}^3 x^i e_i \mapsto \bar{x} := x^0 e_0 - \sum_{i=1}^3 x^i e_i$$

which is the quaternion analogue of complex conjugation. The quadratic form

$$N : (a, b) \rightarrow K, \quad x \mapsto x\bar{x} = \bar{x}x$$

is called the norm form of the quaternion algebra. There is the following isomorphism of quadratic spaces:

$$\left((a, b), N \right) \cong \langle 1, -a, -b, ab \rangle.$$

4.17 Proposition Let $K_{\mathfrak{p}}$ be a $\mathfrak{p}$ -adic field.

- a) Every form of dimension ≥ 5 is isotropic.
- b) Up to isometry there exists exactly one anisotropic 4-dimensional form, namely the norm form of the non-split quaternion algebra. Using Proposition 4.14, this norm form can be written as $\langle 1, -\varepsilon, -\pi, \varepsilon\pi \rangle$.

Therefore quadratic forms can be classified by dimension, determinant and Hasse-Minkowski symbol $s(\alpha_1, \dots, \alpha_n) := \prod_{i < j} s(\alpha_i, \alpha_j)$.

PROOF [Schar], Chap. 6, Thm. 4.2 □

4.18 Hasse-Minkowski theorem (*Local-global principle for quadratic forms.*) *Let K be a global field of characteristic $\neq 2$ and let Q be a n -dimensional quadratic form over K . Then Q is isotropic if and only if Q is isotropic over all completions $K_{\mathfrak{p}}$ with $\mathfrak{p}$ non-archimedean or real.*

PROOF [Schar], Chap. 6, Main Theorem 4.2 □

4.19 Proposition *Let p be a prime number, let $n \geq 1$, let $d \in \mathbb{Q}_p^* / \mathbb{Q}_p^{*2}$ and let $\varepsilon \in \{\pm 1\}$. In order that there exists a quadratic form Q of rank n such that:*

- $\det(Q) = d$ and
- $s(Q) = \varepsilon$,

it is necessary and sufficient that one of the following conditions is fulfilled:

- (i) $n = 1$ and $\varepsilon = 1$
- (ii) $n = 2$ and $\varepsilon = 1$
- (iii) $n = 2$ and $d \neq -1$
- (iv) $n \geq 3$.

PROOF [Serre1], Chap. IV, Prop. 6 □

4.3. The metric tensor

In Riemannian geometry the metric tensor is a smooth family of scalar products. For each point there is one scalar product, i.e. a non-degenerate, positive-definite symmetric bilinear-form living in the tangent space at the point. We would like to provide this notion in algebraic geometry, too.

As usual let us consider a smooth scheme X over a Dedekind scheme S . Let $\omega \in \Omega_{X/S}^1(X)$ be a global differential form and let $(U_i)_{i \in I}$ an affine open covering of X . Consider the $\mathcal{O}_X(U_i)$ -algebra homomorphisms

$$\begin{aligned} \mathcal{O}_X(U_i)[T] &\rightarrow \operatorname{Sym}_{\mathcal{O}_X(U_i)} \left(\Omega_{X/S}^1(U_i) \right) \\ T &\mapsto \omega|_{U_i} \end{aligned}$$

where “Sym” denotes the associated symmetric algebra. This algebra homomorphism corresponds to a X -morphism

$$\omega_i : T_{U_i/S} \rightarrow \mathbb{A}_{U_i}^1,$$

because by Definition 11.16 and Definition 11.24 one has got

$$T_{U_i/S} = \mathbb{V}(\Omega_{U_i/S}^1) = \text{Spec} \left(\text{Sym}_{\mathcal{O}_X(U_i)} \left(\Omega_{X/S}^1(U_i) \right) \right).$$

$$\mathbb{A}_{U_i}^1 = \text{Spec} \mathcal{O}_X(U_i)[T]$$

$T_{U_i/S}$ is called the algebraic geometric tangent bundle, whose basic properties are summarized in chapter 11 of this thesis. As ω is a global section of $\Omega_{X/S}^1$ the morphisms ω_i glue to a X -morphism

$$\omega : T_{X/S} \rightarrow \mathbb{A}_X^1$$

which we also denote by ω with some abuse of notation. Vice versa each such morphism $\omega : T_{X/S} \rightarrow \mathbb{A}_X^1$ gives rise to a global section ω of $\Omega_{X/S}^1$. This global section is the image of the variable T under the associated algebra homomorphism $\mathcal{O}_X[T] \rightarrow \text{Sym}_{\mathcal{O}_X} \left(\Omega_{X/S}^1 \right)$. Thus each global differential form $\omega \in \Omega_{X/S}^1$ can be interpreted as a X -morphism $\omega : T_{X/S} \rightarrow \mathbb{A}_X^1$.

Analogously we can perform with global sections $\omega \in \Omega_{X/S}^{\otimes n}(X)$. Writing

$$\omega = \sum_{1 \leq i_1, \dots, i_n \leq m} \omega_{i_1} \otimes \dots \otimes \omega_{i_n}, \quad \omega_{i_j} \in \Omega_{X/S}^1(X)$$

as a sum of elementary tensors, then ω corresponds to the “multi-linear” morphism

$$\omega : T_{X/S} \times_X \dots \times_X T_{X/S} \rightarrow \mathbb{A}_X^1, \quad (t_1, \dots, t_n) \mapsto \sum_{1 \leq i_1, \dots, i_n \leq m} \omega_{i_1}(t_1) \cdot \dots \cdot \omega_{i_n}(t_n).$$

More precisely this morphism is obtained as follows. By what we have already seen, each $\omega_{i_j} \in \Omega_{X/S}^1(X)$ can be interpreted as a X -morphism $\omega_{i_j} : T_{X/S} \rightarrow \mathbb{A}_X^1$. Thus ω gives rise to a canonical morphism

$$\underbrace{T_{X/S} \times_X \dots \times_X T_{X/S}}_{n\text{-times}} \rightarrow \left(\mathbb{A}_X^1 \times_X \dots \times_X \mathbb{A}_X^1 \right)^{\oplus m}$$

where both fibre products consist of n factors. Composing first with the n -fold multiplication morphism

$$\left(\mathbb{A}_X^1 \times_X \dots \times_X \mathbb{A}_X^1 \right)^{\oplus m} \xrightarrow{\text{mult.}} \left(\mathbb{A}_X^1 \right)^{\oplus m}$$

and then summing up the m factors

$$\left(\mathbb{A}_X^1 \right)^{\oplus m} \xrightarrow{\Sigma} \mathbb{A}_X^1$$

we finally get the X -morphism ω as the composition of these three canonical morphisms. The multi-linear character of this morphism is illustrated in more detail in the beginning of section 11.3.

Let us now assume that $\Omega_{X/S}^1$ is free $\mathcal{O}_X$ -module. For example, this is the case if X is an S -group scheme and if S is in addition the spectrum of a principal ideal domain (let us remark that this will be the case in physical applications). Then choose a base $(\omega^1, \dots, \omega^n)$ of $\Omega_{X/S}^1(X)$ where n is the relative dimension of X over S . In order to construct a metric we consider an element

$$g = \sum_{1 \leq i, j \leq n} g_{ij} \omega^i \otimes \omega^j \in \Omega_{X/S}^{\otimes 2}(X), \quad g_{ij} \in \mathcal{O}_X(X).$$

By what we have seen above g corresponds to a “bilinear” X -morphism

$$g : T_{X/S} \times_X T_{X/S} \rightarrow \mathbb{A}_X^1.$$

More precisely bilinear means the following. Let $\alpha : X' \rightarrow T_{X/S}$ be a X -morphism. Due to the universal property of the fibre product there is a canonical bijection

$$\mathrm{Hom}_X(X', T_{X/S} \times_X T_{X/S}) \cong \mathrm{Hom}_X(X', T_{X/S}) \times \mathrm{Hom}_X(X', T_{X/S}).$$

Thus g induces a canonical map

$$g(\alpha) : \mathrm{Hom}_X(X', T_{X/S}) \times \mathrm{Hom}_X(X', T_{X/S}) \rightarrow \mathrm{Hom}_X(X', \mathbb{A}_X^1).$$

This map is bilinear. Let us make this more transparent by looking at generic fibres.

Physical interpretation: Let $X \rightarrow S$ be smooth of relative dimension n . Let S be a Dedekind scheme with field of fractions K . Let $g : T_{X/S} \times_X T_{X/S} \rightarrow \mathbb{A}_X^1$ be as above. Performing a base change with the canonical inclusion $\mathrm{Spec} K \hookrightarrow S$ we arrive at the archimedean component of g which is given by the X_K -morphism

$$g_K : (T_{X/S} \otimes_S K) \times_{X_K} (T_{X/S} \otimes_S K) \rightarrow \mathbb{A}_{X_K}^1.$$

Recalling the properties of fibre bundles we know that

$$T_{X/S} \otimes_S K = \mathbb{V}(\Omega_{X/S}^1) \times_X X_K = \mathbb{V}(i_K^* \Omega_{X/S}^1) = \mathbb{V}(\Omega_{X_K/K}^1) = T_{X_K/K}$$

where $i_K : X_K \hookrightarrow X$ is the canonical inclusion. Thereby, the first and last equality is by definition, the second one is due to Proposition 11.18, and the third equality is due to Proposition 1.6. Thus we can write g_K in the more transparent way

$$g_K : T_{X_K/K} \times_{X_K} T_{X_K/K} \rightarrow \mathbb{A}_{X_K}^1.$$

Let us evaluate this morphism at L -valued points, where the field L is a separable extension of K . Because of $\mathbb{A}_{X_K}^1(L) = (\mathbb{A}_{\mathbb{Z}}^1 \times_{\mathbb{Z}} X_K)(L) = L \times X_K(L)$ we have got

$$\begin{array}{ccc} & & L \\ & \nearrow p_1 & \\ T_{X_K/K}(L) \times T_{X_K/K}(L) & \xrightarrow{g_K(L)} & L \times X_K(L) \\ & \searrow p_2 & \\ & & X_K(L) \end{array}$$

where p_1 and p_2 are the projections onto the respective factor. Anticipating the notions and statements immediately following Remark 11.25, we obtain

$$T_{X_K/K}(L) = \bigcup_{x \in X_K(L)} T_{X_K/K}(x) = \bigcup_{x \in X_K(L)} T_{X_K, x} \cong \bigcup_{x \in X_K(L)} L^n.$$

If we choose L as a separable algebraic closure $\overline{K}$ of K , this shows that $T_{X_K/K}(\overline{K})$ is actually the differential geometric tangent space (i.e. the family of tangent spaces indexed by points). There is one tangent space for each physical point. Thus $g_K(\overline{K})$ is a family of bilinear forms; for each physical point there is one bilinear form living on the tangent space of this point. More precisely $p_1 \circ g_K(\overline{K})$ is a bilinear form and $p_2 \circ g_K(\overline{K})$ gives the physical point in whose tangent space the bilinear forms lives.

Later we will consider the above bilinear forms

$$g(\alpha) : \text{Hom}_X(X', T_{X/S}) \times \text{Hom}_X(X', T_{X/S}) \rightarrow \text{Hom}_X(X', \mathbb{A}_X^1).$$

for S -valued points $\alpha \in X(S)$ in order to have the notion of a metric at the “adelic” points of X .

So let us continue to construct the metric. We already saw that the global section

$$g = \sum_{1 \leq \mu, \nu \leq n} g_{\mu\nu} \omega^\mu \otimes \omega^\nu \in \Omega_{X/S}^{\otimes 2}(X), \quad g_{\mu\nu} \in \mathcal{O}_X(X).$$

can be interpreted as a bilinear form. The matrix $(g_{\mu\nu})$ is symmetric if and only if the corresponding X -morphism $g : T_{X/S} \times_X T_{X/S} \rightarrow \mathbb{A}_X^1$ is symmetric, i.e. $g(v, w) = g(w, v)$. In order to say what we mean with a non-degenerated bilinear form, we consider the X -functor $\mathcal{T}_{X/S} := \underline{\text{Hom}}_{\mathcal{O}_X}(\Omega_{X/S}^1, \mathcal{O}_X)$ represented by $T_{X/S}$. The X -morphism g induces a canonical map

$$\mathcal{T}_{X/S} \rightarrow (\mathcal{T}_{X/S})^\vee := \underline{\text{Hom}}_{\mathcal{O}_X}(\mathcal{T}_{X/S}, \mathcal{O}_X) \cong \Omega_{X/S}^1, \quad v \mapsto g(\cdot, v).$$

We say the g is non-degenerate if and only if this map is an isomorphism of X -functors. In particular the matrix $(g_{\mu\nu})$ is invertible, i.e. $\det(g_{\mu\nu}) \in \mathcal{O}_X(X)^*$ if g is non-degenerate. We are now prepared to make the

4.20 Definition Let $X \rightarrow S$ be smooth of relative dimension n such that $\Omega_{X/S}^1$ is a free $\mathcal{O}_X$ -module. Let

$$g = \sum_{1 \leq i, j \leq n} g_{ij} \omega^i \otimes \omega^j \in \Omega_{X/S}^{\otimes 2}(X), \quad g_{ij} \in \mathcal{O}_X(X).$$

where $(\omega^1, \dots, \omega^n)$ is a base of $\Omega_{X/S}^1(X)$. Let

$$g : T_{X/S} \times_X T_{X/S} \rightarrow \mathbb{A}_X^1.$$

the associated bilinear X -morphism as constructed above. Then g is called a *spin-2 metric* if and only if g is symmetric and non-degenerate; i.e. if and only if:

- a) The matrix $(g_{\mu\nu})$ is symmetric.
- b) The canonical morphism $\mathcal{T}_{X/S} \rightarrow (\mathcal{T}_{X/S})^\vee$ is an isomorphism of X -functors.

In particular $\det(g_{\mu\nu}) \in \mathcal{O}_X(X)^*$, i.e. the matrix $(g_{\mu\nu})$ is invertible. We denote the inverse matrix of $(g_{\mu\nu})$ by $(g^{\mu\nu})$.

4.21 Proposition *Let $S = \operatorname{Spec} R$ be an affine base scheme and let $X \rightarrow S$ be a smooth S -scheme of finite type. Then there exists a canonical metric g on X , the so called first fundamental form. Locally, this metric looks as follows:*

*In the special case $X = \mathbb{A}_S^n = \operatorname{Spec} R[T_1, \dots, T_n]$, one has got $\Omega_{X/S}^1 = \bigoplus_{i=1}^n \mathcal{O}_X \cdot dT_i$. Then $(g_0)_{\mu\nu} := \operatorname{diag}(\pm 1, \dots, \pm 1)$ induces a trivial metric g_0 on X . If $j : X \rightarrow \mathbb{A}_S^n$ is a closed S -immersion, we may pull back the trivial metric g_0 via j in order to obtain a metric j^*g on X . The metric j^*g is called the first fundamental form.*

PROOF We may choose a finite open covering (U_i) of X by affine schemes U_i together with closed S -immersions $\iota_i : U_i \hookrightarrow \mathbb{A}_S^n$, i.e. $U_i = V(I_i)$ is the zero set of some ideal $I_i \subset R[T_1, \dots, T_n]$. Let $g_i := \iota_i^* g_0$ be the first fundamental form on U_i (as described above). We will show that we may glue the local metrics g_i along the intersections $U_{ij} := U_i \cap U_j$. Due to Lemma 4.22, we know that $U_{ij} = U_i \times_{\mathbb{A}_S^n} U_j$ as scheme. Let $p_k : U_{ij} \rightarrow U_k$, $k = i, j$, be the canonical projection. Then, $p_i^* g_i = p_i^* \iota_i^* g_0 = (\iota_i \circ p_i)^* g_0 = (\iota_j \circ p_j)^* g_0 = p_j^* g_j$. Interpreting g_i as section $g_i \in \Omega_{X/S}^{\otimes 2}(U_i)$, the equation above may be written as $g_i|_{U_{ij}} = g_j|_{U_{ij}}$. Thus, the local metrics g_i glue to a global metric $g \in \Omega_{X/S}^{\otimes 2}(X)$. This metric is called the first fundamental form. $\square$

4.22 Lemma *Let $S = \operatorname{Spec} R$ be an affine base scheme, and let X be a closed subset of $\mathbb{A}_S^n$. Then there is a unique sheaf of ideals $\mathcal{I}$ such that $V(\mathcal{I})$ realizes X as closed sub-scheme of $\mathbb{A}_S^n$ which is smooth over S .*

PROOF We may assume that X is affine. Let $I, J \subset A := \mathcal{O}_{\mathbb{A}_S^n}(\mathbb{A}_S^n)$ be two ideals such that $V(I)$ and $V(J)$ define two closed sub-scheme structures on X . As $V(I)$ and $V(J)$ coincide as topological spaces, one knows from Hilbert's Nullstellensatz that $\operatorname{rad}(I) = \operatorname{rad}(J)$. Due to smoothness, one has got $\operatorname{rad}(I) = I$. This may be seen as follows: Clearly, $I \subset \operatorname{rad}(I)$. Conversely, let $f \in \operatorname{rad}(I)$ and $\bar{f} \in A/I$ be the residue class of f . There is a $m \in \mathbb{N}$ such that $f^m \in I$, i.e. $\bar{f}^m = 0$. Due to smoothness, A/I has no zero-divisors, and it follows that already $\bar{f} = 0$, i.e. $f \in I$. All in all, we obtain $I = \operatorname{rad}(I) = \operatorname{rad}(J) = J$. $\square$

4.23 Remark In exactly the same way as in differential geometry, we may furthermore define the notion of a covariant derivation. Using as well the metric as the notion of the commutator of vector fields (see Definition 11.41), it makes also sense to talk about torsion free and metric preserving connections. Like in differential geometry, we obtain a unique connection which is torsion free and metric preserving, the Levi-Civita connection. It may be derived in terms of the metric by means of the Koszul formula. All in all, we may introduce the notion of curvature for relative schemes $X \rightarrow S$. This establishes a general relativity for schemes, because we may simply write down the equations of general relativity in our algebraic geometric setting. This is done in section 1.2.

Ricci calculus for schemes.

Let $X \rightarrow S$ be a scheme of relative dimension n with spin-2 metric g . Then, locally, we can perform Ricci calculus in the same manner we are used to from differential geometry. For this purpose choose a local base $(\partial_1, \dots, \partial_n)$ of $\mathcal{T}_{X/S}$ and let $(\omega^1, \dots, \omega^n)$ be a local base of $\Omega_{X/S}^1$ such that both bases are dual to each other. Each local section v of $\mathcal{T}_{X/S}$, i.e. each local vector field can be written uniquely as

$$v = \sum_{\mu} v^{\mu} \partial_{\mu}, \quad \text{with local sections } v^{\mu} \text{ of } \mathcal{O}_X.$$

Let us denote the image of v under the isomorphism $\mathcal{T}_{X/S} \rightarrow (\mathcal{T}_{X/S})^{\vee}$ by

$$g(\cdot, v) = \sum_{\mu} v_{\mu} \omega^{\mu}, \quad \text{with local sections } v_{\mu} \text{ of } \mathcal{O}_X.$$

On the other hand we have got the explicit description $g(\cdot, v) = \sum_{\mu, \nu} g_{\mu\nu} \omega^{\nu}(v) \cdot \omega^{\mu} = \sum_{\mu, \nu} g_{\mu\nu} v^{\nu} \omega^{\mu}$. Therefore

$$g_{\mu\nu} v^{\nu} = v_{\mu}$$

where over indices which occur twice is summed. Thus indices are lowered with $g_{\mu\nu}$. Vice versa indices are lifted with $g^{\mu\nu}$:

$$g^{\mu\nu} v_{\nu} = v^{\mu}.$$

Differential forms with values in Lie-algebras

Let $\pi : X \rightarrow S$ be a smooth connected S -scheme which admits global sections (e.g. a universe) and let $G \rightarrow S$ be a smooth S -group scheme. Let $\mathfrak{g} = \text{Lie}(G/S)(S)$ denote the Lie-algebra of G . We know that we can write as well $\mathfrak{g} = (\varepsilon^* \Omega_{G/S}^1)(S)$. In gauge theory we are interested in differential forms which take values in the Lie-algebra $\mathfrak{g}$. This way the fields corresponding to the so called gauge-bosons are described.

The morphism $\pi : X \rightarrow S$ gives a morphism of sheaves $\pi^{-1} \mathcal{O}_S \rightarrow \mathcal{O}_X$. By our premise π has sections; in particular faithfully flat. Then, $\pi^{-1} \mathcal{O}_S = \mathcal{O}_S \circ \pi$ and $\pi^{-1} \mathcal{O}_S \hookrightarrow \mathcal{O}_X$ is injective. Evaluating at global sections, we get a monomorphism of rings $\mathcal{O}_S(S) \hookrightarrow \mathcal{O}_X(X)$. Let us consider $\mathcal{O}_S(S)$ and $\mathfrak{g}$ as a constant sheaves on X , i.e. for all open subsets $U \subset X$ we set:

$$\begin{aligned} \mathcal{O}_S(S)(U) &:= \mathcal{O}_S(S) \\ \mathfrak{g}(U) &:= \mathfrak{g} \end{aligned}$$

We consider $\mathcal{O}_X$ as a $\mathcal{O}_S(S)$ -algebra in the following way: For all open subsets $U \subset X$ we set

$$\mathcal{O}_S(S)(U) \rightarrow \mathcal{O}_X(U), \quad \alpha \mapsto \alpha|_U$$

where we consider α as a global section of $\mathcal{O}_X$. Thus we can consider $\Omega_{X/S}^1$ as a $\mathcal{O}_S(S)$ -module.

4.24 Definition Let $\pi : X \rightarrow S$ be a smooth connected S -scheme which admits global sections (e.g. a universe) and let $G \rightarrow S$ be a smooth S -group scheme. Let $\mathfrak{g}$ be the Lie-algebra of G . A Lie-algebra valued differential form is a section of the $\mathcal{O}_X$ -module

$$\Omega_{X/S}^1 \otimes_{\mathcal{O}_S(S)} \mathfrak{g}.$$

Regarding Proposition 11.30, we have got

$$\begin{aligned} \Omega_{X/S}^1 \otimes_{\mathcal{O}_S(S)} \mathfrak{g} &= \Omega_{X/S}^1 \otimes_{\mathcal{O}_X} (\mathcal{O}_X \otimes_{\mathcal{O}_S(S)} \mathfrak{g}) \\ &= \underline{\text{Hom}}_{\mathcal{O}_X} (\mathcal{T}_{X/S}, \mathcal{O}_X) \otimes_{\mathcal{O}_X} (\mathcal{O}_X \otimes_{\mathcal{O}_S(S)} \mathfrak{g}) \\ &= \underline{\text{Hom}}_{\mathcal{O}_X} (\mathcal{T}_{X/S}, \mathcal{O}_X \otimes_{\mathcal{O}_S(S)} \mathfrak{g}) \end{aligned}$$

Thus elements of $\Omega_{X/S}^1 \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$ can be evaluated at tangent-vectors and the result is a Lie-algebra valued function. We would like to give an equivalent characterization of global sections of $\Omega_{X/S}^1 \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$ in terms of morphisms of schemes as we did above for ordinary differential forms, i.e. sections of $\Omega_{X/S}^1$. So let us consider the fibre-bundle

$$T_{X/S}^{\mathfrak{g}} := \mathbb{V} \left(\Omega_{X/S}^1 \otimes_{\mathcal{O}_S(S)} \mathfrak{g} \right)$$

and let $\mathcal{T}_{X/S}^{\mathfrak{g}}$ be the associated X -functor. In terms of X -functors we can write

$$\begin{aligned} \mathcal{T}_{X/S}^{\mathfrak{g}} &= \underline{\text{Hom}}_{\mathcal{O}_X} \left(\Omega_{X/S}^1 \otimes_{\mathcal{O}_S(S)} \mathfrak{g}, \mathcal{O}_X \right) \\ &= \underline{\text{Hom}}_{\mathcal{O}_X} \left(\Omega_{X/S}^1, \underline{\text{Hom}}_{\mathcal{O}_X} (\mathcal{O}_X \otimes_{\mathcal{O}_S(S)} \mathfrak{g}, \mathcal{O}_X) \right) \\ &= \underline{\text{Hom}}_{\mathcal{O}_X} \left(\Omega_{X/S}^1, (\mathcal{O}_X \otimes_{\mathcal{O}_S(S)} \mathfrak{g})^{\vee} \right) \end{aligned}$$

where we have used again Proposition 11.30. Simply by replacing $\Omega_{X/S}^1$ through $\Omega_{X/S}^1 \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$ in our former calculations, we see that we can identify global sections of $\Omega_{X/S}^1 \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$, i.e. elements of $\Omega_{X/S}^1(X) \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$, with X -morphisms

$$T_{X/S}^{\mathfrak{g}} \rightarrow \mathbb{A}_{X/S}^1.$$

Analogously we can perform with more general tensors with values in the Lie-algebra, i.e. with global sections $\omega \in \Omega_{X/S}^{\otimes n}(X) \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$. Writing

$$\omega = \sum_{1 \leq i_1, \dots, i_n \leq m} \omega_{i_1} \otimes \dots \otimes \omega_{i_{n-1}} \otimes (\omega_{i_n} \otimes r_{i_n}), \quad \omega_{i_j} \in \Omega_{X/S}^1(X), \quad r_{i_n} \in \mathfrak{g}$$

as a sum of elementary tensors, then ω corresponds to the following “multi-linear” morphism.

$$\begin{aligned} \omega : \underbrace{T_{X/S} \times_X \dots \times_X T_{X/S}}_{(n-1)\text{-times}} \times_X T_{X/S}^{\mathfrak{g}} &\rightarrow \mathbb{A}_X^1 \\ (t_1, \dots, t_n) &\mapsto \sum_{1 \leq i_1, \dots, i_n \leq m} \omega_{i_1}(t_1) \cdot \dots \cdot \omega_{i_{n-1}}(t_{n-1}) \cdot (\omega_{i_n} \otimes r_{i_n})(t_n) \end{aligned}$$

4.25 Remark Let K be local field of characteristic zero. Let X_K be a scheme which is locally of finite type over K . If $\overline{K}$ denotes an algebraic closure of K , the set $X_K(\overline{K})$ may be endowed with an analytic structure. In the case $\overline{K} = \mathbb{C}$, we may consider $X_K(\overline{K})$ as complex analytic space, and in the p -adic case $\overline{K} = \mathbb{C}_p$ we obtain a rigid analytic space.

In some analytic theories, the metric is introduced as a powerful notion which contains further geometric information. This is the motivation to introduce this notion on the algebraic level, too (as presented in section 4.3). If X_K is in addition smooth over K (or more generally reduced), we know from Proposition 4.21 that there exists a metric on X_K , the first fundamental form. So, let us assume that there is a metric $g_K : T_{X_K} \times_{X_K} T_{X_K} \rightarrow \mathbb{A}_{X_K}^1$ on X_K . Then we consider the following category: The objects are pairs (X_K, g_K) and a morphism $(Y_K, h_K) \rightarrow (X_K, g_K)$ is defined as a morphism $f_K : Y_K \rightarrow X_K$ of schemes such that $h_K = f_K^* g_K$. This category might be called the category of *Riemannian schemes* over K (of course, the same definition makes sense if we replace K by an arbitrary base scheme). We may consider the category of smooth schemes (or reduced schemes which are locally of finite type) as a full subcategory of Riemannian schemes by endowing the respective scheme with the first fundamental form as metric.

Let us now evaluate Riemannian schemes (X_K, g_K) at $\overline{K}$ -valued points and study the analytic objects that we obtain this way. In order to do this, we assume that X_K is smooth. As mentioned above, the evaluation at $\overline{K}$ -valued points of a scheme X_K yields a set which is the zero set of some polynomials, and which may be endowed with the structure of an analytic space. But if we start from Riemannian schemes instead, it is possible that the resulting analytic object describes the zero set of functions which are *not* necessarily algebraic but may be transcendent.

4.26 Example Let $K = \mathbb{R}$ and $X_K := \mathbb{A}_K^2$ with coordinates (t, r) . With respect to the global base dt, dr of $\Omega_{X_K/K}^1$ choose the metric $g_K = \begin{pmatrix} p(r)^2 & 0 \\ 0 & p'(r)^2 \end{pmatrix}$ on X_K , where $p(r)$ is a polynomial in r , and $p'(r) := \frac{\partial p(r)}{\partial r}$ denotes the first partial derivative of $p(r)$ with respect to r . Evaluation of the Riemannian scheme (X_K, g_K) at $\overline{K}$ -valued points yields a complex analytic space, whose K -valued points constitute the Riemannian manifold $M = (\mathbb{R}^2, g_{\mu\nu})$ with $g_{\mu\nu} = \text{diag}(p(r)^2, q(r)^2)$. We claim that M is isomorphic (in the category of Riemannian manifolds) to a zero set of non-algebraic, transcendent functions. In order to prove this consider an embedding

$$\phi : \mathbb{R}^2 \rightarrow \mathbb{R}^4, \quad (t, r) \mapsto (t, r, \varphi_r \varphi_t, \varphi_r \psi_t)$$

where $\varphi_r, \varphi_t, \psi_t \in C^\infty(\mathbb{R})$ will be chosen later. Thereby, φ_r only depends on r , and $\varphi_t, \psi_t \in C^\infty(\mathbb{R})$ only depend on t . We want to choose φ_r, φ_t and ψ_t in such a way that $\phi(\mathbb{R}^2)$ endowed with the first fundamental form is isomorphic to M . If we introduce the abbreviations $\dot{\varphi}_t = \frac{\partial \varphi_t}{\partial t}$ and $\varphi'_r = \frac{\partial \varphi_r}{\partial r}$, we obtain the following basis-vectors ∂_t, ∂_r of the

tangent space of $\phi(\mathbb{R}^2)$:

$$\partial_t = \begin{pmatrix} 1 \\ 0 \\ \varphi_r \dot{\varphi}_t \\ \varphi_r \dot{\psi}_t \end{pmatrix} \quad \text{and} \quad \partial_r = \begin{pmatrix} 0 \\ 1 \\ \varphi'_r \varphi_t \\ \varphi'_r \psi_t \end{pmatrix}.$$

Choosing the trivial metric $\text{diag}(1, 1, 1, 1)$ in $\mathbb{R}^4$ and providing $\phi(\mathbb{R}^2)$ with the first fundamental form, we obtain the following differential equations.

$$\begin{aligned} p^2 &= g_{tt} = \varphi_r^2(\dot{\varphi}_t^2 + \dot{\psi}_t^2) \\ 0 &= g_{tr} = \varphi_r \varphi'_r(\varphi_t \dot{\varphi}_t + \psi_t \dot{\psi}_t) \\ p'^2 &= g_{rr} = \varphi_r'^2(\varphi_t^2 + \psi_t^2) \end{aligned}$$

One easily checks that $\varphi_r := p(r)$, $\varphi_t := \sin(t)$ and $\psi_t := \cos(t)$ is a solution of these equations. Thus, if (t, r, y, z) are the variables of $\mathbb{R}^4$, we see that we may realize $M \cong \phi(\mathbb{R}^2)$ as zero set in $\mathbb{R}^4$ of the C^∞ -functions $y - p(r)\sin(t)$ and $z - p(r)\cos(t)$. However, these functions are manifestly not algebraic.

Some background material from algebraic number theory

We already introduced the notion of a valuation of an algebraic number field in Definition 4.11. Important is the following result.

4.27 Lemma *Let $K \hookrightarrow L$ be an algebraic extension of fields. Let v be a valuation on K , and let w be a valuation on L which extends v . Then there exists a K -embedding $\tau : L \rightarrow \overline{K_v}$ such that*

$$w = \bar{v} \circ \tau,$$

where $\bar{v}$ is the canonical continuation of v to an algebraic closure $\overline{K_v}$ of the completion K_v of K with respect to v . Furthermore, if L_w denotes the completion of L with respect to w , one has got

$$L_w = LK_v.$$

PROOF The first statement is [Neu], Chap. II, Fortsetzungssatz 8.1. Therefore, it remains to prove the equality $L_w = LK_v$. As a finite extension of a complete field is complete again (see [Neu], Chap. II, Thm. 4.8), the finite field extension $LK_v \subset L_w$ of K_v is complete with respect to w . On the other hand $L \subset LK_v$, and thus LK_v must already coincide with the completion L_w of L with respect to w . $\square$

Let us recall that a prime spot $\mathfrak{p}$ of an algebraic number field K is an equivalence class of valuations of K . The non-archimedean equivalence classes are called finite primes, the archimedean ones are called infinite primes. We write $\mathfrak{p} \nmid \infty$ (resp. $\mathfrak{p} \mid \infty$) if $\mathfrak{p}$ is finite

(resp. infinite). In the case $\mathfrak{p} \nmid \infty$, we write $\mathfrak{p} \mid p$ if the residue field $\kappa(\mathfrak{p})$ corresponding to $\mathfrak{p}$, is of characteristic p . Recall that

$$\kappa(\mathfrak{p}) := \mathcal{O}_{\mathfrak{p}} / \mathfrak{m}_{\mathfrak{p}},$$

where $\mathcal{O}_{\mathfrak{p}} := \{\alpha \in K \mid \mathfrak{p}(\alpha) \geq 0\}$ and where $\mathfrak{m}_{\mathfrak{p}} \subset \mathcal{O}_{\mathfrak{p}}$ is the maximal ideal. In analogy to the case of finite prime spots exposed in Lemma 4.27, the infinite prime spots are given by embeddings $\tau : K \hookrightarrow \mathbb{C}$. An infinite prime spot $\mathfrak{p}$ is called real or complex depending on whether the completion $K_{\mathfrak{p}}$ is isomorphic to $\mathbb{R}$ or $\mathbb{C}$, and we define

$$\kappa(\mathfrak{p}) := K_{\mathfrak{p}}.$$

We may associate to each prime spot $\mathfrak{p}$ of K a canonical homomorphism

$$v_{\mathfrak{p}} : K^* \rightarrow \mathbb{R}.$$

If $\mathfrak{p}$ is finite, we define $v_{\mathfrak{p}}$ to be the normed $\mathfrak{p}$ -adic valuation on K given by $v_{\mathfrak{p}}(K^*) = \mathbb{Z}$. If $\mathfrak{p}$ is infinite, we define $v_{\mathfrak{p}}(a) := -\log |\tau a|$, where $\tau : K \rightarrow \mathbb{C}$ is the embedding defining $\mathfrak{p}$.

4.28 Definition If L/K is a finite extension of K , let us denote the prime spots of L with $\mathfrak{P}$. We will write $\mathfrak{P} \mid \mathfrak{p}$ if the restriction of $\mathfrak{P}$ to K gives $\mathfrak{p}$. In this situation we define:

a) the ramification index $e_{\mathfrak{P}|\mathfrak{p}}$ and the inertia index $f_{\mathfrak{P}|\mathfrak{p}}$ by:

$$\begin{aligned} e_{\mathfrak{P}|\mathfrak{p}} &:= \begin{cases} (\mathfrak{P}(L^*) : \mathfrak{p}(K^*)) & \text{if } \mathfrak{p} \mid p \\ 1 & \text{if } \mathfrak{p} \mid \infty \end{cases} \\ f_{\mathfrak{P}|\mathfrak{p}} &:= [\kappa(\mathfrak{P}) : \kappa(\mathfrak{p})] \\ f_{\mathfrak{p}} &:= f_{\mathfrak{p}|p} \end{aligned}$$

b) the absolute norm $\mathfrak{N}$ by:

$$\mathfrak{N}(\mathfrak{p}) := \begin{cases} p^{f_{\mathfrak{p}}} & \text{if } \mathfrak{p} \mid p \\ e^{f_{\mathfrak{p}}} & \text{if } \mathfrak{p} \mid \infty \end{cases}$$

where $e = \sum_n 1/n!$ is Euler's number.

c) the $\mathfrak{p}$ -adic norm $|\cdot|_{\mathfrak{p}} : K \rightarrow \mathbb{R}$ by:

$$|a|_{\mathfrak{p}} := \begin{cases} \mathfrak{N}(\mathfrak{p})^{-v_{\mathfrak{p}}(a)} & \text{if } \mathfrak{p} \mid p \text{ and } a \neq 0 \\ 0 & \text{if } \mathfrak{p} \mid p \text{ and } a = 0 \\ |\tau a| & \text{if } \mathfrak{p} \mid \infty \text{ is real} \\ |\tau a|^2 & \text{if } \mathfrak{p} \mid \infty \text{ is complex} \end{cases}$$

4.29 Proposition *For arbitrary prime spots $\mathfrak{P} \mid \mathfrak{p}$ the following identities hold:*

$$\begin{aligned} \sum_{\mathfrak{P} \mid \mathfrak{p}} e_{\mathfrak{P} \mid \mathfrak{p}} f_{\mathfrak{P} \mid \mathfrak{p}} &= \sum_{\mathfrak{P} \mid \mathfrak{p}} [L_{\mathfrak{P}} : K_{\mathfrak{p}}] = [L : K] \\ \mathfrak{N}(\mathfrak{P}) &= \mathfrak{N}(\mathfrak{p})^{f_{\mathfrak{P} \mid \mathfrak{p}}} \\ v_{\mathfrak{P}}(a) &= e_{\mathfrak{P} \mid \mathfrak{p}} v_{\mathfrak{p}}(a) && \text{for } a \in K^*, \\ v_{\mathfrak{p}}(N_{L_{\mathfrak{P}} \mid K_{\mathfrak{p}}}(a)) &= f_{\mathfrak{P} \mid \mathfrak{p}} v_{\mathfrak{P}}(a) && \text{for } a \in L^*, \\ |a|_{\mathfrak{P}} &= |N_{L_{\mathfrak{P}} \mid K_{\mathfrak{p}}}(a)|_{\mathfrak{p}} && \text{for } a \in L^*. \end{aligned}$$

PROOF [Neu], Chap. III, Satz 1.2 □

4.30 Proposition *Let K be an algebraic number field. For all $a \in K^*$ there are only finitely many prime spots $\mathfrak{p}$ such that $|a|_{\mathfrak{p}} \neq 1$. Furthermore,*

$$\prod_{\mathfrak{p}} |a|_{\mathfrak{p}} = 1.$$

PROOF [Neu], Chap. III, Satz 1.3 □

4.4. Metrical fields of higher spin

In general relativity gravity is described by a field of spin 2. In this section we will argue that in principle it is mathematically possible that there also exists a metrical field of spin-3. The resulting model is then at least ten dimensional. The global results of this thesis are formulated in a language which leaves open the possibility to work with pure spin-2 gravity or with a combined spin-2 and spin-3 metrical field. Nevertheless, those parts of this thesis which contain computations in local coordinates will work with pure spin-2 gravity only. Therefore, this section 4.4 is thought as a brief prospect for metrical fields with higher spin and will not be that detailed as the exposition concerning spin-2 gravity in section 4.1.

Let us consider a model of type (GR) $(X \rightarrow S, g)$ of relative dimension n . In the preceding section 4.1 we assumed that gravity is describable by a spin-2 metric

$$g : T_{X/S} \times_X T_{X/S} \rightarrow \mathbb{A}_X^1$$

(see Definition 4.20). If $(\omega^1, \dots, \omega^n)$ is a local base of $\Omega_{X/S}^1$, then g corresponds to a non-degenerate symmetric bilinear form

$$g = \sum_{1 \leq \mu, \nu \leq n} g_{\mu\nu} \omega^\mu \otimes \omega^\nu \in \Omega_{X/S}^{\otimes 2}(X), \quad g_{\mu\nu} \in \mathcal{O}_X(X).$$

and is thus given by a matrix $(g_{\mu\nu})$. This definition is motivated by our intuitive experience that e.g. the length of vector $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ should be measured using the theorem of Pythagoras in order to get a “physically sensible” result:

$$\|x\|_2 := \sqrt{\sum_{i=1}^n x_i^2}.$$

But in principle one could also interpret

$$\|x\|_r := \left(\sum_{i=1}^n x_i^r \right)^{\frac{1}{r}} \quad \text{with } r \in \mathbb{R}, \ 0 < r < \infty,$$

as a length of the vector x as this metric yields the same topology on $\mathbb{R}^n$. As we would interpret the index r as the spin of an corresponding gauge boson, we are reduced to values $r \in \{2, 3, 4, \dots\} \subset \mathbb{N}$. We have to exclude $r = 1$ as we want to relate gravity with curvature. The general form of a the metric in $\mathbb{R}^n$ would then be written as a symmetric form of homogeneous degree r .

$$\|x\|_r^r := \sum_{\mu_1, \dots, \mu_r} g_{\mu_1 \dots \mu_r} \cdot x_{\mu_1} \dots x_{\mu_r} \quad \text{with } r \in \{2, 3, 4, \dots\}.$$

In our setting the spin- r field would be given by a symmetric rank r tensor $g_{\mu_1 \dots \mu_r}$ or more precisely by a morphism

$$g : \underbrace{T_{X/S} \times_X \dots \times_X T_{X/S}}_{r\text{-times}} \rightarrow \mathbb{A}_X^1.$$

Locally this could then be written as

$$g = \sum_{1 \leq \mu_1, \dots, \mu_r \leq n} g_{\mu_1 \dots \mu_r} \omega^{\mu_1} \otimes \dots \otimes \omega^{\mu_r} \in \Omega_{X/S}^{\otimes r}(X), \quad g_{\mu_1, \dots, \mu_r} \in \mathcal{O}_X(X).$$

As the tensor product is associative and commutative for symmetric tensors, we can decompose $g_{\mu_1 \dots \mu_r}$ into a tensor-product of symmetric tensors of smaller rank, e.g.:

$$g_{\mu_1 \dots \mu_i \dots \mu_r} = g_{\mu_1 \dots \mu_i} \otimes g_{\mu_{i+1} \dots \mu_r}.$$

According to this decomposition we would interpret the “spin- r field” as some kind of overlap of a “spin- i field” $g_{\mu_1 \dots \mu_i}$ and a “spin- $(r-i)$ field” $g_{\mu_{i+1} \dots \mu_r}$. As r is restricted to values $r \in \{2, 3, 4, \dots\} \subset \mathbb{N}$ we see that the “spin-2 field” $g_{\mu\nu}$ and the “spin-3 field” $g_{\mu\nu\lambda}$ are the only fields which are indecomposable in this sense. They are irreducible in this regard. Therefore only $g_{\mu\nu}$ and $g_{\mu\nu\lambda}$ can be interpreted as elementary particles.

This means that next to the massless spin-2 graviton $g_{\mu\nu}$ of general relativity there could also exist a spin-3 field $g_{\mu\nu\lambda}$. Due to the principles of quantum field theory, this should be an repulsive force which is limited to small distances if we provide it with

a big mass. In the same way as in the case of the pure spin-2 gravity of the previous section 4.1, let us now analyze the effects of a metrical spin-3 field on the dimensionality of space-time. As we will only work with the spin-2 graviton in the section on gauge theory, we will do this very shortly and not that detailed as in the spin-2 case.

The number-theoretic background for the analysis of a spin-3 field is the theory of cubic forms. Of course the cubic form $g_{\mu\nu\lambda}$ describing the potential of the spin-3 field should be non-singular. But then there is a central result due to D.R. Heath-Brown:

4.31 Proposition *a) Every non-singular cubic form over the rational numbers in at least 10 variables represents zero.*

b) There exist non-singular cubic forms over the rational numbers in 9 variables that do not represent zero.

PROOF For item *a*) see [Hea], and for item *b*) see [Mor]. □

By similar arguments as given in the previous section 4.1 on pure spin-2 gravity, this implies:

4.32 Corollary *Let $X \rightarrow S$ be smooth of relative dimension n . If there is a spin-3 field then $n \geq 10$.*

5. Discreteness of geometry and quantization of gauge fields

Let $(X \rightarrow S, g)$ be a model of type (SR) (see Definition 0.7). The classical choice for the base is $S = \operatorname{Spec} \mathbb{C}$. In this case, $X(S)$ may be considered as a manifold, and the points of $X(S)$ form a continuum. However, as we represent the adelic point of view, we make a different choice which also takes into consideration the finite prime spots. Therefore, from now on, let $K \subset \mathbb{R}$ be an algebraic number field with ring of integral numbers $\mathcal{O}_K$, and let $S = \operatorname{Spec} \mathcal{O}_K$. Recall that $X_K(K)$ denotes the archimedean component of $X(S)$ (with X_K being the generic fibre of $X \rightarrow S$).

Within this chapter, we will finally illustrate that $X(S)$ indeed carries a discrete geometry in the “adelic” case $S = \operatorname{Spec} \mathcal{O}_K$ (in contrast to the situation over the real or complex numbers). Because of the canonical bijection $X_K(K) \cong X(S)$, the same is true for the archimedean component $X_K(K)$. More precisely, both $X_K(K)$ and $X(S)$ are finitely generated abelian groups. If the rank of these groups is zero, they are actually finite groups.

In particular, gauge fields on $X_K(K)$ may be treated by means of the methods of lattice gauge theory. This is particularly interesting, because lattice gauge theory offers the possibility to perform a non-perturbative, well-defined quantization of gauge fields.

5.1. The discrete geometry of the arithmetic models

From now on let us consider a smooth, separated K -scheme X_K of relative dimension n over an algebraic number field K . Let us assume that X_K gives rise to a model of type (SR) $(X \rightarrow S, g)$ (see Definition 0.7). In this section, we will study the archimedean component $X_K(K)$ of X . At first, we will illustrate that $X_K(K)$ indeed carries a discrete structure. The argument is as follows. Due to Theorem 3.12, X_K is an extension of an Abelian variety A_K by a torus T_K . Following the ideas of chapter 3, we interpret the torus as the gauge group part and A_K as the space-time part of X_K . As we are interested in the structure of space-time, we will therefore assume that X_K is an Abelian variety. Therefore, the Mordell-Weil theorem 5.1 tells us that the archimedean limit $X_K(K)$ of $X(S)$ is indeed finitely generated abelian group. Because of the bijection $X_K(K) \cong X(S)$, the same is true for the “adelic” world $X(S)$.

5.1 Theorem *Let K be an algebraic number field and let A_K be an algebraic variety over K . Then the set $A_K(K)$ of K -valued points of A_K is a finitely generated abelian group.*

As usually we will let $\mathbb{Z}$ denote the additive group of integers, and we will let $\mathbb{Z}_m$ denote the cyclic group $\mathbb{Z}/m\mathbb{Z}$ of integers mod m . Then the fundamental theorem of finitely generated abelian groups tells us that $X_K(K)$ looks like

$$X_K(K) \cong \mathbb{Z}^d \oplus \mathbb{Z}_{p_1^{\nu_1}} \oplus \cdots \oplus \mathbb{Z}_{p_s^{\nu_s}}$$

for some prime numbers $p_i \in \mathbb{N}$ and integers $d, s, \nu_i \in \mathbb{N}$. As we are interested in a physical theory which is completely free of infinities, it is appealing to demand that $d = 0$, because in this special case, $X_K(K) \cong X(S)$ is actually a finite group. Consequently, there are only finitely many space-time points.

In order to exploit some more physical properties of $X_K(K)$, let us assume that $X := E_0 \times_S \cdots \times_S E_n$ is the fibred product of n smooth elliptic curves E_i over S endowed with the first fundamental form g . We know from Theorem 2.9 that $(X \rightarrow S, g)$ is a model of type (SR). In the case $d > 0$, the set $X_K(K)$ is already dense in the continuum $X_K(\mathbb{R})$. If $d = 0$, $X_K(K)$ is a group of finite order. In the special case $K = \mathbb{Q}$, the Nagell-Lutz theorem tells us that the coordinates of the points of $X_{\mathbb{Q}}(\mathbb{Q})$ are actually integers. Therefore the generator $\ell := 1 \in \mathcal{O}_K$ of the group $\mathcal{O}_K = \mathbb{Z}$ appears as smallest possible length (Planck length) as one should expect.

Next to the existence of a discrete geometry, $(X \rightarrow S, g)$ sometimes has the interesting property to consists of only finitely many points. We already met this phenomenon in the case that X_K is the product of elliptic curves. Therefore, let us next assume that X_K decomposes into a fibred product

$$X_K = C_K^{(1)} \times_K \cdots \times_K C_K^{(n)}$$

of one-dimensional projective, smooth curves $C_K^{(i)}$, $i = 1, \dots, n$. For example, in the case $n = 4$, think of $C_K^{(1)}$ as time coordinate and think of $C_K^{(2)}, C_K^{(3)}$ and $C_K^{(4)}$ as space coordinates. First notice that the genus of the curves cannot be smaller than 1. This may be seen as follows: Assume that the genus of a curve $C_K^{(i)}$ is smaller than one for some i . However, $C_K^{(i)}$ is present on the level of archimedean space-time if and only if it contains a non-virtual point, i.e. if and only if $C_K^{(i)}(K) \neq \emptyset$. Then Proposition 5.3 implies that $C_K^{(i)}$ is already the one-dimensional projective space $\mathbb{P}_K^1$ over K . Thereby, we also made use of the fact that the arithmetic and the geometric genus of the curve $C_K^{(i)}$ agree:

5.2 Remark Let C be a smooth, geometrically connected, projective curve over a field k . Let us remind the reader that

$$p_a(C) := \dim_k H^1(C, \mathcal{O}_C)$$

is called the *arithmetic* genus, and that

$$p_g(C) := \dim_k H^0(C, \Omega_{C/k}^1)$$

is called the *geometric* genus of C . Then one can prove that $p_a(C) = p_g(C)$.

PROOF [Liu], Rem. 7.3.28 □

5.3 Proposition *Let C be a geometrically integral projective curve over a field k of arithmetic genus $p_a \leq 0$. Then we have the following properties:*

- a) *The curve C is a smooth conic over k , i.e. C is smooth over k and there exists a homogeneous polynomial of degree 2 such that $C = V_+(F) \subset \mathbb{P}_k^2$.*
- b) *We have $C \cong \mathbb{P}_k^1$ if and only if $C(k) \neq \emptyset$.*

PROOF [Liu], Prop. 7.4.1 □

But the projective spaces $\mathbb{P}_K^r$, $r \in \mathbb{N}$, cannot be the archimedean component of a model of type (GR), because not every K -automorphism of $\mathbb{P}_K^r$ extends to an S -automorphism of $\mathbb{P}_S^r$. For instance consider the following counterexample:

5.4 Example Let R be a discrete valuation ring with field of fractions K . Consider a K -automorphism $u_K : \mathbb{P}_K^r \xrightarrow{\sim} \mathbb{P}_K^r$. Using a set of homogeneous coordinates $x_0, \dots, x_r$ of $\mathbb{P}_K^r$, we can describe u_K by

$$x_i \mapsto \sum_{j=0}^r a_{ij} x_j, \quad i = 0, \dots, r,$$

where $A := (a_{ij})$ is a matrix in $\mathrm{Gl}_{r+1}(K)$. We may assume that all coefficients a_{ij} belong to R . Then, by the theory of elementary divisors, there are matrices $S, T \in \mathrm{Gl}_{r+1}(K)$ and integers $0 \leq n_0 \leq \dots \leq n_r$ such that SAT is the diagonal matrix $\mathrm{diag}(\pi^{n_0}, \dots, \pi^{n_r})$. Hence there exist sets of homogenous coordinates $x_0, \dots, x_r$ and $x'_0, \dots, x'_r$ of $\mathbb{P}_R^r$, such that u_K is described by

$$x_i \mapsto \pi^{n_i} x'_i,$$

where we may assume $n_0 = 0$.

If $n_0 = \dots = n_r = 0$, it is clear that $u_K : \mathbb{P}_K^r \xrightarrow{\sim} \mathbb{P}_K^r$ extends to an automorphism $u : \mathbb{P}_R^r \xrightarrow{\sim} \mathbb{P}_R^r$. However, if $n_0 = \dots = n_s = 0$ and $n_{s+1}, \dots, n_r > 0$ for some $s < r$, then u_K extends only to an R -rational map $u : \mathbb{P}_R^r \dashrightarrow \mathbb{P}_R^r$. Namely, u is defined on the R -dense open subscheme $V \subset \mathbb{P}_R^r$ which consists of the generic fibre and of the open part $V_k \subset \mathbb{P}_k^r$ complementary to the linear subspace Q_k where $x_0, \dots, x_s$ vanish. In fact, if Q'_k is the linear subspace in $\mathbb{P}_k^r$ where $x'_{s+1}, \dots, x'_r$ vanish, we can view u_k as projection of $\mathbb{P}_k^r$ to Q'_k with center Q_k .

Therefore the projective spaces $\mathbb{P}_K^r$, $r \in \mathbb{N}$, cannot occur. This reflects the fact that models of type (GR) cannot be flat spaces with trivial topology, and may be regarded as projective version of the fact that e.g. Néron models cannot contain affine spaces (Proposition 3.6).

Therefore, the genus of the curves $C_K^{(i)}$ is bigger than zero. This is the case of a non-trivial vacuum topology. In the case of genus bigger than one, we may finally evoke Falting's theorem:

5.5 Faltings's theorem *Any non-singular projective curve of genus $g > 1$ defined over an algebraic number field K contains only finitely many K -valued points.*

All in all, in the case of curves $C_K^{(i)}$ of genus bigger than one, we see that

$$X_K(K) = C_K^{(1)}(K) \times \dots \times C_K^{(n)}(K)$$

is a finite set. Only in the case of genus one, it is still possible that $X_K(K)$ contains infinitely many points. However, we have furthermore proved the following result.

5.6 Non-trivial vacuum structure. *Let us assume that the archimedean limit X_K decomposes into a product*

$$X_K = C_K^{(1)} \times_K \dots \times_K C_K^{(n)}$$

of one-dimensional projective, smooth curves $C_K^{(i)}$, $i = 1, \dots, n$. Then the genus of the curves is strictly bigger than zero, i.e. X_K carries a non-trivial topology. Due to Corollary 2.8, the curvature tensor of X_K vanishes. Therefore, X_K is a vacuum which carries a non-trivial topology.

The next step is the incorporation of gauge fields.

5.2. Lattice gauge theory in the arithmetic setting

In the regime of high energy physics, gravity may be neglected. Therefore, one usually chooses Minkowski space $\mathbb{R}^n$ as space-time manifold and considers gauge fields upon $\mathbb{R}^n$. In a next step one has to quantize these gauge field. Essentially, there are two different approaches. First, there is a perturbative approach based on Feynman diagrams. Second, there is the non-perturbative approach of lattice gauge theory. The first one is manifestly Lorentz-invariant, but in order to quantize the gauge fields (e.g. by means of the Fadeev-Popov method) one has to make a gauge fixing. This breaks the gauge invariance and one has to make sure that physical entities are independent from the chosen gauge. In contrast to this, lattice gauge theory conserves gauge invariance. But one replaces the continuum of Minkowski space by a lattice, and this way Lorentz invariance is broken.

However, Minkowski space-time is not the only solution of Einstein's field equations of general relativity which describes an empty, static universe. Every direct product of one dimensional manifolds is as well a vacuum solution, because the curvature tensor vanishes globally. Nevertheless the different vacua will in general differ with respect to their topological structure. While Minkowski space-time is the product of curves of genus zero (and therefore of trivial vacuum structure), it is as well admissible to consider a product of curves of higher genus. In the case of genus one, we obtain a product of elliptic curves.

In this section we will illustrate that a space-time manifold consisting of a product of elliptic curves (or more generally an Abelian variety) possesses a canonical "lattice" which

is even invariant under algebraic diffeomorphisms (see Definition 5.7, b)). Therefore, in this case, lattice gauge theory offers a possibility to quantize gauge field in such a way that neither gauge invariance nor space-time symmetries are broken.

From now on, we will provide as well the flat space $\mathbb{R}^n$ as the affine space $\mathbb{A}_K^n$ with the euclidean metric $\text{diag}(1, \dots, 1)$, i.e. instead of working with the hyperbolic Minkowski space-time we will consider an euclidean space-time. This is usually done in lattice gauge theory. Therefore, from now on, all manifolds will be considered as Riemannian manifolds.

5.7 Definition a) Let M be a differentiable manifold and $i : \Gamma \hookrightarrow M$ a subset (i denotes the canonical inclusion). Let D be a set of diffeomorphisms of M . Then Γ is called *invariant under D* if the following holds: For every $\varphi \in D$ there is a bijection $\varphi|_\Gamma : \Gamma \xrightarrow{\sim} \Gamma$ of sets such that the following diagram is commutative.

$$\begin{array}{ccc} \Gamma & \xrightarrow{\varphi|_\Gamma} & \Gamma \\ \downarrow i & & \downarrow i \\ M & \xrightarrow{\varphi} & M \end{array}$$

b) Let $K \subset \mathbb{R}$ be an algebraic number field and let X_K be a smooth K -scheme. Let $i : \Gamma \hookrightarrow X_K(\mathbb{R})$ be a subset (i denotes the canonical inclusion). Then Γ is called *K -isomorphism invariant* if the following holds: For every K -isomorphism $\varphi : X_K \rightarrow X_K$ there is a bijection $\varphi(\mathbb{R})|_\Gamma : \Gamma \xrightarrow{\sim} \Gamma$ of sets such that the following diagram is commutative.

$$\begin{array}{ccc} \Gamma & \xrightarrow{\varphi(\mathbb{R})|_\Gamma} & \Gamma \\ \downarrow i & & \downarrow i \\ X_K(\mathbb{R}) & \xrightarrow{\varphi(\mathbb{R})} & X_K(\mathbb{R}) \end{array}$$

Thereby, $\varphi(\mathbb{R}) : X_K(\mathbb{R}) \rightarrow X_K(\mathbb{R})$ denotes the morphism which is obtained by evaluation of φ at $\mathbb{R}$ -valued points.

Due to the smoothness of X_K over K , we may endow $X_K(\mathbb{R})$ with the structure of a differentiable manifold. Thus, if Γ is a K -isomorphism invariant subset, then it is invariant under the set of diffeomorphisms $\varphi(\mathbb{R}) : X_K(\mathbb{R}) \rightarrow X_K(\mathbb{R})$ of $X_K(\mathbb{R})$ which are induced by isomorphisms $\varphi : X_K \rightarrow X_K$ of schemes.

In the special case of the flat space-time $\mathbb{R}^n$ we may consider the set of Lorentz-transformations. A Lorentz-transformation is linear diffeomorphism which describes a rotation in $\mathbb{R}^n$. Each Lorentz-transformation may be written as a finite product of matrices

of the following type.

$$L_\theta := \begin{pmatrix} \ddots & & & & & & \\ & 1 & 0 & 0 & 0 & 0 & 0 \\ & 0 & 1 & 0 & 0 & 0 & 0 \\ & 0 & 0 & \cos(\theta) & \sin(\theta) & 0 & 0 \\ & 0 & 0 & -\sin(\theta) & \cos(\theta) & 0 & 0 \\ & 0 & 0 & 0 & 0 & 1 & 0 \\ & 0 & 0 & 0 & 0 & 0 & 1 \\ & & & & & & \ddots \end{pmatrix}$$

Let D_{Lor} denote the set of all Lorentz-transformations. A subset $\Gamma \subset \mathbb{R}^n$ is called *Lorentz invariant* if it is invariant under D_{Lor} in the sense of Definition 5.7, a). However, whereas the notion of Lorentz invariance is especially adapted to $\mathbb{R}^n$, the notion of K -isomorphism invariance applies to every scheme X_K over K . But in the special case of $\mathbb{R}^n$ one can show that K -isomorphism invariance already implies Lorentz-invariance for a quite big class of sets Γ (see Proposition 5.8). Therefore, in the setting of non-trivial manifolds $X_K(\mathbb{R}) \neq \mathbb{R}^n$, where the notion of Lorentz-invariance does in general not make sense, it is natural to look for K -isomorphism invariant structures.

5.8 Proposition *Let $X_K = \mathbb{A}_K^n$ and consider $X_K(\mathbb{R}) \cong \mathbb{R}^n$ as flat manifold in a canonical way. Let $\Gamma \subset X_K(\mathbb{R})$ be a closed or an open subset. If Γ is K -isomorphism invariant, it is Lorentz-invariant, too.*

PROOF Let us first assume that $\Gamma \subset X_K(\mathbb{R})$ is a closed subset. Choose $x \in \Gamma$. As every Lorentz transformation may be written as a finite product of matrices L_θ , it suffices to show, that $L_\theta(x) \in \Gamma$ for all the Lorentz transformations of type L_θ . Thus it remains to show that $\inf_{\gamma \in \Gamma} |L_\theta(x) - \gamma| = 0$, because $\Gamma \subset X_K(\mathbb{R})$ is a closed subset. Thereby, $|\cdot|$ denotes the canonical norm on $\mathbb{R}^n$.

In order to prove this consider the sphere of radius one $S^1 \subset \mathbb{R}^2$. For every $z \in S^1$ there is a $\theta \in [0, 2\pi]$ such that $z = (\cos(\theta), \sin(\theta)) \in \mathbb{R}^2$. By Lemma 5.9, we know that the set of K -valued points of S^1 is dense in S^1 (with respect to the canonical topology on S^1). Consequently, we may choose a point $z_\varepsilon = (\cos(\theta_\varepsilon), \sin(\theta_\varepsilon)) \in K^2$ such that $|\cos(\theta) - \cos(\theta_\varepsilon)| < \varepsilon$ and $|\sin(\theta) - \sin(\theta_\varepsilon)| < \varepsilon$ for every $\varepsilon > 0$. In particular,

$$|L_\theta(x) - L_{\theta_\varepsilon}(x)| \leq \sqrt{2}\varepsilon|x| \quad \text{for all } \varepsilon > 0.$$

But each Lorentz transformations L_{θ_ε} is induced by a K -isomorphisms of X_K , because the matrix L_{θ_ε} has only entries in K . Thus $L_{\theta_\varepsilon}(x) \in \Gamma$ due to our assumption, and it follows that $\inf_{\gamma \in \Gamma} |L_\theta(x) - \gamma| = 0$.

Let us now assume that $\Gamma \subset X_K(K)$ is open. Then the complement $\Gamma^c := X_K(K) \setminus \Gamma$ of Γ in $X_K(K)$ is a closed subset. Let us first show that Γ^c is K -isomorphism invariant. In order to do this, assume that there is a K -isomorphism $\varphi : X_K \rightarrow X_K$ and a $y \in \Gamma^c$ such that $x := \varphi(y) \in \Gamma$. But then $\varphi^{-1}(x) = y \notin \Gamma$ which contradicts the K -isomorphism

invariance of Γ . Consequently, Γ^c is even Lorentz invariant by what we have already shown. Let us now prove the Lorentz-invariance of Γ . For this purpose choose a point $x \in \Gamma$ and assume that $y := L_\theta(x) \notin \Gamma$ for some Lorentz transformation L_θ . Therefore, the Lorentz transformation $L_{-\theta} = L_\theta^{-1}$ maps $y \in \Gamma^c$ to $x \in \Gamma$. This contradicts the Lorentz-invariance of Γ^c . Consequently, Γ is Lorentz invariant. $\square$

5.9 Lemma *Let $K \subset \mathbb{R}$ be an algebraic number field. Let $S^1 := \{(x, y) \in \mathbb{R}^2 | x^2 + y^2 = 1\} \subset \mathbb{R}^2$ be the sphere of radius one endowed with its canonical topology. Let $S^1(K) := \{(x, y) \in K^2 | x^2 + y^2 = 1\} \subset S^1$ be the set of K -valued points of S^1 . Then $S^1(K)$ is a dense subset of S^1 .*

PROOF First notice that for any $t \in K$

$$z_t := \left(\frac{1-t^2}{1+t^2}, \frac{2t}{1+t^2} \right) \in S^1(K).$$

Identifying $\mathbb{R}^2$ with the complex numbers $\mathbb{C}$ in a canonical way, we may write $z_t = e^{i\varphi_t}$ for some angle φ_t . If we let $t \rightarrow 0$, we see that there are points $z_t \in S^1(K)$ with arbitrary small, but non-vanishing angle $\varphi_t \neq 0$.

Let now $z = e^{i\varphi} \in S^1$ and $\varepsilon > 0$. By what we have seen, there is an $m \in \mathbb{N}$ and a point $z_t = e^{i\varphi_t} \in S^1(K)$ such that $|\varphi - m\varphi_t| \leq \varepsilon$ (simply choose φ_t sufficiently small). But $e^{im\varphi_t} \in S^1(K)$, because $e^{im\varphi_t} = (e^{i\varphi_t})^m$. Therefore we are done because $\varepsilon > 0$ is arbitrary. $\square$

However, in our “adelic” approach to physics, space-time is given by the set $X_K(K)$ of K -valued points of a K -scheme X_K . As motivated above, one should expect for physical reasons that $X_K(K)$ is K -isomorphism invariant. This is indeed always true:

5.10 Proposition *Let $K \subset \mathbb{R}$ be an algebraic number field, and let X_K be a K -scheme. Then $X_K(K) \subset X_K(\mathbb{R})$ is K -isomorphism invariant (see Definition 5.7, b)).*

PROOF Let $\mathfrak{C}$ be the category of K -schemes. Each object $X \in \mathfrak{C}$ gives rise to its functor of points $h_X : \mathfrak{C} \rightarrow (\text{Sets})$ which associates to any $T \in \mathfrak{C}$ the set $h_X(T) := X(T) := \text{Hom}(T, X)$ of T -valued points of X . Each morphism $X \rightarrow X'$ in $\mathfrak{C}$ induces a morphism $h_X \rightarrow h_{X'}$ of functors by the composition of morphisms in $\mathfrak{C}$. In this way one gets a covariant functor

$$f : \mathfrak{C} \rightarrow \text{Hom}(\mathfrak{C}^0, (\text{Sets}))$$

of $\mathfrak{C}$ to the category of covariant functors from $\mathfrak{C}^0$ (the dual of $\mathfrak{C}$) to the category of sets which (by Proposition 11.43) defines $\mathfrak{C}$ as full subcategory of $\text{Hom}(\mathfrak{C}^0, (\text{Sets}))$.

In particular, each K -isomorphism $\varphi : X_K \rightarrow X_K$ gives rise to a natural transformation $f(\varphi) : h_{X_K} \rightarrow h_{X_K}$. The latter means that for all K -morphisms $T \rightarrow T'$ there is a commutative diagram

$$\begin{array}{ccc} X_K(T') & \xrightarrow{\sim} & X_K(T') \\ \downarrow & & \downarrow \\ X_K(T) & \xrightarrow{\sim} & X_K(T) \end{array}$$

whereby the horizontal maps are isomorphisms. If we consider the special case of the canonical K -morphism $\text{Spec } \mathbb{R} \rightarrow \text{Spec } K$ corresponding to the inclusion $K \subset \mathbb{R}$, we obtain the commutative diagram

$$\begin{array}{ccc} X_K(K) & \xrightarrow{\sim} & X_K(K) \\ \downarrow i & & \downarrow i \\ X_K(\mathbb{R}) & \xrightarrow{\sim} & X_K(\mathbb{R}) \end{array}$$

Thereby $i : X_K(K) \hookrightarrow X_K(\mathbb{R})$ denotes the canonical inclusion. This means exactly that $X_K(K)$ is K -isomorphism invariant. $\square$

5.11 Remark a) Consider the affine space $X_K = \mathbb{A}_K^n$. In quantum field theory, one usually chooses $\mathbb{A}_K^n(\mathbb{R})^n \cong \mathbb{R}^n$ as underlying space-time manifold. As already mentioned the lattices $\Gamma \cong \mathbb{Z}^n \subset \mathbb{R}^n$ are obviously not Lorentz-invariant. In particular, we know from Proposition 5.8 that they are not K -isomorphism invariant. Conversely, the canonical K -isomorphism invariant set $\mathbb{A}_K^n(K)$ is neither a finitely generated abelian group nor a lattice, because $\mathbb{A}_K^n(K) = K^n$ is a dense subset of $\mathbb{R}^n$.

b) Let us now consider a product $X_K = E_K^{(1)} \times_K \dots \times_K E_K^{(n)}$ of elliptic curves over K . Like Minkowski space-time, also $X_K(\mathbb{R}) = E_K^{(1)}(\mathbb{R}) \times \dots \times E_K^{(n)}(\mathbb{R})$ is a solution of Einstein's equations which describes a vacuum without gravity. But this time we know from Theorem 5.1 that $X_K(K)$ is a finitely generated abelian group. If the rank of this group is zero, $X_K(K)$ is a discrete set. Furthermore, the set $X_K(K)$ is K -isomorphism invariant.

All in all, it makes sense to interpret the set $X_K(K)$ as vacuum (as it is suggested from the adelic point of view anyway). Gauge fields may be quantized in a non-perturbative way by means of lattice gauge theory.

c) If there is an Abelian variety A_K over K , which is solution of the Einstein equations, the statements of b) remain essentially true. The only difference is that gravity need not be trivial, i.e., in general $A_K(K)$ is not a vacuum.

Let us from now on consider the Abelian variety $X_K := E_K^{(1)} \times_K \dots \times_K E_K^{(n)}$ defined as a product of smooth elliptic curves over K . Let us consider the K -isomorphism invariant set $X_K(K) \subset X_K(\mathbb{R})$ as vacuum. If $X_K(K)$ is a group of rank zero, it is a discrete set. In the continuum, gauge fields are solutions of the continuum Yang-Mills equations. In the lattice formulation, the respective expressions are obtained from the continuum Yang-Mills theory by replacing the infinitesimal differential operators by finite difference operators. There is a canonical way to introduce difference operators: Each elliptic curve $E_K^{(i)}$ induces a one dimensional Riemannian manifold $M^i \cong E_K^{(i)}(\mathbb{R})$ (if we forget the point at infinity). For each manifold M^i we may choose an embedding $\gamma^i : \mathbb{R} \rightarrow \mathbb{R}^2$. The corresponding first fundamental form on M^i induces a Levi-Civita connection, and we

may perform parallel transports of tangent vectors along γ^i . If $M_{\gamma^i(t)}^i$ denotes the tangent space of M^i at $\gamma^i(t)$, the parallel transport from $M_{\gamma^i(t)}^i$ to $M_{\gamma^i(s)}^i$ is usually denoted by $\tau_{s,t}^i$. Then the covariant derivation in the continuum may be written as

$$\nabla_{(\gamma^i)'(t)} \mathbf{v} = \lim_{h \rightarrow 0} \frac{1}{h} \left(\tau_{t,t+h}^i \mathbf{v}(t+h) - \mathbf{v}(t) \right).$$

The set $E_K^{(i)}(K)$ is a finite, discrete subgroup of the continuum $E_K^{(i)}(\mathbb{R})$. Consequently, there is a finite set $\{t_1, \dots, t_m\}$ of real numbers $t_j < t_{j+1} \in \mathbb{R}$ whose image under γ^i is $E_K^{(i)}(K)$. At the K -valued point $\gamma^i(t_j) \in E_K^{(i)}(K)$, we may therefore introduce the following discrete notions.

- a) $(\nabla_i f)(t_j) := \frac{1}{t_{j+1} - t_j} \left(f(t_{j+1}) - f(t_j) \right)$ for all functions f .
- b) $(\bar{\nabla}_i f)(t_j) := \frac{1}{t_j - t_{j-1}} \left(f(t_j) - f(t_{j-1}) \right)$ for all functions f .
- c) $(\nabla_i \mathbf{v})(t_j) := \frac{1}{t_{j+1} - t_j} \left(\tau_{t_j, t_{j+1}}^i \mathbf{v}(t_{j+1}) - \mathbf{v}(t_j) \right)$ for all vector-fields $\mathbf{v}$.
- d) $(\bar{\nabla}_i \mathbf{v})(t_j) := \frac{1}{t_j - t_{j-1}} \left(\mathbf{v}(t_j) - \tau_{t_j, t_{j-1}}^i \mathbf{v}(t_{j-1}) \right)$ for all vector-fields $\mathbf{v}$.
- e) $\int f := \sum_j (t_{j+1} - t_j) f(t_j)$

On the set $X_K(K) = E_K^{(1)}(K) \times \dots \times E_K^{(n)}(K)$, ∇_i resp. $\bar{\nabla}_i$ is the discrete analogue of the partial derivative in the i -th direction.

These definitions naturally extend the classical notions of discrete calculus in Minkowski space, and are the starting point of a lattice gauge theory for Abelian varieties. However, it is beyond the scope of this thesis to work out this lattice gauge theory explicitly. Instead, in a first step, we will treat Yang-Mills theory over arbitrary base schemes. This is done in Part II of this thesis.

6. Appendix I

6.1. Néron models

We already mentioned in section 0.2 that a model of type (GR) $(X \rightarrow S, g)$ is almost a Néron model. A model of type (SR) (see Definition 0.7) actually is a Néron model. These Néron models are universal objects which we will introduce in this section.

6.1 Definition Let X_K be a smooth and separated K -scheme of finite type. A *Néron model* of X_K is an S -model X which is smooth, separated and of finite type, and which satisfies the following universal property, the so called Néron mapping property:

For each smooth S -scheme Y and each K -morphism $u_K : Y_K \rightarrow X_K$ there is an unique S -morphism $u : Y \rightarrow X$ extending u_K .

It is highest non-trivial to prove the existence of the Néron model, and in general it will not exist without further assumptions. At least, due to its universal property, it is clear that the Néron model is unique up to canonical isomorphism. Due to its uniqueness and its universal property, the formation of Néron models is a functor.

6.2 Remark If the archimedean limit X_K admits a Néron model X , and if X_K in addition carries a group structure (e.g. if $X \rightarrow S$ is a model of type (SR)), this group structure extends to X due to the Néron mapping property. Interpreting the formation of the inverse with respect to the group structure as a simultaneous parity and time-reversal operation, we see that there is a canonical notion of parity and time-reversal on X , too. Thus, parity and time-reversal lift to the “adelic” level.

We finish this section with the statement of a further fundamental property of Néron models whose significance will become clear in the next chapter, too. It is the so called *extension property for étale points*. Physically, this property guarantees that the archimedean zero-dimensional space-time points appear as projections of the “adelic” space-time points to their archimedean component.

6.3 Definition Let X be a scheme over a Dedekind scheme S . Then we say that X satisfies the extension property for étale points at a closed point $s \in S$ if, for all étale local $\mathcal{O}_{S,s}$ -algebra R' with field of fractions K' , the canonical map $X(R') \rightarrow X_K(K')$ is surjective.

6.4 Proposition Let S be a Dedekind scheme with field of fractions K , and let X_K be a smooth and separated K -scheme of finite type which admits a Néron model X . Furthermore, let S' be a second Dedekind scheme with field of fractions K' such that $S' \rightarrow S$ is étale. Then there is a bijection

$$X(S') \cong X_K(K').$$

PROOF Due to the Néron mapping property, each morphism $\mathrm{Spec} K' \rightarrow X_K$ extends to a unique morphism $S' \rightarrow X$. $\square$

Important is the fact that Néron models are local on the base which is stated by means of the following proposition.

6.5 Proposition *Let S be a Dedekind scheme and let X be an S -scheme of finite type. Then the following assertions are equivalent:*

- a) *X is a Néron model of its generic fibre.*
- b) *For each closed point $s \in S$, $X \times_S \mathrm{Spec} \mathcal{O}_{S,s}$ is a Néron model of its generic fibre.*

PROOF [BLR], Prop. 1.2/4 $\square$

6.6 Proposition *Let X be a smooth R -group scheme of finite type or a torsor under a smooth R -group scheme of finite type. Then the following conditions are equivalent:*

- a) *X is a Néron model of its generic fibre X_K .*
- b) *X is separated and the canonical map $X(R^{sh}) \rightarrow X_K(K^{sh})$ is surjective.*
- c) *The canonical map $X(R^{sh}) \rightarrow X_K(K^{sh})$ is bijective.*

PROOF [BLR], Thm. 7.1/1 $\square$

The surjectivity of the canonical map $X(R^{sh}) \rightarrow X_K(K^{sh})$ is due to the definition of the strict henselisation equivalent to the extension property for étale points in the case of a local base scheme $\mathrm{Spec} R$. Furthermore, due to the valuative criterion of separatedness, the surjection is even a bijection as stated in Proposition 6.6 c).

6.2. Further statements on the archimedean component

Let us consider a model of type (GR) $(X \rightarrow S, g)$ in the sense of Definition 0.4. Furthermore assume that $K \subset \mathbb{R}$ is an algebraic number field. The purpose of this subsection is to illustrate in more detail how the archimedean continuum limit $\mathfrak{X}_\infty$ (see section 0.3) emerges. Let us for simplicity assume that $X \rightarrow S$ is representable by a smooth S -scheme of relative dimension n . Consider the archimedean component of X , i.e. the generic fibre X_K of X . Due to smoothness, there are local $\mathrm{Spec} K$ -immersion $X_K \hookrightarrow \mathbb{A}_K^n$ (see Definition 1.10). Let us for simplicity assume that this local immersion is already a global, closed immersion. Consequently, $X_K = V(I) = \mathrm{Spec} K[T_1, \dots, T_n]/I$ is the zero set of an ideal I . Recall that for any field $L \supset K$, $X_K(L) := \mathrm{Hom}_{\mathrm{Spec} K}(\mathrm{Spec} L, X_K) \cong \mathrm{Hom}_K(K[T_1, \dots, T_n]/I, L)$. Thus, due to the universal property of quotients, we see that

$$X_K(L) = \{(x_1, \dots, x_n) \in L^n \mid p(x_1, \dots, x_n) = 0 \text{ for all } p \in I\}.$$

Choosing $L = \mathbb{R}$, we obtain a set which may be endowed with the structure of a differentiable manifold. This is the archimedean continuum limit $\mathfrak{X}_\infty := X_K(\mathbb{R})$ of X . However, a point $x_\infty \in \mathfrak{X}_\infty$ is in general not induced by a point of the underlying scheme X_K . As we will illustrate below, only those points $x_\infty \in \mathfrak{X}_\infty$ origin from prime ideals of X_K , whose coordinates take values in an algebraic closure $\overline{K}$ of K . These point are elements of the set $X_K(K^{\text{sep}}) = X_K(\overline{K})$, where K^{sep} is a separable algebraic closure of K . Thereby, we may write $K^{\text{sep}} = \overline{K}$ because $\text{char} K = 0$.

Let $x = (x_1, \dots, x_n) \in X_K(K^{\text{sep}})$. Then we can associate to it the maximal ideal $\mathfrak{m}_x := (T_1 - x_1, \dots, T_n - x_n) \subset K^{\text{sep}}[T_1, \dots, T_n]$. Using Taylor expansion around x we see that $I \cdot K^{\text{sep}}[T_1, \dots, T_n] \subset \mathfrak{m}_x$, and therefore $\mathfrak{m}_x$ corresponds to a closed point of the scheme X_K . Let us denote this point by x , too. More precisely any element of $X_K(K^{\text{sep}})$ is uniquely determined by the data consisting of a closed point $x \in X_K$ and a K -algebra homomorphism $K \hookrightarrow K^{\text{sep}}$. Alternatively, one can prove the following result.

6.7 Proposition *Let X_K be a scheme over a field K . Then there is a bijection*

$$X_K(K) \cong \{x \in X_K \mid k(x) = K\}$$

PROOF Let $\sigma \in X_K(K)$ and let x be the image of the point of $\text{Spec } K$. The homomorphism σ_x^* induces a field homomorphism $k(x) \rightarrow K$. As $k(x)$ is K -algebra, this implies that $k(x) = K$. Let conversely be $x \in X_K$ with $k(x) = K$. Composing the closed immersion $\text{Spec } k(x) \rightarrow \text{Spec } \mathcal{O}_{X_K, x}$ with the canonical morphisms $\text{Spec } \mathcal{O}_{X_K, x} \rightarrow X_K$, one obtains the desired section $\text{Spec } K \rightarrow X_K$ whose image is x . $\square$

However, due to the algebraic nature of the underlying scheme, the set $X_K(K^{\text{sep}} \cap \mathbb{R})$ is already a dense subset of the manifold $\mathfrak{X}_\infty$ with respect to the canonical topology on $\mathfrak{X}_\infty$. In particular, $\mathfrak{X}_\infty$ may be reconstructed from $X_K(K^{\text{sep}})$. On the other hand, we saw that every point of $X_K(K^{\text{sep}}) \subset M$ gives rise to a maximal ideal, i.e. *closed* point of X_K . But these points also lie dense in X_K with respect to Zariski topology:

6.8 Proposition *If X is a smooth scheme over a field k , the set of closed points x of X such that $k(x)$ is a separable extension of k is dense in X .*

PROOF [BLR], Cor. 2.2/13 $\square$

This motivates the following notion.

6.9 Definition Let $f : X \rightarrow S$ be a smooth scheme. Let $x \in X$ and $s := f(x)$. Then x is called a *physical point*, if:

- a) x is closed in the fibre $X_s = X \times_S \text{Spec } k(s)$.
- b) $k(s) \hookrightarrow k(x)$ is a separable.

6.10 Remark a) Let X_K be a variety over K (e.g. our smooth space-time continuum) and let K'/K a Galois extension (e.g. $K' = K^{\text{sep}}$) with Galois group G acting on X_K in the canonical way: Any $\sigma \in G$ induces an automorphism of

K -schemes $\mathrm{Spec} K \rightarrow \mathrm{Spec} K$. Thus there is a canonical action of G on $X'_K := X_K \times_{\mathrm{Spec} K} \mathrm{Spec} K'$ which is the identity on the first component. If X_K is as in a), then for any $\sigma \in G$ and for any $x = (x_1, \dots, x_n)$ we have $\sigma(x) = \sigma(x_1), \dots, \sigma(x_n)$. The set of classes $G \backslash X_K(K')$ injects into X_K .

More precisely the physical points of X_K are Galois orbits of physical points of $X_K(K^{\mathrm{sep}})$ (i.e. of the points of the archimedean continuum limit of space-time). Thus X_K occurs in a canonical way as a base space for some kind of “gauge theory” (of space-time) whose “gauge group” is a Galois group.

b) Intuitively, we may think of the bijections

$$X_K(K) \cong \{x \in X_K \mid k(x) = K\} \quad \text{or} \quad X_K(K^{\mathrm{sep}}) \cong \{\text{physical points of } X_{K^{\mathrm{sep}}}\}$$

as some kind of “wave-particle duality” on space-time level. For example, the elements of $X_K(K^{\mathrm{sep}})$ are sections of the structure morphism $X_{K^{\mathrm{sep}}} \rightarrow K^{\mathrm{sep}}$. But morphisms of schemes have similar symmetries like the differential geometric morphisms of vector bundles (e.g. one needs a one-cocycle condition in order to glue). Therefore, $X_K(K^{\mathrm{sep}})$ represents the wave interpretation of space-time points, because sections of vector bundles are classically interpreted as fields.

But the physical points of $X_{K^{\mathrm{sep}}}$ are points of a topological space. They represent the particle interpretation of space-time points.

Part II.

Arithmetic Yang-Mills theory

7. Some essential techniques

In ordinary (i.e. $\mathbb{R}$ -valued) Yang-Mills theory, gauge fields are described by co-vector-fields. If we want to generalize the $\mathbb{R}$ -valued, differential geometric Yang-Mills theory to arbitrary commutative rings R (or even base schemes S), we therefore have to make use of the notion of the algebraic geometric tangent bundle. In particular, we need a suitable notion of pull-back and push-forward of tensor fields in the realm of algebraic geometry. This chapter is devoted to the development of such a notion (see section 7.2). We start with some introductory remarks on general coordinate transformations in the setting of algebraic geometry.

7.1. General coordinate transformations

Consider a classical physical situation: $S = \text{Spec } \mathbb{R}$ and $\psi : X' \rightarrow X$ is a morphism of smooth algebraic varieties X, X' over $\mathbb{R}$ (Definition 1.10). Then the sets of $\mathbb{R}$ -valued points $X(\mathbb{R}), X'(\mathbb{R})$ may be considered as differential manifolds, and ψ gives rise to a morphism $\psi(\mathbb{R}) : X(\mathbb{R}) \rightarrow X'(\mathbb{R})$ of differential manifolds (which was illustrated in the physical interpretation after Definition 1.10). We saw within the bounds of the physical interpretation following Definition 1.3 that on $\mathbb{R}$ -valued points $x \in X(\mathbb{R})$ the pull-back map $\psi^* \Omega_{X/S}^1 \rightarrow \Omega_{X'/S}^1$ of differential forms coincides with the ordinary, differential geometric notion of pull-back of 1-forms: It is described by the Jacobian matrix $D\psi$. If we choose especially $X = X'$, then $X(\mathbb{R}) = X'(\mathbb{R})$ and $\psi(\mathbb{R}) : X(\mathbb{R}) \rightarrow X(\mathbb{R})$ is called a diffeomorphism (of $X(\mathbb{R})$ in the differential geometric sense), if $D\psi$ is an invertible matrix. However, physics should be general covariant, i.e. if we pass from one observer to another one by means of a coordinate transformation, the physical laws should be invariant. General relativity is even only determined up to diffeomorphisms.

It is our goal to evoke a Yang-Mills theory over arbitrary numbers (given by a commutative ring R) which is therefore no longer limited to the real numbers $\mathbb{R}$. In particular, we have to find the algebraic geometric analogue of diffeomorphisms. Starting from the differential geometric characterization of diffeomorphisms as morphisms with invertible Jacobi matrix and recalling our analysis at the beginning of this section, we will see in Corollary 7.2 that the algebraic geometric analogues of diffeomorphisms are exactly the étale morphisms.

7.1 Definition A morphism of schemes $f : X \rightarrow S$ is called *étale* (at a point) if it is smooth (at the point) of relative dimension 0 (see Definition 1.10 for the notion of smoothness).

7.2 Corollary *Let $f : X \rightarrow Y$ be an S -morphism. Let x be a point and let $y = f(x)$. Assume that X is smooth over S at x and that Y is smooth over S at y . Then the following conditions are equivalent:*

- a) *f is étale at x .*
- b) *The canonical homomorphism $(f^*\Omega_{Y/S}^1)_x \rightarrow (\Omega_{X/S}^1)_x$ is bijective.*

PROOF [BLR], Cor. 2.2/10 □

7.3 Corollary *Let $f : X \rightarrow Y$ be an S -morphism of smooth S -schemes X and Y . Then is equivalent:*

- a) *f is étale.*
- b) *The canonical homomorphism $f^*\Omega_{Y/S}^1 \rightarrow \Omega_{X/S}^1$ is bijective.*
- c) *The canonical homomorphism $(f^*\Omega_{Y/S}^1)_x \rightarrow (\Omega_{X/S}^1)_x$ is bijective for all $x \in X$.*

This feature of étale morphism has a far reaching consequence and leads to a phenomenon which does not occur in differential geometry: We already mentioned that we should only look at “space-time ” and “physical objects” up to coordinate transformation. Thus from a physical point of view one should not distinguish between X' and X if $X' \rightarrow X$ is surjective and étale. But in contrast to the differential geometric setting, an étale, surjective morphism $X' \rightarrow X$ will in general not be an isomorphism if we work over arbitrary rings R (instead of e.g. the complex numbers $\mathbb{C}$). Thus, X and X' are different with respect to Zariski topology. But from the physical point of view, we should not distinguish between X and X' . Therefore, Zariski topology is physically not appropriate. At least, one knows that étale morphisms are open due to flatness. But, we have to extend the Zariski topology in such a way that X' is “open” in X with respect to the bigger “topology” if $X' \rightarrow X$ is étale. If $X' \rightarrow X$ is étale, surjective we want to consider X' as an “open” covering of X consisting of only one element. In such a case we will consider them as equal in a certain way: “ $X' = X$ ”. We may say this mathematically more correctly as follows:

We have to work with respect to étale topology instead of Zariski topology.

Etale topology is an example of a so called Grothendieck topology whose definition we are going to recall next.

7.4 Definition A *Grothendieck topology* $\mathfrak{T}$ consists of a category $\text{Cat } \mathfrak{T}$ and a set $\text{Cov } \mathfrak{T}$ of families $(U_i \rightarrow U)_{i \in I}$ of morphisms in $\text{Cat } \mathfrak{T}$, called *covering*, such that the following holds:

- a) If $\Phi : U \rightarrow V$ is an isomorphism in $\text{Cat } \mathfrak{T}$, then $(\Phi) \in \text{Cov } \mathfrak{T}$.
- b) If $(U_i \rightarrow U)_{i \in I}$ and $(V_{ij} \rightarrow U_i)_{j \in J_i}$ for $i \in I$ belong to $\text{Cov } \mathfrak{T}$, then the same is true for the composition $(V_{ij} \rightarrow U_i \rightarrow U)_{i \in I, j \in J_i}$.

- c) If $(U_i \rightarrow U)_{i \in I}$ is in $\text{Cov } \mathfrak{T}$ and if $V \rightarrow U$ is a morphism in $\text{Cat } \mathfrak{T}$, then the fibred products $U_i \times_U V$ exist in $\text{Cat } \mathfrak{T}$, and $(U_i \times_U V \rightarrow V)_{i \in I}$ belongs to $\text{Cov } \mathfrak{T}$.

We may think of the objects of $\text{Cat } \mathfrak{T}$ as of the open sets of our topology and of the morphisms of $\text{Cat } \mathfrak{T}$ as of the inclusions of open sets. Furthermore, a family $(U_i \rightarrow U)_{i \in I}$ of $\text{Cov } \mathfrak{T}$ has to be interpreted as a covering of U by the U_i and a fibred product of $U_i \times_U V$ as the intersection of U_i and V . Thinking along this lines an ordinary topological space X is canonically equipped with a Grothendieck topology: $\text{Cat } \mathfrak{T}$ is the category of open subsets of X with inclusions as morphisms, and $\text{Cov } \mathfrak{T}$ consists of all open covers of open subsets of X .

7.5 Definition For a scheme X , let $\text{Cat } \mathfrak{T}$ be the category of étale X -schemes with étale morphisms as morphisms, and let $\text{Cov } \mathfrak{T}$ be the set of families $(U_i \rightarrow U)_{i \in I}$ of étale morphisms such that $U = \bigcup_{i \in I} U_i$. Then $\mathfrak{T}$ is called the *étale topology* on X .

It should be pointed out that the “intersection” of “open” sets is dealt with in condition c) of Definition 7.4, whereas we have refrained from giving any sense to the union of “open” sets. In fact, even in examples, where the union of “open” sets does make sense, such a union will not necessarily yield an “open” set again.

The notion of a Grothendieck topology has been designed in such a way that the notion of presheaf or sheaf can easily be adapted to such a situation:

7.6 Definition Let $\mathfrak{T}$ be a Grothendieck topology and $\mathfrak{C}$ be a category which admits cartesian products. A *presheaf* on $\mathfrak{T}$ with values in $\mathfrak{C}$ is a contravariant functor $\mathcal{F} : \text{Cat } \mathfrak{T} \rightarrow \mathfrak{C}$. We call $\mathcal{F}$ a *sheaf* if the diagram

$$\mathcal{F}(U) \rightarrow \prod_{i \in I} \mathcal{F}(U_i) \rightrightarrows \prod_{i, j \in I} \mathcal{F}(U_i \times_U U_j)$$

is exact for any covering $(U_i \rightarrow U)_{i \in I}$ in $\text{Cov } \mathfrak{T}$.

Having defined the “open subsets” of a scheme X with respect to étale topology, we may now introduce the notion of neighborhoods of points.

7.7 Definition A morphism $X' \rightarrow X$ of schemes is called an *étale neighborhood* of a point $x \in X$, if $X' \rightarrow X$ is étale and if x is contained in the image of X' .

The following Proposition 7.8 finally gives the justification, why it is physically admissible to consider étale morphisms $X \rightarrow Y$ as inclusions.

7.8 Proposition *Let $f : X \rightarrow Y$ be locally of finite type. Let $x \in X$ and $y := f(x)$. If f is étale at x (resp. unramified at x), there exists an étale neighborhood $Y' \rightarrow Y$ of y such that, locally at each point of $X := X \times_Y Y'$ above x , the morphism $f' : X' \rightarrow Y'$ obtained by base change $Y' \rightarrow Y$ is an open immersion (resp. an immersion).*

PROOF [BLR], Prop. 2.3/8

□

Therefore, the algebraic geometric analogue of the differential geometric diffeomorphism should be the étale morphism. However, if we perform a coordinate transformation or any “deformation” of space-time, the number of points should be invariant under coordinate transformations. In this situation, we should therefore restrict attention to the class of finite, étale surjective morphisms $f : X \rightarrow Y$ which have the property that there are bijections $Y_K(K') = X_K(K')$ and $Y(S') = X(S')$ for all étale S -schemes S' with field of fractions K' . Thereby, X_K and Y_K denote as usually the respective generic fibres. However, let us point to the fact that it can happen that f changes the topology.

Working with respect to étale topology, there is the *theorem of the inverse function* for smooth schemes. This is crucial if we want to perform analysis, and we have to make use of it later when we develop Yang-Mills theory over rings. It takes the following form.

7.9 Proposition *Let $f : X \rightarrow S$ be a smooth morphism. Let s be a point of S , and let x be a closed point of the fibre $X_s = X \times_S \text{Spec } k(s)$ such that $k(x)$ is a separable extension of $k(s)$. Then there exists an étale morphism $g : S' \rightarrow S$ and a point $s' \in S'$ above s such that the morphism $f' : X \times_S S' \rightarrow S'$ obtained from f by the base change $S' \rightarrow S$ admits a section $h : S' \rightarrow X \times_S S'$, where $h(s')$ lies above x , and where $k(s') = k(x)$.*

PROOF [BLR], Prop. 2.2/14 □

7.10 Proposition *Let X be an S -scheme. Then the notion of physical points is stable under étale base change.*

PROOF Consider a physical point $x \in X$ over $s \in S$, and let $S' \rightarrow S$ be an étale morphism. There is a closed immersion $\{x\} \hookrightarrow X_s = X \otimes_S k(s)$. Therefore the morphism $S' \times_S \text{Spec } k(x) \hookrightarrow (X \times_S S') \times_S k(s)$ obtained by base change with $S' \rightarrow S$ is a closed immersion, too. Furthermore, $S' \times_S \text{Spec } k(x) \rightarrow \text{Spec } k(x)$ is étale. In particular, $S' \times_S \text{Spec } k(x)$ is a finite set. More precisely it is a finite disjoint union of separable field extensions of $k(x)$, and thus $S' \times_S \text{Spec } k(x)$ is equipped with discrete topology. It follows that each point of $S' \times_S \text{Spec } k(x)$ is a closed point. Let $h(s') \in S' \times_S \text{Spec } k(x)$ be a point over $s' \in S'$, where s' is over s . Due to the closed immersion $S' \times_S \text{Spec } k(x) \hookrightarrow (X \times_S S') \times_S k(s)$, $h(s')$ is a point of $(X \times_S S') \times_{S'} k(s')$ which is closed in $(X \times_S S') \times_S k(s)$. But then, the continuity of the canonical inclusion

$$\begin{array}{ccc} (X \times_S S') \times_S k(s) & \xlongequal{\quad} & (X \times_S S') \times_{S'} (S' \times_S k(s)) \\ & & \uparrow \\ (X \times_S S') \times_{S'} k(s') & \xlongequal{\quad} & (X \times_S S') \times_{S'} (k(s') \times_S k(s)) \end{array}$$

shows that $h(s')$ is already closed in $(X \times_S S') \times_{S'} k(s')$.

It remains to show that $k(h(s'))$ is separable over $k(s')$. By assumption $S' \rightarrow S$ is étale and it follows that $k(s) \hookrightarrow k(s')$ is separable. Furthermore $X \times_S S' \xrightarrow{\text{pr}_1} X$ is étale, too. But $h(s')$ is mapped to x under this morphism which implies that $k(x) \hookrightarrow k(h(s'))$ is

separable. Finally $k(s) \hookrightarrow k(x)$ is separable, because $x \in X_s$ is a physical point. Due to the canonical commutative diagram

$$\begin{array}{ccc} k(h(s')) & \longleftarrow & k(x) \\ \uparrow & & \uparrow \\ k(s') & \longleftarrow & k(s) \end{array}$$

$k(h(s'))$ has to be separable over $k(s')$, because separable field extensions are transitive. All in all we see that all points, which lie over the physical point x after étale base change, are physical points, too. $\square$

7.2. Pull-back and push-forward of tensor fields

During this section, we assume $f : X \rightarrow Y$ to be a smooth morphism of smooth S -schemes. We already introduced the pull-back of differential forms in Definition 1.3. Within this section we will prove at first that smooth schemes allow a local (with respect to étale topology) push-forward of vector fields. Finally we will use these notions as a basis for an inductive definition of pull-back and push-forward of tensor fields. We begin with the statement of some preparing results.

7.11 Lemma *If X is a smooth scheme over a field k , the set of closed points $x \in X$ such that $k(x)$ is a separable extension of k is dense in X . Recall that we call these points physical points.*

PROOF [BLR], Cor. 2.2/13 $\square$

For all S -schemes $Z \rightarrow S$, let us denote the fibre over a point $s \in S$ by Z_s . For each morphism $f : X \rightarrow Y$ of S -schemes and for all $s \in S$, denote by $f_s : X_s \rightarrow Y_s$ the induced morphism on the fibre over $s \in S$. With this notation we may state the following result.

7.12 Proposition *Let $f : X \rightarrow Y$ be a morphism of S -schemes such that X_s and Y_s are algebraic varieties over $k(s)$ for all $s \in S$. Let $x \in X$ be a physical point. Then $y := f(x)$ is a physical point, too.*

PROOF By assumption, the morphism $X \rightarrow Y$ induces a family of morphisms of algebraic varieties $X_s \rightarrow Y_s$ over $k(s)$, $s \in S$. Let x be a physical point, $y = f(x)$. Looking at the local rings, we have got commutative diagrams

$$\begin{array}{ccc} k(x) & \longleftarrow & k(y) \\ \uparrow & \nearrow & \\ k(s) & & \end{array}$$

As $k(s) \hookrightarrow k(x)$ is a separable algebraic extension, the same is true for $k(s) \hookrightarrow k(y)$. In particular $y \in Y_s$ is a closed point in Y_s , because for algebraic varieties Z over a field k one knows the following fact (e.g. due to [Liu], Ex. 2.5.9): $z \in Z$ is closed if and only if $k \hookrightarrow k(z)$ is algebraic. $\square$

7.13 Proposition *Let $f : X \rightarrow Y$ be a smooth morphism of S -schemes. Let $y \in f(X)$ be a physical point. Then there exists a physical point $x \in X$ which is mapped to y .*

PROOF Let s be the image of y under the structure morphism $Y \rightarrow S$. We first remark that we write x_y (resp. x_s) if we consider $x \in X$ as a point of X_y (resp. of X_s). As $X_y \rightarrow k(y)$ is smooth and $X_y \neq \emptyset$, we know by Lemma 7.11 that there exists a $x_y \in X_y$ such that x_y is closed in X_y , and such that $k(y) \hookrightarrow k(x_y)$ is separable. By [Liu], Ex. 2.5.9 this is equivalent to the statement that $k(y) \hookrightarrow k(x_y)$ is a separable algebraic extension. If s denotes the image of y under the structure morphism $Y \rightarrow S$ we know furthermore by assumption that: $k(s) \hookrightarrow k(y_s)$ is separable algebraic. So we are done by the following lemma if we apply [Liu], Ex. 2.5.9 once again. $\square$

7.14 Lemma *Let $f : X \rightarrow Y$ be a morphism of schemes. Let $x \in X$ and $y := f(x)$. If we denote by x_y the point x considered as a point of the fibre X_y , then:*

$$k(x_y) = k(x).$$

PROOF Considered as a scheme, the fibre X_y is isomorphic to $X \times_Y \text{Spec } k(y)$. Therefore we get for the local ring at $x_y \in X_y$

$$\mathcal{O}_{X_y, x_y} = \mathcal{O}_{X, x} \otimes_{\mathcal{O}_{Y, y}} k(y) = \mathcal{O}_{X, x} / \mathfrak{m}_y \mathcal{O}_{X, x}.$$

In particular the maximal ideal $\mathfrak{m}_{x_y}$ of $\mathcal{O}_{X_y, x_y}$ is given by $\mathfrak{m}_{x_y} = \mathfrak{m}_x / \mathfrak{m}_y \mathcal{O}_{X, x}$. Therefore

$$k(x_y) = \mathcal{O}_{X_y, x_y} / \mathfrak{m}_{x_y} = (\mathcal{O}_{X, x} / \mathfrak{m}_y \mathcal{O}_{X, x}) / (\mathfrak{m}_x / \mathfrak{m}_y \mathcal{O}_{X, x}) = \mathcal{O}_{X, x} / \mathfrak{m}_x = k(x). \quad \square$$

Consider now the smooth morphism $f : X \rightarrow Y$ of S -schemes. Let us consider a (local) vector field $\mathfrak{v} : U \rightarrow T_{X/S}$ on X . We would like to define a local push-forward of $\mathfrak{v}$ under f to a local vector field $f_* \mathfrak{v}$ on Y .

This should be done in a canonical way using the pull-back of co-vector fields. Due to the duality between vector fields and co-vector fields (see Remark 11.25) we may identify $\mathfrak{v} : U \rightarrow T_{X/S}$ with its corresponding $\mathcal{O}_U$ -module-homomorphism $\mathfrak{v} : \Omega_{X/S}^1 \rightarrow \mathcal{O}_U$ which we will also denote by v . Also using the contravariant functor $\mathbb{V}$ introduced in Proposition 11.18, we will finally arrive at the following diagram as explained below:

$$\begin{array}{ccccccc} f(U) & \xleftarrow{f} & U & \xrightarrow{\mathfrak{v}} & T_{X/S} & \xrightarrow{T_{X/S}(f)} & T_{Y/S} \times_Y X \\ f^{-1}\mathcal{O}_{f(U)} & \xrightarrow{f^*} & \mathcal{O}_U & \xleftarrow{\mathfrak{v}} & \Omega_{X/S}^1 & \xleftarrow{f^*} & f^*\Omega_{Y/S}^1 \end{array} \quad (7.1)$$

Here f^* denotes as well the canonical morphism $f^{-1}\mathcal{O}_{f(U)} \rightarrow \mathcal{O}_U$ as the pull-back of co-vector fields $\omega \in \Omega_{Y/S}^1$

$$\begin{array}{ccc} f^*\Omega_{Y/S}^1 & \xrightarrow{f^*} & \Omega_{X/S}^1 \\ \omega \otimes 1 & \longmapsto & f^*\omega \end{array},$$

which we sometimes simply abbreviate by $\omega \mapsto f^*\omega$. The pull-back is defined globally, i.e. if we consider a co-vector field ω living on all of Y , we can pull it back to a co-vector field $f^*\omega$ living on *all* of X .

In contrast to this, (even globally defined) vector fields $\mathbf{v}$ may in general only be pushed forward to vector fields living on (étale-)open subsets of $f(X)$. This is analogously to the situation upon real-valued manifolds. As an example consider the projection $p : S^2 \subset \mathbb{R}^3 \rightarrow \mathbb{R}^2$, $(x, y, z) \mapsto (x, y)$ and a vector field on the two-dimensional sphere S^2 . Let us denote the corresponding flow by ϕ_t . But in general, $p \circ \phi_t$ will not be injective. Then, the push-forward does not exist globally. More precisely, we have to face the following problem, whenever we are trying to define of a push-forward $f_*\mathbf{v}$ of a vector field $\mathbf{v}$: In general there is only the following canonical diagram

$$f(U) \xleftarrow{f} U \xrightarrow{\mathbf{v}} T_{X/S} \xrightarrow{pr \circ T_{X/S}(f)} T_{Y/S},$$

but in general there is even no chance to get a local section of f . Thus we will not find a Y -morphism $V \rightarrow T_{Y/S}$ with $V \subset f(U)$ Zariski-open which is induced by $\mathbf{v}$. Thus there is no chance to define the push-forward f_*v of the given vector field $\mathbf{v}$ within Zariski-topology.

The way out of trouble is performing a local coordinate transformation, i.e. performing an étale base change $Y' \rightarrow Y$. Intuitively one may think that a badly chosen coordinate system may lead to singularities which do not exist in “reality”, but which are due to the bad choice of the coordinates. Such “imaginary” singularities may be resolved by performing a suitable coordinate transformation. As an example consider a flat space-time $\mathbb{R}^2$ with metric $ds^2 = dx^2 + dy^2$ and introduce the new coordinate system $(x', y') = (x^3/27, y)$. Then $dx' = d(x^3/27) = 1/9 \cdot x^2 dx = x'^{2/3} dx$ and therefore

$$ds^2 = x'^{-4/3} dx'^2 + dy^2$$

which is singular at $x' = 0$. But in truth there is “no” physically relevant difference between the points $(0, y)$ and $(1, y)$ of empty space-time $\mathbb{R}^2$.

So let us perform an étale base change $Y' \rightarrow Y$ as follows: Let $y \in f(U)$. Thus $f^{-1}(y) = X_y \neq \emptyset$, and by Lemma 7.11 we may choose a closed point $x \in X_y$ such that $k(y) \hookrightarrow k(x)$ is separable, i.e. we may choose a physical point $x \in X_y$. Then we are in the situation of Proposition 7.9, and we infer that there exists an étale morphism $g : Y' \rightarrow Y$ such that $f' : X \times_Y Y' \rightarrow Y'$ has a section h . Thereby f' is the morphism induced by f by base change with $g : Y' \rightarrow Y$.

We want to use this section in order to define a push-forward of $\mathbf{v}$. For this purpose let us first prove that the canonical diagram (9.1)

$$\begin{array}{ccccccc} Y & \xleftarrow{f} & X & \xrightarrow{\mathbf{v}} & T_{X/S} & \xrightarrow{T_{X/S}(f)} & T_{Y/S} \times_Y X \\ f^{-1}\mathcal{O}_Y & \xrightarrow{f^*} & \mathcal{O}_X & \xleftarrow{\mathbf{v}} & \Omega_{X/S}^1 & \xleftarrow{f^*} & f^*\Omega_{Y/S}^1 \end{array}$$

is stable under local coordinate transformations, i.e. compatible with étale base change.

So let us apply the above base change $g : Y' \rightarrow Y$ to the diagram (9.1), and let $f' : X' \rightarrow Y'$ be as above ($X' := X \times_Y Y'$). We claim that we arrive at the diagram

$$\begin{array}{ccccccc} Y' & \xleftarrow{f'} & X' & \xrightarrow{\mathbf{v}'} & T_{X'/S} & \xrightarrow{T_{X'/S}(f')} & T_{Y'/S} \times_{Y'} X' \\ f'^{-1}\mathcal{O}_Y & \xrightarrow{f'^*} & \mathcal{O}_{X'} & \xleftarrow{\mathbf{v}'} & \Omega_{X'/S}^1 & \xleftarrow{f'^*} & f'^*\Omega_{Y'/S}^1 \end{array} \quad (7.2)$$

PROOF Consider the cartesian diagram

$$\begin{array}{ccc} X' & \xrightarrow{f'} & Y' \\ j \downarrow & & \downarrow g \\ X & \xrightarrow{f} & Y \end{array}$$

As g is étale, j is étale, too (because $j = \text{id}_X \times g$). Thus we know from Corollary 7.3 that

$$\begin{aligned} g^*\Omega_{Y'/S}^1 &= \Omega_{Y/S}^1, \\ j^*\Omega_{X'/S}^1 &= \Omega_{X/S}^1. \end{aligned}$$

These relations and the equation $f \circ j = g \circ f'$, which expresses the commutativity of the above cartesian diagram, yield

$$\begin{aligned} (f \circ j)^*\Omega_{Y/S}^1 &= (g \circ f')^*\Omega_{Y/S}^1 \\ &= f'^*(g^*\Omega_{Y/S}^1) \\ &= f'^*\Omega_{Y'/S}^1. \end{aligned}$$

Also using Proposition 11.18, we deduce:

- $T_{X/S} \times_Y Y' = T_{X/S} \times_X X' = \mathbb{V}(j^*\Omega_{X/S}^1) = \mathbb{V}(\Omega_{X'/S}^1) = T_{X'/S}$.
- $(T_{Y/S} \times_Y X) \times_{Y'} Y' = T_{Y/S} \times_Y X' = \mathbb{V}((f \circ j)^*\Omega_{Y/S}^1) = \mathbb{V}(f'^*\Omega_{Y'/S}^1) = T_{Y'/S} \times_{Y'} X'$.

So we see that we arrive at the diagram (9.2) if we apply the base change $g : Y' \rightarrow Y$ to the diagram (9.1). $\square$

Let us emphasize that the base-scheme S is *not* effected by this process, i.e. f' is actually a restriction of f (with respect to étale topology). The “adelic” space-time points (which are isomorphic to S) remain unaltered; we are only looking closer at a region of the universe. Thus, with some abuse of notation, we may suppress the “dashes” in (9.2), and we may assume (after eventually shrinking Y and performing a coordinate transformation) that f has a section h . Then we derive the following canonical morphism $\widetilde{f_*\mathbf{v}} : Y \rightarrow T_{Y/S} \times_Y X$ induced by f and $\mathbf{v}$:

$$\widetilde{f_*\mathbf{v}} : Y \xrightleftharpoons[f]{h} X \xrightarrow{\mathbf{v}} T_{X/S} \xrightarrow{T_{X/S}(f)} T_{Y/S} \times_Y X$$

This induces a canonical morphism on the sheaves

$$\begin{aligned} (\widetilde{f_*\mathfrak{v}}) : f^*\Omega_{Y/S}^1 &\xrightarrow{f^*} \Omega_{X/S}^1 \xrightarrow{\mathfrak{v}} \mathcal{O}_X \xrightarrow{h^*} h_*\mathcal{O}_Y, \\ \omega \otimes 1 &\mapsto f^*\omega \mapsto \mathfrak{v}(f^*\omega) \mapsto (\widetilde{f_*\mathfrak{v}})(\omega \otimes 1) \end{aligned}$$

where $\omega \in \Omega_{Y/S}^1$. So far we found a canonical morphism $\widetilde{f_*\mathfrak{v}} : f^*\Omega_{Y/S}^1 \rightarrow h_*\mathcal{O}_Y$ such that

$$(\widetilde{f_*\mathfrak{v}})(\omega \otimes 1) = (h^* \circ \mathfrak{v} \circ f^*)(\omega) = h^*(\mathfrak{v}(f^*\omega)) \stackrel{\text{by duality}}{=} h^*((f^*\omega)(\mathfrak{v})).$$

We would like to interpret $\widetilde{f_*\mathfrak{v}}$ as our push-forward of $\mathfrak{v}$, but actually $\widetilde{f_*\mathfrak{v}}$ is not even a vector field. Luckily, Proposition 7.15 shows that the $\mathcal{O}_X$ -module homomorphism $\widetilde{f_*\mathfrak{v}} : f^*\Omega_{Y/S}^1 \rightarrow h_*\mathcal{O}_Y$ corresponds to a uniquely determined $\mathcal{O}_Y$ -module homomorphism $f_*\mathfrak{v} : \Omega_{Y/S}^1 \rightarrow \mathcal{O}_Y$, i.e. a vector field on Y . This will be a push-forward of $\mathfrak{v}$ under f .

7.15 Proposition *Let $f : X \rightarrow Y$ be a morphism of schemes and let $\mathcal{G}$ be a $\mathcal{O}_Y$ -module on Y . Then the pull-back $\mathcal{O}_X$ -module $f^*\mathcal{G}$ on X fulfills the universal property that there is a functorial isomorphism*

$$\mathrm{Hom}_{\mathcal{O}_Y}(\mathcal{G}, f_*\mathcal{F}) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{O}_X}(f^*\mathcal{G}, \mathcal{F})$$

functorial in $\mathcal{O}_X$ -modules $\mathcal{F}$ on X .

PROOF [Liu], Ex. 5.1.1 □

Let us describe this functorial isomorphism shortly. For this purpose, we assume that $\mathcal{G}$ is quasi-coherent. Let $\varphi \in \mathrm{Hom}_{\mathcal{O}_Y}(\mathcal{G}, f_*\mathcal{F})$, i.e. φ consists of a collection of morphisms

$$\varphi(V) : \mathcal{G}(V) \longrightarrow f_*\mathcal{F}(V) := \mathcal{F}(f^{-1}(V)), \quad V \text{ in } Y \text{ open,}$$

which is compatible with restrictions. For open subsets $U \subset X$ with $U \subset f^{-1}(V)$, this yields morphisms

$$\varphi(V) : \mathcal{G}(V) \longrightarrow \mathcal{F}(f^{-1}(V)) \xrightarrow{\mathrm{res}} \mathcal{F}(U), \quad V \text{ in } Y \text{ open,}$$

where “res” denotes the canonical restriction. Let now U be an affine open subset of X such that $f(U)$ is contained in an affine open subset V of Y . Then one knows (e.g. by [Liu], Prop. 5.1.14) that

$$f^*\mathcal{G}|_U \cong (\mathcal{G}(V) \otimes_{\mathcal{O}_Y(V)} \mathcal{O}_X(U))^\sim.$$

We consider $\mathcal{F}(U)$ as an $\mathcal{O}_Y(V)$ -module by means of the canonical morphism $\mathcal{O}_Y(V) \longrightarrow \mathcal{O}_X(f^{-1}(V)) \xrightarrow{\mathrm{res}} \mathcal{O}_X(U)$. The $\mathcal{O}_Y(V)$ -linear map $\varphi(V) : \mathcal{G}(V) \rightarrow \mathcal{F}(U)$ gives rise to a $\mathcal{O}_Y(V)$ -bilinear map

$$\mathcal{G}(V) \times \mathcal{O}_X(U) \rightarrow \mathcal{F}(U), \quad (g, a) \mapsto \varphi(V)(g) \cdot a.$$

Using the universal property of tensor products, we finally arrive at a $\mathcal{O}_X(U)$ -linear map

$$\tilde{\varphi}(U) : \mathcal{G}(V) \otimes_{\mathcal{O}_Y(V)} \mathcal{O}_X(U) \rightarrow \mathcal{F}(U), \quad g \otimes a \mapsto \varphi(V)(g) \cdot a.$$

As $(f^*\mathcal{G})(U) = \mathcal{G}(V) \otimes_{\mathcal{O}_Y(V)} \mathcal{O}_X(U)$, the $\mathcal{O}_X(U)$ -linear maps $\tilde{\varphi}(U)$ give rise to the searched $\mathcal{O}_X$ -module homomorphism $\tilde{\varphi} \in \text{Hom}_{\mathcal{O}_X}(f^*\mathcal{G}, \mathcal{F})$. Thus the functorial isomorphism of Proposition 7.15 may be described as follows:

$$\text{Hom}_{\mathcal{O}_Y}(\mathcal{G}, f_*\mathcal{F}) \xrightarrow{\sim} \text{Hom}_{\mathcal{O}_X}(f^*\mathcal{G}, \mathcal{F}), \quad \varphi \mapsto \tilde{\varphi}, \text{ where } \tilde{\varphi}(g \otimes a) := \varphi(g) \cdot a.$$

Now we may apply this result to our morphism $\widetilde{f_*\mathfrak{v}} : f^*\Omega_{Y/S}^1 \rightarrow h_*\mathcal{O}_Y$. Let us denote its image under the functorial isomorphism of Proposition 7.15 by $f_*\mathfrak{v} : \Omega_{Y/S}^1 \rightarrow f_*h_*\mathcal{O}_Y$. One knows that the direct image of sheaves is a covariant functor (see e.g. [Liu], Ex. 2.2.6). In particular $f_*h_*\mathcal{O}_Y = (f \circ h)_*\mathcal{O}_Y = \mathcal{O}_Y$ because h is a section of f , i.e. $f \circ h = \text{id}$. All in all we arrive at a $\mathcal{O}_Y$ -module homomorphism $f_*\mathfrak{v} : \Omega_{Y/S}^1 \rightarrow \mathcal{O}_Y$, i.e. at a vector field on Y , which by construction fulfills the equation $f_*\mathfrak{v}(\omega) = \widetilde{(f_*\mathfrak{v})}(\omega \otimes 1) = (h^* \circ \mathfrak{v} \circ f^*)(\omega)$, i.e.:

$$f_*\mathfrak{v} = h^* \circ \mathfrak{v} \circ f^*$$

7.16 Remark a) The smoothness of the morphism $f : X \rightarrow Y$ was only needed in order to guarantee the existence of local sections of f (where local is meant with respect to étale topology). If we assume that there is already a section h of f , we may drop the smoothness assumption of f . Then the construction of the push-forward works as described above, and every global vector field $\mathfrak{v} \in \Gamma(T_{X/S}/X)$ on X yields a *global* vector field $f_*\mathfrak{v} = h^* \circ \mathfrak{v} \circ f^*$ on Y .

b) If $f : X \rightarrow Y$ is an isomorphism, then $h := f^{-1}$ is a global section of f , and the push-forward of every vector field $\mathfrak{v} \in \Gamma(T_{X/S}/X)$ on X fulfills the equation:

$$f_*\mathfrak{v} = (f^{-1})^* \circ \mathfrak{v} \circ f^*.$$

This is exactly the definition of the push-forward of vector fields which is considered in differential geometry. Thus the differential geometric notion is contained as a special case.

c) Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be morphisms with sections i_f and i_g , and let $\mathfrak{v} \in \Gamma(T_{X/S}/X)$. Then

$$(g \circ f)_*\mathfrak{v} = g_*(f_*\mathfrak{v}).$$

This may be seen as follows:

$$\begin{aligned} ((g \circ f)_*\mathfrak{v})(\omega) &= (i_f \circ i_g)^* \circ \mathfrak{v} \circ (g \circ f)^*(\omega) = i_g^* \circ i_f^* \circ \mathfrak{v} \circ f^* \circ g^*(\omega). \\ (g_*(f_*\mathfrak{v}))(\omega) &= i_g^*((f_*\mathfrak{v})(g^*\omega)) = i_g^*(i_f^*(\mathfrak{v}(f^*g^*\omega))). \end{aligned}$$

7.17 Definition Let X and Y be smooth S -schemes, and let $f : X \rightarrow Y$ be a S -morphism which locally (with respect to étale topology on Y) admits a section $h : Y \hookrightarrow X$ (e.g. f is smooth). Let $\omega \in \Omega_{Y/S}^1$ be a co-vector field on Y and $v : X \rightarrow T_{X/S}$ be an vector field on X . Then, locally (in étale topology), there exists a vector field $f_*\mathbf{v}$ on Y fulfilling the following identity

$$(f_*\mathbf{v})(\omega) = h^*(\mathbf{v}(f^*\omega)).$$

This vector field is called the *push-forward* of $\mathbf{v}$ under f (with respect to h).

If $y \in Y$ and if we take the inductive limit over all étale neighborhoods of y in Y (see Definition 7.7), we get a push-forward $(f_*\mathbf{v})(y) \in \text{Hom}_{\mathcal{O}_{Y,y}}(\Omega_{Y/S,y}^1, \mathcal{O}_{Y,y})$ on stalks

$$(f_*\mathbf{v})_y(\omega_y) := (f_*\mathbf{v})(\omega)(y) := (f_*\mathbf{v})(y)(\omega) = h_y^*(\mathbf{v}(f^*\omega)(h(y))) = h_y^*(\mathbf{v}_{h(y)}(f^*\omega)_{h(y)}),$$

where $h_y^* : \mathcal{O}_{X,h(y)} \rightarrow \mathcal{O}_{Y,y}$ is the canonical map.

7.18 Remark On stalks there is also a well-defined notion of push-forward for sections of smooth and separated morphism. In order to see this, let $\pi : X \rightarrow Y$ be a smooth and separated morphism of S -schemes, and let $s : Y \hookrightarrow X$ be a section of π . In particular, s is a closed immersion. As already described above, every vector field $\mathbf{v} \in T_{Y/S}$ gives rise to a $\mathcal{O}_Y$ -linear morphism

$$\begin{array}{ccccccc} \widetilde{(s_*\mathbf{v})} : & s^*\Omega_{X/S}^1 & \xrightarrow{s^*} & \Omega_{Y/S}^1 & \xrightarrow{\mathbf{v}} & \mathcal{O}_Y, \\ & \omega \otimes 1 & \mapsto & s^*\omega & \mapsto & \mathbf{v}(s^*\omega) \end{array}$$

which corresponds to a $\mathcal{O}_X$ -linear morphism $\Omega_{X/S}^1 \rightarrow s_*\mathcal{O}_Y$. If $x = s(y)$ is in the image of s , then we get on stalks at x an $\mathcal{O}_{X,x}$ -linear morphism

$$\Omega_{X/S,x}^1 \longrightarrow (s_*\mathcal{O}_Y)_x = \mathcal{O}_{Y,y}.$$

The last equality is due to the fact that

$$(s_*\mathcal{O}_Y)_x = \begin{cases} 0 & \text{if } x \notin s(Y) \\ \mathcal{O}_{Y,y} & \text{if } x = s(y) \end{cases},$$

because s is a closed immersion (see e.g. [Liu], Prop. 2.2.24). Due to the flatness of π , the canonical morphism $\pi_x^* : \mathcal{O}_{Y,y} \rightarrow \mathcal{O}_{X,x}$ is injective, because flat morphisms of local rings are injective. Thus we may embed $\mathcal{O}_{Y,y}$ into $\mathcal{O}_{X,x}$ and finally get a $\mathcal{O}_{X,x}$ -linear homomorphism

$$(s_*\mathbf{v})_x : \Omega_{X/S,x}^1 \longrightarrow \mathcal{O}_{X,x}.$$

This gives a well-defined stalkwise push-forward of vector fields for sections s of smooth and separated morphisms. Again we obtain the formula

$$(s_*\mathbf{v})_x = \pi_x^* \circ \mathbf{v}_y \circ s_y^*,$$

where $x = s(y)$.¹ In particular we may push-forward vector fields along constant morphism. Thereby an S -morphism $f : X \rightarrow Y$ of S -schemes is called *constant*, if there is a section $s : S \hookrightarrow Y$ such that $f = s \circ \pi$, where $\pi : X \rightarrow S$ is the canonical morphism.

7.19 Definition Let $f : X \rightarrow Y$ be an isomorphism of smooth S -schemes. Let $\mathbf{v} : Y \rightarrow T_{Y/S}$ be a vectorfield on Y and $\omega \in \Omega_{X/S}^1$ a covectorfield on X . Then we define:

- a) The pull-back $f^*\mathbf{v}$ of $\mathbf{v}$ to X to be the push-forward $(f^{-1})_*\mathbf{v}$.
- b) The push-forward $f_*\omega$ of ω to Y to be the pull-back $(f^{-1})^*\omega$.

7.20 Definition Let X and Y be smooth S -schemes. A *r -times covariant, s -times contravariant (relative) tensor field* on Y over S is a global section of

$$(\mathcal{T}_{Y/S})_s := \underbrace{(\Omega_{Y/S}^1)^\vee \otimes \dots \otimes (\Omega_{Y/S}^1)^\vee}_{r\text{-times}} \otimes \underbrace{\Omega_{Y/S}^1 \otimes \dots \otimes \Omega_{Y/S}^1}_{s\text{-times}}.$$

Let $f : X \rightarrow Y$ be an S -morphism and $h : Y \hookrightarrow X$ a local section of f (where local is meant with respect to étale topology on Y). Then, whenever the following makes sense, we define:

- a) Let $\mathbf{t} \in (\mathcal{T}_{X/S})_s^r$, $\mathbf{t} = \mathbf{v}_1 \otimes \dots \otimes \mathbf{v}_r \otimes \omega^1 \otimes \dots \otimes \omega^s$, be a tensor field, and let $\nu^1, \dots, \nu^r \in \Omega_{Y/S}^1$ and let $\mathbf{w}_1, \dots, \mathbf{w}_s \in (\Omega_{Y/S}^1)^\vee$. Then

$$(f_*\mathbf{t})(\nu^1, \dots, \nu^r, \mathbf{w}_1, \dots, \mathbf{w}_s) := \prod_{i=1}^r h^*(\mathbf{v}_i(f^*\nu^i)) \cdot \prod_{j=1}^s \mathbf{w}_j(h^*\omega^j)$$

is called the *push-forward* of the tensor field $\mathbf{t}$.

¹Viewing an element $\alpha \in \mathcal{O}_X$ as a function $\alpha : X \rightarrow \prod_{x \in X} \mathcal{O}_{X,x}$, $x \mapsto \alpha_x$, we may even write

$$(s_*\mathbf{v})(\omega) = (\mathbf{v}(s^*\omega)) \circ \pi \quad \text{for all } \mathbf{v} \in \mathcal{T}_{Y/S}, \omega \in \Omega_{X/S}^1.$$

This may be seen as follows. As we are working on stalks, we may assume that X and Y are affine, say $X = \text{Spec } B$ and $Y = \text{Spec } A$. For $x \in X$ and $y = \pi(x) \in Y$ let B_x and A_y be the localization of B and A at the corresponding prime ideals. Then there is a canonical commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{\pi^*} & B \\ \downarrow & & \downarrow \\ A_y & \hookrightarrow & B_x \end{array} \quad \begin{array}{ccc} \alpha & \xrightarrow{\quad} & \pi^*(\alpha) \\ \downarrow & & \downarrow \\ \alpha(\pi(x)) & \xrightarrow{\quad} & \pi^*(\alpha)(x) \end{array}$$

As $A_y \rightarrow B_x$ is injective, we may identify A_y with a subring of B_x , whence yielding the desired equation $\pi^*(\alpha) = \alpha \circ \pi$ of functions from X to $\prod_{x \in X} \mathcal{O}_{X,x}$.

- b) Let $\mathbf{t} \in (\mathcal{T}_{Y/S})_s^0$, $\mathbf{t} = \omega^1 \otimes \dots \otimes \omega^s$, be a tensor field consisting of tensor powers of differential forms, and let $\mathbf{w}_1, \dots, \mathbf{w}_s \in (\Omega_{X/S}^1)^\vee$. Then

$$(f^*\mathbf{t})(\mathbf{w}_1, \dots, \mathbf{w}_s) := \prod_{j=1}^s \mathbf{w}_j(f^*\omega^j)$$

is called the *pull-back* of the tensor field $\mathbf{t}$.

- c) Let $\mathbf{t} \in (\mathcal{T}_{Y/S})_s^r$ and let f be an isomorphism. Then $f^*\mathbf{t} := (f^{-1})_*\mathbf{t}$ is called the *pull-back* of the tensor field $\mathbf{t}$.

7.2.1. Some formulas

Within this subsection let us collect some formulas concerning the pull-back or push-forward with respect to some types of morphisms. More precisely let us consider

- a) constant morphism.
b) morphisms of the type (f, g) or $f \times g$ involving fibre products.

7.21 Proposition *Let $f : X \rightarrow Y$ be a morphism of smooth S -schemes which allows a section $h : Y \hookrightarrow X$. Let $S' \rightarrow S$ be an unramified morphism. Assume that there is a factorization*

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ p \downarrow & \nearrow s & \\ S' & & \end{array}$$

Then $f^\omega = 0$ for all $\omega \in \Omega_{Y/S}^1$. In particular, $f_*\mathbf{v} = 0$ for all $\mathbf{v} \in \Gamma(X, T_{X/S})$, and this equation holds for any choice of the section h . (Recall that $f_*\mathbf{v}$ denotes the push-forward of $\mathbf{v}$ under f with respect to h in the sense of Definition 7.17.)*

PROOF $(f_*\mathbf{v})(\omega) = h^*(\mathbf{v}(f^*\omega))$ for all $\omega \in \Omega_{Y/S}^1$. Thus it suffices to show that $f^*\omega = 0$ for all $\omega \in \Omega_{Y/S}^1$. The above diagram induces a commutative diagram

$$\begin{array}{ccc} f^*\Omega_{Y/S}^1 & \longrightarrow & \Omega_{X/S}^1 \\ \parallel & & \uparrow \\ p^*(s^*\Omega_{Y/S}^1) & \longrightarrow & \pi^*\Omega_{S'/S}^1 \end{array}$$

By Proposition 1.9, $\Omega_{S'/S}^1 = 0$ and we are done. $\square$

The “adelic” space-time points are elements of $X(S)$. Thus we see that constant maps $X(S) \rightarrow Y(S)$ have zero differential (as we are used to from differential geometry). Let us illustrate this in more detail

7.22 Definition Let $f : X \rightarrow Y$ be a morphism of smooth S -schemes.

- a) f is called *constant*, if there exists a factorization

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ p \downarrow & \nearrow s & \\ S & & \end{array}$$

where $s : Y \hookrightarrow Y$ is a section of the structure morphism $Y \rightarrow S$.

- b) f is called *locally constant*, if any point $y \in Y$ possesses an étale neighborhood V , such that $f \times \text{id} : X \times_Y V \rightarrow V$ is constant.

7.23 Proposition Let $f : X \rightarrow Y$ a smooth morphism of smooth S -schemes. Let S be a Dedekind scheme and let X and Y be Néron models of their generic fibre. Let y be a physical point of Y over $s \in S$ which lies in the image of f . Let $x \in X$ be a physical point which is mapped to y under f (such a point exists by Proposition 7.13).

Then x extends to an S -valued point of X , whose image under f is an S -valued point $\alpha : S \hookrightarrow Y$ of Y containing y . In particular, there exists a constant morphism $g : X \rightarrow Y$ which maps X to α . Furthermore, there exists an étale environment $V \rightarrow Y$ of y , such that the restriction $g|_V = g \times \text{id}_V : X \times_Y V \rightarrow V$ of g induces a vanishing push-forward:

$$(g|_V)_* \mathfrak{v} = 0 \quad \text{for all } \mathfrak{v} \in \Gamma(X \times_Y V, T_{X \times_Y V/S}).$$

With some abuse of notation, we denote the constant morphism g by $y : X \rightarrow Y$.

PROOF As we are working with respect to étale topology, the local rings are strictly henselian. Let us first prove that x lifts to an S -valued point of X . If s is the generic point of S , this is true due to the Néron mapping property (Definition 6.1). Therefore let s be a closed point of S . Then x lifts to a $\mathcal{O}_{S,s}$ -valued point x' of X by Proposition 1.20 which itself may be lifted to an S -valued point β of X (again apply Néron mapping property in order to extend the point of x' which lies over the generic point of S). Due to the uniqueness assertion of the Néron mapping property, the point of the image of β which lies over s is x . If $p : X \rightarrow S$ is the structure morphism, $\alpha \circ p$ is the searched constant morphism, where $\alpha := f \circ \beta$.

By Proposition 7.9, there exists an étale open environment $V \rightarrow Y$ of y , such that $f|_V = f \times \text{id} : X \times_Y V \rightarrow V$ admits a section $h|_V$. By construction, $h|_V$ is also a section of $g|_V$. Thus the push-forward of vector fields under $g|_V$ exists. In order to prove that it is zero, it suffices to show that the corresponding pull-back of differential forms is zero, because $(g|_V)_* \mathfrak{v} = (h|_V)^* \circ \mathfrak{v} \circ (g|_V)^*$. Like in the proof of Proposition 7.21, the diagram

$$\begin{array}{ccc} X \times_Y V & \longrightarrow & V \\ \downarrow & & \downarrow \varphi_V \\ X & \longrightarrow & Y \\ \downarrow & \nearrow & \\ S & & \end{array}$$

shows that the pull-back of differential forms $(g|_V)^*$ is zero. Thereby one has to use that $(\varphi_V)^*$ is an isomorphism, because φ_V is étale. $\square$

7.24 Proposition *Let $f : X \rightarrow Y$, $g : X \rightarrow Z$ be smooth morphisms of S -schemes. Let $x \in X$ be a physical point and also denote by $f(x)$ (resp. $g(x)$) the constant map $X \rightarrow Y$, $q \mapsto f(x)$ (resp. $X \rightarrow Z$, $q \mapsto g(x)$) (see Proposition 7.23). Then in the stalk at $(f(x), g(x)) = (f, g)(x)$ there holds the equation*

$$(f, g)_* = (f(x), g)_* + (f, g(x))_*.$$

PROOF We have a commutative diagram

$$\begin{array}{ccccc} & & Y & & \\ & \nearrow f & \uparrow p_1 & & \\ X & \xrightarrow{(f, g)} & Y \times_S Z & & \\ & \searrow g & \downarrow p_2 & & \\ & & Z & & \end{array}$$

Therefore, we also get a commutative diagram

$$\begin{array}{ccc} (f, g)^* \Omega_{Y \times_S Z/S}^1 & \xrightarrow{\quad} & \Omega_{X/S}^1 \\ \wr \uparrow & & \uparrow \\ (f, g)^* (p_1^* \Omega_{Y/S}^1 \oplus p_2^* \Omega_{Z/S}^1) & \xlongequal{\quad} & f^* \Omega_{Y/S}^1 \oplus g^* \Omega_{Z/S}^1 \\ \\ p_1^* \omega_1 + p_2^* \omega_2 & \xrightarrow{\quad} & f^* \omega_1 + g^* \omega_2 \\ \uparrow & & \uparrow \\ (\omega_1, \omega_2) & \xlongequal{\quad} & (\omega_1, \omega_2) \end{array}$$

Thus we can write:

$$(f, g)^*(\omega_1, \omega_2) = f^* \omega_1 + g^* \omega_2.$$

In particular, this equation holds in the stalk at $(f(x), g(x))$. But in the stalk at $(f(x), g(x))$, we obtain analogously equations

$$\begin{aligned} (f(x), g)^*(\omega_1, \omega_2) &= f(x)^* \omega_1 + g^* \omega_2 = g^* \omega_2 \quad \text{and} \\ (f, g(x))^*(\omega_1, \omega_2) &= f^* \omega_1 + g(x)^* \omega_2 = f^* \omega_1. \end{aligned}$$

Thereby, we made use of the fact that the pull-back of differential forms under constant morphisms is zero (Proposition 7.21). This finishes the proof. $\square$

7.25 Proposition *Let $f : X \rightarrow Y$ and $g : Z \rightarrow T$ be smooth morphisms of S -schemes. Let $x \in X$ and $z \in Z$ be physical points and also denote by $f(x)$ (resp. $g(z)$) the constant maps $X \rightarrow Y$, $q \mapsto f(x)$ (resp. $Z \rightarrow T$, $q \mapsto g(z)$). Then in the stalk at $(f(x), g(z)) = (f \times g)(x, z)$ there holds the equation:*

$$(f \times g)_* = (f(x) \times g)_* + (f \times g(z))_*.$$

PROOF We have a commutative diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \pi_1 \uparrow & & \uparrow p_1 \\ X \times_S Z & \xrightarrow{f \times g} & Y \times_S T \\ \pi_2 \downarrow & & \downarrow p_2 \\ Z & \xrightarrow{g} & T \end{array}$$

Therefore, we also get a commutative diagram

$$\begin{array}{ccc} (f \times g)^* \Omega_{Y \times_S T/S}^1 & \xrightarrow{\quad} & \Omega_{X \times_S Z/S}^1 \\ \wr \uparrow & & \uparrow \wr \\ (f \times g)^*(p_1^* \Omega_{Y/S}^1 \oplus p_2^* \Omega_{T/S}^1) & \xrightarrow{\quad} & \pi_1^* \Omega_{X/S}^1 \oplus \pi_2^* \Omega_{Z/S}^1 \\ \parallel & & \\ \pi_1^* f^* \Omega_{Y/S}^1 \oplus \pi_2^* g^* \Omega_{T/S}^2 & & \end{array}$$

$$\begin{array}{ccc} p_1^* \omega_1 + p_2^* \omega_2 & \xrightarrow{\quad} & (f \times g)^* p_1^* \omega_1 + (f \times g)^* p_2^* \omega_2 \\ \uparrow & & \uparrow \\ (\omega_1, \omega_2) & \xrightarrow{\quad} & (f^* \omega_1, g^* \omega_2) \\ \parallel & & \\ (\omega_1, \omega_2) & & \end{array}$$

Thus we can write:

$$(f \times g)^*(\omega_1, \omega_2) = (f^* \omega_1, g^* \omega_2) = \pi_1^* f^* \omega_1 + \pi_2^* g^* \omega_2$$

In particular, this equation holds in the stalk at $(f \times g)(x, z) = (f(x), g(z))$. But in the stalk at $(f(x), g(z))$, we obtain analogously equations

$$\begin{aligned} (f(x) \times g)^*(\omega_1, \omega_2) &= \pi_1^* f(x)^* \omega_1 + \pi_2^* g^* \omega_2 = \pi_2^* g^* \omega_2 \quad \text{and} \\ (f \times g(z))^*(\omega_1, \omega_2) &= \pi_1^* f^* \omega_1 + \pi_2^* g(z)^* \omega_2 = \pi_1^* f^* \omega_1. \end{aligned}$$

Thereby, we made use of the fact that the pull-back of differential forms under constant morphisms is zero (Proposition 7.21). This finishes the proof. $\square$

8. Gauge theory

8.1. The classical, differential geometric theory

The fundamental object underlying classical, differential geometric Yang-Mills theory are principal bundles. Before we state the definition of the algebraic geometric analogue of principle bundles in the next section 8.2, let us briefly recall some differential geometric notions which are essential for gauge theory. Throughout this section let G be a Liegroup, and let P be a differentiable manifold.

8.1 Definition Let $\psi : P \times G \rightarrow P$ be a differentiable map and denote by $\psi_g : P \rightarrow P$ resp. $\psi_p : G \rightarrow P$ the induced maps $\psi_g(\cdot) := \psi(\cdot, g)$ resp. $\psi_p(\cdot) := \psi(p, \cdot)$ where $g \in G$ and $p \in P$. Let us also introduce the abbreviation $p \cdot g := \psi(p, g)$. Then ψ is called a *right group action* if and only if:

- a) $\psi_g : P \rightarrow P$ is a diffeomorphism for all $g \in G$.
- b) $\psi_a \circ \psi_b(p) = \psi_{ba}(p)$; i.e.: $(p \cdot b) \cdot a = p \cdot (b \cdot a)$.

8.2 Definition A group action $\psi : P \times G \rightarrow P$ is called *free*, if and only if the following holds: If there exists a $p \in P$ and a $g \in G$ such that $p \cdot g = p$, then $g = e$ is the unit element of G .

8.3 Proposition Let G act on P on the right by means of $\psi : P \times G \rightarrow P$. Let $A_e \in \mathfrak{g}$, where $\mathfrak{g}$ denotes the Lie-algebra of G , and let $\exp : \mathfrak{g} \rightarrow G$ be the exponential map. Then A_e induces a vectorfield $(A_e)_*$ on P , the so called *killing-vectorfield* or *fundamental vectorfield*. More precisely:

$$((A_e)_*f)(p) = \left. \frac{d}{dt} \right|_{t=0} (f \circ \psi_{\exp(tA_e)}(p))$$

for all $f \in C^\infty(P)$ and $p \in P$.

8.4 Remark Let G act on P on the right by means of $\psi : P \times G \rightarrow P$. Denote by $0 \in \Gamma(P, TP)$ the zero section of tangent bundle TP of P . Let $A_e \in \mathfrak{g}$ and denote by A the unique left invariant vectorfield on G corresponding to A_e . Then we may consider the vectorfield $0 \times A$ living on $P \times G$, and we get for all $p \in P$:

$$((\psi_p)_*A_e)(p) = ((A_e)_*)(p) = (\psi_*(0 \times A))(p),$$

where $(\psi_*(0 \times A))(p) = ((\psi_*)_{(p \cdot g^{-1}, g)}(0_{p \cdot g^{-1}} \times A_g))(p)$ for all $g \in G$.

PROOF Denote by $\psi_p : G \rightarrow P$, $g \mapsto \psi_p(g) := p \cdot g$ the translation by p . For all $g \in G$ let $A_g := A(g) \in T_g G$ be the tangent vector of A at g , and consider the integral curve $g(t) := g \cdot \exp(tA_e) = L_g \circ \exp(tA_e)$ representing A_g (where L_g is the left translation by g). Then we obtain

$$\left(((\psi_p)_* A_g)(f) \right)(p \cdot g) = \frac{d}{dt} \Big|_{t=0} f \circ \psi_p \circ g(t)$$

for all $f \in C^\infty(P)$. In particular

$$\left(((\psi_p)_* A_e)(f) \right)(p) = \frac{d}{dt} \Big|_{t=0} f \circ \psi_p(\exp(tA_e)) = \frac{d}{dt} \Big|_{t=0} f \circ \psi_{\exp(tA_e)}(p) = ((A_e)_* f)(p).$$

This shows the first equality. On the other hand, $(0 \times A)(p, g) = 0_p \times A_g$ for all $(p, g) \in P \times G$. If $p(t)$ is an integral curve of 0_p , i.e. $p(t) = p$ for all $t \in \mathbb{R}$, then we get for all $f \in C^\infty(P)$:

$$\begin{aligned} \left(\psi_*(0_p \times A_g)(f) \right)(p \cdot g) &= \frac{d}{dt} \Big|_{t=0} f \circ \psi(g(t), p(t)) = \frac{d}{dt} \Big|_{t=0} f \circ \psi(g(t), p) \\ &= \frac{d}{dt} \Big|_{t=0} f \circ \psi_p(g(t)) = ((\psi_p)_* A_g)(f)(p \cdot g). \end{aligned}$$

As ψ is a group action, we know that $(p \cdot g) \cdot h = p \cdot (gh)$ for all $p \in P$ and $g, h \in G$. It follows that

$$\begin{aligned} \psi_{p \cdot h} \circ (h^{-1}g)(t) &= \psi_{p \cdot h}(h^{-1}g \cdot \exp(tA_e)) = (p \cdot h) \cdot (h^{-1}g \cdot \exp(tA_e)) \\ &= p \cdot (g \cdot \exp(tA_e)) = \psi_p \circ g(t), \end{aligned}$$

and thus we obtain

$$((\psi_p)_* A_g)(f)(p \cdot g) = \frac{d}{dt} \Big|_{t=0} f \circ \psi_p(g(t)) = \frac{d}{dt} \Big|_{t=0} f \circ \psi_{p \cdot g}(\exp(tA_e)) = ((A_e)_* f)(p \cdot g).$$

All in all we see that

$$\left(\psi_*(0_p \times A_g) \right)_{p \cdot g}(f) = ((A_e)_*)_{p \cdot g}(f) = \left(\psi_*(0_{p \cdot h} \times A_{h^{-1}g}) \right)_{p \cdot g}(f)$$

for all $p \in P$ and $g, h \in G$. Thus it indeed makes sense to write

$$\left(\psi_*(0 \times A) \right)(p \cdot g) = \psi_*(0_p \times A_g).$$

□

8.2. Torsors

Now we are going to transfer these structures to algebraic geometry, in order to provide the basis for a Yang-Mills theory over arbitrary commutative rings. Let G be a group scheme over X and P and X -scheme. Then an action of G on P is a X -morphism

$$\psi : P \times_X G \rightarrow P$$

that induces an action of the group $G(T)$ on the set $P(T)$ for all S -scheme T (especially $(p \cdot g) \cdot h = p \cdot (g \cdot h)$ for all $p \in P(T)$, $g, h \in G(T)$).

8.5 Proposition *Let G act on a scheme P . Let $\pi : P \rightarrow X$ be the structure morphism. Then it is equivalent:*

- a) *The scheme P is faithfully flat and locally of finite-type over X and there is an isomorphism*

$$P \times_X G \xrightarrow{\sim} P \times_X P, \quad (p, g) \mapsto (p, p \cdot g).$$

- b) *There is a covering $(U_i \rightarrow X)$ for the flat topology¹ on X , such that, for each i , there is an isomorphism*

$$\chi_i : \pi^{-1}(U_i) := P_{U_i} := U_i \times_X P \rightarrow U_i \times_X G =: G_{U_i}$$

respecting the G_{U_i} -action.

PROOF [Milne], Chap. III, Prop. 4.1 □

8.6 Remark

- Let G be a smooth group scheme. Then any torsor under G is smooth (see Proposition 8.8). Using the fact that any X -torsor P under G , which allows a X -section, is trivial, Proposition 1.19 shows that we may choose the covering $(U_i \rightarrow X)$ in b) for the étale topology. This is important in regard to physical applications as we already saw in section 7.1 that we have to work with respect to étale topology in order to incorporate the principle of general covariance. In physical situations we will furthermore assume that there exists a covering $(U_i \rightarrow X)$ consisting of only finitely many maps $U_i \rightarrow X$. This will allow us to glue sheaves. Then the global and the family of local versions of a physical theory are equivalent (as one would expect).
- Condition b) says that there are commutative diagrams

$$\begin{array}{ccc} \pi^{-1}(U_i) & \xrightarrow{\chi_i} & U_i \times_X G \\ & \searrow \pi|_{U_i} := \text{id}_{U_i} \times \pi & \swarrow pr_1 \\ & U_i & \end{array}$$

$$\begin{array}{ccccc} P_{U_i} \times_{U_i} G_{U_i} & \equiv & U_i \times_X P \times_X G & \xrightarrow{\chi_i \times \text{id}} & U_i \times_X G \times_X G & \equiv & G_{U_i} \times_{U_i} G_{U_i} \\ & & \downarrow \text{id} \times \psi & & \downarrow \text{id} \times m & & \\ & & U_i \times_X P & \xrightarrow{\chi_i} & U_i \times_X G & & \end{array}$$

¹See Definition 7.4 for the notion of a Grothendieck topology.

where $m : G \times_X G \rightarrow G$ denotes the group multiplication. Evaluated at T -valued points p of $\pi^{-1}(U_i)$ and g of G (where T is an S -scheme), the latter commutative diagram may be stated as follows:

$$(pr_2 \circ \chi_i)(p \cdot g) = (pr_2 \circ \chi_i(p)) \cdot g.$$

Thereby pr_2 denotes the canonical projection $U_i \times_X G \rightarrow G$.

- Condition *a*) implies that the group action $\psi : P \times_X G \rightarrow P$ is free (see Definition 8.2) and transitive on fibres over X . This may be seen as follows: The morphism $P \times_X G \xrightarrow{\sim} P \times_X P$, $(p, g) \mapsto (p, p \cdot g)$ is an isomorphism by definition. Due to injectivity, the preimage of a pair $(p, p \cdot g)$ with $p \cdot g = p$ consists of the pair (p, e) , where e denotes the unit element of G . This shows that the action is free. The surjectivity implies that for all $p, p' \in P$ with $\pi(p) = \pi(p')$ (i.e. for all $(p, p') \in P \times_X P$) there exists a $g \in G$ such that $p' = pg$.

8.7 Definition A scheme P , on which acts G , and that satisfies the equivalent statements of Proposition 8.5 is called a *principal homogeneous space* or an *X -torsor under G* .

In physical applications we are going to consider smooth schemes provided with étale topology. Thus we see that the notion of torsors is exactly corresponds to the G -principal bundles in ordinary real-valued differential geometry.

8.8 Proposition *If G is smooth respectively étale over X , then so also any X -torsor under G .*

PROOF [Milne], Chap. III, Prop. 4.2 □

8.9 Proposition *Let us denote the set of all isomorphism classes of X -torsors under G by $PHS(G/X)$. Then there is a canonical injection $PHS(G/X) \hookrightarrow \check{H}^1(X_{fl}, G)$. This map is an isomorphism if G is affine over X (i.e. the preimage of every affine subscheme of X is an affine subscheme of G).*

PROOF Let us shortly summarize how a X -torsor P under G gives rise to a 1-cocycle. For the details of the proof see [Milne], Chap. III, Cor. 4.7. As we are interested in physical applications let us directly work with étale topology. For this purpose we assume that G is smooth. Then P is smooth, too, by Proposition 8.8, and due to Proposition 1.19 we may assume that there exists an étale covering $\mathfrak{U} := (U_i \rightarrow X)$ that trivializes P , so that $P(U_i)$ is non-empty for all i . Then choose a section $s_i \in P(U_i)$ for all i . As the chosen covering trivializes P , there exist unique $g_{ij} \in G(U_{ij})$ (where $U_{ij} := U_i \times_X U_j$), such that $s_i|_{U_{ij}} \cdot g_{ij} = s_j|_{U_{ij}}$. Omitting restriction signs we therefore obtain $s_i g_{ij} g_{jk} = s_k = s_i g_{ik}$, and so $g_{ij} g_{jk} = g_{ik}$, because the G -action on P is free. One says that the family (g_{ij}) is a 1-cocycle. If $s'_i \in P(U_i)$ is another family of sections, then there is a family (h_i) , $h_i \in G(U_i)$, such that $s_i = s'_i \cdot h_i$ for all i , because the chosen covering trivializes P . Again omitting

restriction signs, we get $s'_i h_i g_{ij} = s_i g_{ij} = s_j = s'_j h_j = s'_i g'_{ij} h_j$, and so $g'_{ij} = h_i g_{ij} h_j^{-1}$. One says that (g_{ij}) and (g'_{ij}) are cohomologous 1-cocycles. The property of 1-cocycles to be cohomologous is an equivalence relation, and the set of cohomology classes is denoted by $\check{H}^1(\mathfrak{U}, G)$. Of course the cohomology class is unaltered if P is replaced by an isomorphic torsor. Thus P defines an element of $\check{H}^1(X_{\text{ét}}, G)$ which is defined to be $\varinjlim \check{H}^1(\mathfrak{U}, G)$ where the limit runs over all coverings $\mathfrak{U}$ obtained by refinement. $\square$

8.10 Translations by S -valued points on torsors.

Let X be a smooth S -scheme, let G be a smooth S -group scheme and let $G_X := G \times_S X$. Let $\psi : P \times_X G_X \rightarrow P$ be the group action of G_X on a X -torsor P under G_X . Consider an S -valued point $\alpha : S \hookrightarrow P$ of P . Let $f : P \rightarrow P$ and $g : P \rightarrow G$ be S -morphisms. Then $f_\alpha := f \circ \alpha$ and $g_\alpha := g \circ \alpha$ are S -valued points, too.

The group action induces two kinds of translation by S -valued points. First, we consider the translation $\psi_{f(\alpha)}$ by the S -valued point f_α of P :

$$\psi_{f(\alpha)} : G \xrightarrow{(\pi, \text{id})} S \times_S G \xrightarrow{f_\alpha \times \text{id}} P \times_S G \xrightarrow{\psi} P.$$

In formulas we may write $\psi_{f(\alpha)} := \psi \circ (f_\alpha \times \text{id}) \circ (\pi, \text{id}) = \psi \circ (f_\alpha \circ \pi, \text{id})$, where $\pi : G \rightarrow S$ is the canonical morphism. If $f(\alpha)$ denotes the constant map making the diagram

$$\begin{array}{ccc} G & \xrightarrow{f(\alpha)} & P \\ \downarrow \pi & \nearrow f \circ \alpha & \\ S & & \end{array}$$

commutative, then we may state the translation $\psi_{f(\alpha)}$ in the following way:

$$\psi_{f(\alpha)} = \psi \circ (f(\alpha), \text{id}) : G \rightarrow P.$$

Analogously we define the translation $\psi_{g(\alpha)}$ by the S -valued point g_α of G :

$$\psi_{g(\alpha)} : P \xrightarrow{(\text{id}, \pi)} P \times_S S \xrightarrow{\text{id} \times g_\alpha} P \times_S G \xrightarrow{\psi} P,$$

where π is this time the canonical morphism $P \rightarrow S$. So we get $\psi_{g(\alpha)} := \psi \circ (\text{id} \times g_\alpha) \circ (\text{id}, \pi) = \psi \circ (\text{id}, g_\alpha \circ \pi)$. Using the constant map $g(\alpha)$ making the diagram

$$\begin{array}{ccc} P & \xrightarrow{g(\alpha)} & G \\ \downarrow \pi & \nearrow g \circ \alpha & \\ S & & \end{array}$$

commutative, we arrive at the identity

$$\psi_{g(\alpha)} = \psi \circ (\text{id}, g(\alpha)) : P \rightarrow P.$$

Furthermore, if $h_P : P \rightarrow P$ and $h_G : P \rightarrow G$ are morphisms of S -schemes, we get identities:

$$\begin{aligned} \psi_{f(\alpha)} \circ h_G &= \psi \circ (f(p), h_G). \\ \psi_{g(\alpha)} \circ h_P &= \psi \circ (h_P, g(p)). \end{aligned}$$

8.3. Gauge transformations

Throughout this section let X be a smooth S -scheme, and let G be a smooth X -group scheme.

8.11 Definition Let P be a X -torsor P under G . A X -automorphism $\hat{\theta} : P \rightarrow P$ is called a *gauge transformation* or a *vertical automorphism*, if $\hat{\theta}$ is compatible with the group action ψ of G on P ; i.e., there are the following commutative diagrams.

$$\begin{array}{ccc} P \times_X G & \xrightarrow{\psi} & P \\ \hat{\theta} \times id \downarrow & & \downarrow \hat{\theta} \\ P \times_X G & \xrightarrow{\psi} & P \end{array} \quad \hat{\theta} : \begin{array}{ccc} P & \xrightarrow{\sim} & P \\ & \searrow & \swarrow \\ & X & \end{array}$$

We denote by the group of gauge transformations by $\mathcal{G}$.

Furthermore, let us call a morphism $u : P \rightarrow G$ *equivariant*, if $u(p \cdot g) = g^{-1}u(p)g$ for all T -valued points $p \in P(T)$ and $g \in G(T)$ and for all S -schemes T .

8.12 Proposition *There is a one-to-one correspondence between gauge transformations and equivariant X -morphisms $u : P \rightarrow G$.*

PROOF As usual let us denote the formation of the inverse respectively the multiplication on the group scheme G by ι respectively m . Let assume that an equivariant X -morphisms u is given. Then we define the following X -morphism $\hat{\theta}_u$.

$$\begin{array}{ccccc} \hat{\theta}_u : & P & \xrightarrow{(id, u)} & P \times_X G & \xrightarrow{\psi} & P \\ & p & \mapsto & (p, u(p)) & \mapsto & p \cdot u(p) \end{array}$$

Of course, $\hat{\theta}_u$ is a X -isomorphism with inverse $\hat{\theta}_{u^{-1}}$, where $u^{-1} := \iota \circ u : G \rightarrow G$. Furthermore $\hat{\theta}_u$ is compatible with the group action of G on P . This may be seen as follows: By assumption, $u(p \cdot g) = g^{-1}u(p)g$ and therefore

$$\hat{\theta}_u(p \cdot g) = (p \cdot g) \cdot u(p \cdot g) = (p \cdot g) \cdot (g^{-1}u(p)g) = (p \cdot u(p)) \cdot g = \hat{\theta}_u(p) \cdot g.$$

Thus $\hat{\theta}_u$ is indeed a gauge transformation.

Conversely, let now a gauge transformation $\hat{\theta}$ be given. Then we define the following X -morphism $u_{\hat{\theta}}$.

$$u_{\hat{\theta}} : P \xrightarrow{id \times \hat{\theta}} P \times_X P \xrightarrow{\sim} P \times_S G \xrightarrow{pr_2} G$$

Thereby $P \times_X P \xrightarrow{\sim} P \times_S G$ denotes the inverse of the canonical isomorphism induced by the group action ψ (see Proposition 8.5 a)). Let us show that this morphism is equivariant. By definition of $u_{\hat{\theta}}$, we have got

$$(p \cdot g, \hat{\theta}(p \cdot g)) = (p \cdot g, \hat{\theta}(p) \cdot g) = (p \cdot g, (p \cdot u_{\hat{\theta}}(p)) \cdot g) = (p \cdot g, (p \cdot g) \cdot (g^{-1}u_{\hat{\theta}}(p)g)) \in P \times_X P.$$

Thus the image of $p \cdot g$ under $u_{\hat{\theta}}$ is indeed $g^{-1}u_{\hat{\theta}}(p)g$. Furthermore, the given associations are inverse to each other:

$$p \cdot u_{\hat{\theta}_u}(p) = \hat{\theta}_u(p) = p \cdot u(p), \quad \text{i.e.} \quad u_{\hat{\theta}_u}(p) = u(p) \text{ for all } p,$$

because the group action is free. On the other hand

$$\hat{\theta}_{u_{\hat{\theta}}}(p) = p \cdot u_{\hat{\theta}}(p) = \hat{\theta}(p) \quad \text{for all } p. \quad \square$$

8.4. Horizontality

Throughout this section let X be a smooth S -scheme, and let G be a smooth S -group scheme. Furthermore let X and G be separated, let $G_X := G \times_S X$ and consider an X -torsor P under G_X (see Proposition 8.5). Let T be an arbitrary S -scheme. We already know from Proposition 8.8 that P is smooth and separated over X and therefore also over S . Denote by $\mathfrak{g} := (\varepsilon^* \Omega_{G/S}^1)(S)$ the Lie-algebra of G , where $\varepsilon : S \hookrightarrow G$ denotes the unit section.

By definition of torsors there is a commutative diagram

$$\begin{array}{ccccc} & & & P & \\ & & \nearrow \psi & \uparrow p_2 & \\ P \times_S G & \xlongequal{\quad} & P \times_X G_X & \xrightarrow[\pi_1 \times \psi]{\sim} & P \times_X P \\ & & \searrow \pi_1 & \downarrow p_1 & \\ & & & P & \end{array}$$

where π_i and p_i are the canonical projections of the respective schemes onto the i -th factor, and where $\pi_1 \times \psi$ is an isomorphism. As usual, ψ denotes the group action. Then the diagram shows that the diagonal $\Delta : P \rightarrow P \times_X P$ induces a canonical global section $s := (\pi_1 \times \psi)^{-1} \circ \Delta$ of ψ which is given by $p \mapsto (p, 1)$ on T -valued points. Therefore, by Remark 7.16, every global vector field $\mathfrak{v}$ on $P \times_X G$ may be pushed forward to a global vector field $\psi_* \mathfrak{v}$ on P which is given by

$$\psi_* \mathfrak{v} = s^* \circ \mathfrak{v} \circ \psi^*.$$

Drawing inspiration from Remark 8.4, we use this push-forward, in order to define the killing-vector field. For this purpose consider an element $\mathfrak{v}_e \in \mathfrak{g}$ and denote by $\mathfrak{v}$ the uniquely determined left-invariant vector field induced by $\mathfrak{v}_e$ (see Proposition 11.57). Furthermore, let $0 \in \Gamma(P, T_{P/S})$ be the vector field which is constant zero (i.e. the zero-section). Recalling the identity $T_{P \times_S G/S} = T_{P/S} \times_S T_{G/S}$ from Proposition 11.18 or as well the isomorphism $\pi_1^* \Omega_{P/S}^1 \oplus \pi_2^* \Omega_{G/S}^1 \xrightarrow{\sim} \Omega_{P \times_S G/S}^1$ from Proposition 1.7, we may interpret the cartesian product $0 \times \mathfrak{v}$ as a vector field on $P \times_X G$. Then we define a map

$$\begin{aligned} \tilde{\sigma} : \mathfrak{g} &\longrightarrow \Gamma(P, T_{P/S}) = \text{Hom}_{\mathcal{O}_P}(\Omega_{P/S}^1, \mathcal{O}_P), \\ \mathfrak{v}_e &\longmapsto \psi_*(0 \times \mathfrak{v}) \end{aligned}$$

which turns out to be $\mathcal{O}_S(S)$ -linear. By tensoring over $\mathcal{O}_S(S)$ with $\mathcal{O}_P$, $\tilde{\sigma}$ induces a $\mathcal{O}_P$ -linear map

$$\begin{aligned} \sigma : \mathcal{O}_P \otimes_{\mathcal{O}_S(S)} \mathfrak{g} &\longrightarrow \Gamma(P, T_{P/S}) = \text{Hom}_{\mathcal{O}_P}(\Omega_{P/S}^1, \mathcal{O}_P) \\ r \otimes \mathfrak{v}_e &\longmapsto r \cdot \psi_*(0 \times \mathfrak{v}) = r \cdot \tilde{\sigma}(\mathfrak{v}_e). \end{aligned}$$

8.13 Remark By Proposition 11.56, $\mathfrak{g}$ carries the structure of a $\mathcal{O}_S(S)$ -Lie-algebra by means of the $\mathcal{O}_S(S)$ -linear Lie-bracket $[\cdot, \cdot]$. Therefore we may endow $\mathcal{O}_P \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$ with the following structure of a $\mathcal{O}_P$ -Lie-algebra: For $r \in \mathcal{O}_P$ and $\mathfrak{v}_e \in \mathfrak{g}$ let

$$[r \otimes \mathfrak{v}_e, \cdot] : \mathcal{O}_P \otimes_{\mathcal{O}_S(S)} \mathfrak{g} \rightarrow \mathcal{O}_P \otimes_{\mathcal{O}_S(S)} \mathfrak{g}, \quad s \otimes \mathfrak{w}_e \mapsto rs \otimes [\mathfrak{v}_e, \mathfrak{w}_e].$$

This map is $\mathcal{O}_P$ -linear and induces a Lie-bracket on $\mathcal{O}_P \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$.

8.14 Proposition $\tilde{\sigma} : \mathfrak{g} \rightarrow \Gamma(P, T_{P/S})$ is $\mathcal{O}_S(S)$ -linear. In particular σ is $\mathcal{O}_P$ -linear.

PROOF Let $\mathfrak{v}_e, \mathfrak{w}_e \in \mathfrak{g}$. The canonical map $\varepsilon^* \Omega_{G/S}^1 \longrightarrow p_* \Omega_{G/S}^1$, $\mathfrak{v}_e \mapsto \mathfrak{v}$, extending sections in $\varepsilon^* \Omega_{G/S}^1$ to left invariant sections of $\Omega_{G/S}^1$, is a $\mathcal{O}_S(S)$ -linear map (as illustrated in the proof of Proposition 11.57 or simply by Proposition 11.58). In particular, we know that

$$\begin{aligned} \mathfrak{v}_e + \mathfrak{w}_e &\mapsto \mathfrak{v} + \mathfrak{w} & \text{and} \\ a \cdot \mathfrak{v}_e &\mapsto a \cdot \mathfrak{v} & \text{for all } a \in \mathcal{O}_S(S) \end{aligned}$$

under this map. Due to the decomposition $\pi_1^* \Omega_{P/S}^1 \oplus \pi_2^* \Omega_{G/S}^1 \xrightarrow{\sim} \Omega_{P \times_S G/S}^1$ of Proposition 1.7, we know that $(0 \times (\mathfrak{v} + \mathfrak{w})) = (0 \times \mathfrak{v}) + (0 \times \mathfrak{w})$. Thus we conclude that

$$\begin{aligned} \tilde{\sigma}(a\mathfrak{v}_e + a'\mathfrak{w}_e)(\omega) &= \psi_*(0 \times (a\mathfrak{v} + a'\mathfrak{w}))(\omega) \\ &= s^*((0 \times (a\mathfrak{v} + a'\mathfrak{w}))(\psi^*\omega)) \\ &= s^*((0 \times a\mathfrak{v})(\psi^*\omega)) + s^*((0 \times a'\mathfrak{w})(\psi^*\omega)) \\ &= s^*(a(0 \times \mathfrak{v})(\psi^*\omega)) + s^*(a'(0 \times \mathfrak{w})(\psi^*\omega)) \\ &= a s^*((0 \times \mathfrak{v})(\psi^*\omega)) + a' s^*((0 \times \mathfrak{w})(\psi^*\omega)) \\ &= a \tilde{\sigma}(\mathfrak{v}_e)(\omega) + a' \tilde{\sigma}(\mathfrak{w}_e)(\omega). \end{aligned}$$

for all $\omega \in \Omega_{P/S}^1$. This shows that $\tilde{\sigma}$ is $\mathcal{O}_S(S)$ -linear. As $\sigma(r \otimes \mathfrak{v}_e) = r \cdot \tilde{\sigma}(\mathfrak{v}_e)$ by definition, we are done. $\square$

8.15 Definition Let P be a X -torsor under G_X . Let $\mathfrak{v}_e \in \mathfrak{g}$. Then the vector field $\tilde{\sigma}(\mathfrak{v}_e) = \sigma(1 \otimes \mathfrak{v}_e) \in \Gamma(T_{P/S}/P)$ is called the *killing-vector field* of the group action.

Up to now, we considered the killing-vector field globally on all of P . Let us now, analyze the local properties of $\tilde{\sigma}(\mathfrak{v}_e)$.

By means of the canonical isomorphism stated in Proposition 11.30 a), let us interpret $\tilde{\sigma}(\mathfrak{v}_e)$ as a differential form. Locally (with respect to étale topology), P allows

a split $P = X \times_S G$. So, as we are now interested in a local consideration, let us assume that P splits and let

$$\begin{aligned}\rho_1 : X \times_S G &\rightarrow X \\ \rho_2 : X \times_S G &\rightarrow G\end{aligned}$$

be the canonical projections. Due to the isomorphism $\rho_1^* \Omega_{X/S}^1 \oplus \rho_2^* \Omega_{G/S}^1 \xrightarrow{\sim} \Omega_{P/S}^1$ (Proposition 1.7), we may decompose $\tilde{\sigma}(\mathbf{v}_e)$ into a direct sum $(\omega_1, \omega_2) \in \rho_1^* \Omega_{X/S}^1 \oplus \rho_2^* \Omega_{G/S}^1$. We claim that the component ω_1 is equal to zero. Therefore $\tilde{\sigma}(\mathbf{v}_e)$ lives only in the group part G of P and is killed if it is pushed forward to the base X using the structure morphism $\pi : P \rightarrow X$.

Let us call the X -component in the local split $P = X \times_S G$ the *horizontal* component, and let us call the group part G the *vertical* component. In particular, we see that (locally) the killing vector field lives in the vertical component of P . But before defining the notion of horizontality and verticality for vector fields, let us first introduce the notion of a connection form.

8.16 Definition Let P be an X -torsor under G_X , let $\sigma : \mathfrak{g} \rightarrow \Gamma(T_{P/S}/P)$ be the killing-form, and let d be the exterior differential (see Theorem 11.39). Furthermore, let ω be a global section ω of $\Omega_{P/S}^1 \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$. Then we introduce the following two $\mathcal{O}_P$ -linear morphisms $\text{hor}, \text{ver} \in \text{Hom}_{\mathcal{O}_P}(\Omega_{P/S}^1, \Omega_{P/S}^1)$:

$$\begin{aligned}\text{ver} &:= \sigma \circ \omega. \\ \text{hor} &:= \text{id} - \sigma \circ \omega.\end{aligned}$$

ω is called a *connection form* (or more shortly a *connection*) if and only if:

- a) $\omega \circ \sigma = \text{id} : \mathcal{O}_P \otimes_{\mathcal{O}_S(S)} \mathfrak{g} \rightarrow \mathcal{O}_P \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$.
- b) $\psi_a^* \omega = \text{Ad}(a^{-1})\omega$ for all S -valued points $a \in G(S)$.²
- c) $\omega(\text{hor } \mathbf{v}) = 0$ for all $\mathbf{v} \in \Gamma(T_{P/S}/P)$.
- d) $\omega : \mathcal{T}_{P/S} \rightarrow \mathcal{O}_P \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$ is a homomorphism of Lie-algebras.³

8.17 Definition A vector field $\mathbf{v} \in \Gamma(T_{P/S}/P)$ is called *horizontal* (resp. *vertical*), if $\mathbf{v} = \text{hor } \mathbf{v}$ (resp. $\mathbf{v} = \text{ver } \mathbf{v}$).

8.18 Proposition Let $\omega \in \Omega_{P/S}^1 \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$ be a connection form. Let ψ_a be the translation by an S -valued point a (induced by the group action ψ). Then:

² This equation has to be understood in the sense of $(\psi_a^* \omega)(\mathbf{v}) = \text{Ad}(a^{-1})(\omega(\mathbf{v}))$ for all $\mathbf{v} \in \Gamma(T_{P/S}/P)$, where ψ_a denotes the translation by the S -valued point a (see 8.10 above). The map Ad was introduced in subsection 11.4.3.

³ This means that $\omega([\mathbf{v}, \mathbf{w}]) = [\omega(\mathbf{v}), \omega(\mathbf{w})]$ for all vector fields $\mathbf{v}, \mathbf{w} \in \mathcal{T}_{P/S}$. Thereby $\mathcal{T}_{P/S}$ is considered as a Lie-algebra with the commutator as Lie-bracket (see Definition 11.41). The Lie-algebra structure of $\mathcal{O}_P \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$ was introduced in Remark 8.13.

$$\begin{aligned} a) \quad \text{hor} &= \text{hor} \circ \text{hor} & \text{hor} \circ \text{ver} &= 0. \\ \text{ver} &= \text{ver} \circ \text{ver} & \text{ver} \circ \text{hor} &= 0. \end{aligned}$$

b) $(\psi_a)_*\mathbf{v}$ is horizontal for all horizontal vector fields $\mathbf{v}$ on P .
 $(\psi_a)_*\mathbf{v}$ is vertical for all vertical vector fields $\mathbf{v}$ on P .

$$\begin{aligned} c) \quad \text{hor} \circ (\psi_a)_* &= (\psi_a)_* \circ \text{hor} \\ \text{ver} \circ (\psi_a)_* &= (\psi_a)_* \circ \text{ver} \end{aligned}$$

PROOF If ω is a connection form, then $\omega \circ \sigma = \text{id}$. Therefore $\text{ver} \circ \text{ver} = \sigma \circ \omega \circ \sigma \circ \omega = \sigma \circ \omega = \text{ver}$ and $\text{hor} \circ \text{hor} = (\text{id} - \text{ver}) \circ (\text{id} - \text{ver}) = \text{id} - 2 \cdot \text{ver} + \text{ver} \circ \text{ver} = \text{id} - \text{ver} = \text{hor}$. In particular, it follows that $\text{hor} \circ \text{ver} = (\text{id} - \text{ver}) \circ \text{ver} = 0 = \text{ver} \circ (\text{id} - \text{ver}) = \text{ver} \circ \text{hor}$. In order to prove b) consider first a horizontal vectorfield $\mathbf{v}$. Then

$$\begin{aligned} \text{ver}((\psi_a)_*\mathbf{v}) &= \sigma \circ \omega((\psi_a)_*\mathbf{v}) = \sigma((\psi_a^{-1})^*((\psi_a^*\omega)(\mathbf{v}))) = \sigma((\psi_a^{-1})^*((\text{Ad}(a^{-1})\omega)(\mathbf{v}))) \\ &= \sigma(\text{Ad}(a^{-1})(\psi_a^{-1})^*(\omega(\mathbf{v}))). \end{aligned}$$

Thus $\text{hor}((\psi_a)_*\mathbf{v}) = (\psi_a)_*\mathbf{v}$, because $\omega(\mathbf{v}) = 0$ for horizontal vector fields $\mathbf{v}$.

Let now be $\mathbf{v}$ be a vertical vector field, i.e. $\mathbf{v} = \text{ver}(\mathbf{v}) = \sigma(\omega(\mathbf{v}))$. Writing $\omega(\mathbf{v}) = r \otimes \mathbf{w}_e$ with $r \in \mathcal{O}_P$ and $\mathbf{w}_e \in \mathfrak{g}$ and denoting the unique left-invariant vector field associated to $\mathbf{w}_e$ by $\mathbf{w}$, we obtain

$$(\psi_a)_*\mathbf{v} = (\psi_a)_*(\sigma(\omega(\mathbf{v}))) = (\psi_a)_*(r \cdot \psi_*(0 \times \mathbf{w})) \stackrel{(*)}{=} ((\psi_a^{-1})^*r) \cdot (\psi_a)_*\psi_*(0 \times \mathbf{w}),$$

where $(*)$ will be proven below. Recalling the notion of the right-translation τ'_a by a on G from subsection 11.4.4, we may write $\psi_a \circ \psi = \psi \circ (\text{id} \times \tau'_a) : P \times_S G \rightarrow P$, $(p, g) \mapsto p \cdot (ga)$, and we get

$$(\psi_a)_*\mathbf{v} = ((\psi_a^{-1})^*r) \cdot \psi_*(\text{id} \times \tau'_a)_*(0 \times \mathbf{w}) = ((\psi_a^{-1})^*r) \cdot \psi_*(0 \times (\tau'_a)_*\mathbf{w})$$

As left- and right-translations on G commute with each other, $(\tau'_a)_*\mathbf{w}$ is left invariant again. Therefore $(\tau'_a)_*\mathbf{w}$ is the unique left invariant vector field on G corresponding to an element $((\tau'_a)_*\mathbf{w})_e \in \mathfrak{g}$, and, using Definition 8.16 c), this yields

$$(\psi_a)_*\mathbf{v} = \sigma((\psi_a^{-1})^*r \otimes ((\tau'_a)_*\mathbf{w})_e) = (\sigma \circ \omega \circ \sigma)((\psi_a^{-1})^*r \otimes ((\tau'_a)_*\mathbf{w})_e) = \text{ver}((\psi_a)_*\mathbf{v}).$$

In order to finish the proof of b), it remains to explain the equality $(*)$. Let $\mathbf{t}$ be a vector field on P and let $\alpha \in \Omega_{X/S}^1$. Then

$$\begin{aligned} ((\psi_a)_*(r \cdot \mathbf{t}))(\alpha) &= (\psi_a^{-1})^*((r \cdot \mathbf{t})(\psi_a^*\alpha)) = (\psi_a^{-1})^*(\mathbf{t}(r \cdot \psi_a^*\alpha)) \\ &= (\psi_a^{-1})^*(\mathbf{t}(\psi_a^*((\psi_a^{-1})^*r \cdot \alpha))) = ((\psi_a)_*\mathbf{t})((\psi_a^{-1})^*r \cdot \alpha) \quad (*) \\ &= (\psi_a^{-1})^*r \cdot ((\psi_a)_*(\mathbf{t}))(\alpha). \end{aligned}$$

Finally let us prove c). But this is an immediate consequence of b):

$$\begin{aligned} \text{hor}(\psi_a)_*\mathbf{v} &= \text{hor}(\psi_a)_*(\text{hor} \mathbf{v} + \text{ver} \mathbf{v}) = \text{hor}((\psi_a)_*\text{hor} \mathbf{v}) + \text{hor}((\psi_a)_*\text{ver} \mathbf{v}) = (\psi_a)_*(\text{hor} \mathbf{v}). \\ \text{ver}(\psi_a)_*\mathbf{v} &= \text{ver}(\psi_a)_*(\text{hor} \mathbf{v} + \text{ver} \mathbf{v}) = \text{ver}((\psi_a)_*\text{hor} \mathbf{v}) + \text{ver}((\psi_a)_*\text{ver} \mathbf{v}) = (\psi_a)_*(\text{ver} \mathbf{v}). \end{aligned}$$

□

8.5. Covariant derivation

8.19 Definition Let α be a differential k -form on P with values in $\mathfrak{g}$, i.e. a global section of $\Omega_{P/S}^k \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$. Furthermore, let d be the exterior differential (see Theorem 11.39). Then each connection form ω on P defines a covariant derivative $D\alpha$

$$(D\alpha)(\mathbf{v}_1, \dots, \mathbf{v}_{k+1}) := (d\alpha)(\text{hor } \mathbf{v}_1, \dots, \text{hor } \mathbf{v}_{k+1})$$

with vector fields $\mathbf{v}_1, \dots, \mathbf{v}_{k+1} \in \Gamma(T_{P/S}/P)$.

$D\alpha$ is a differential 2-form on P with values in $\mathfrak{g}$, because $\text{hor} \in \text{Hom}_{\mathcal{O}_P}(\Omega_{P/S}^1, \Omega_{P/S}^1)$.

8.20 Definition Let ω be a curvature form. Then we write $\Omega := D\omega$.

$D\omega$ has some physically important properties which we are going to summarize next. For this purpose we need the following lemma.

8.21 Lemma *Let $\phi : X \rightarrow X'$ be an isomorphism of S -schemes. Then the exterior differential d commutes with ϕ :*

$$\phi^* d = d\phi^*.$$

PROOF First, we prove that $[\phi_* \mathbf{v}, \phi_* \mathbf{w}] = \phi_* [\mathbf{v}, \mathbf{w}]$ (see Definition 11.41 for the notion of the commutator). This may be seen as follows:

$$(\phi_* \mathbf{v})_d(f) = (\phi_* \mathbf{v})(df) = (\phi^{-1})^*(\mathbf{v}(\phi^*(df))) = (\phi^{-1})^*(\mathbf{v}(d(\phi^* f))),$$

where the last equality is due to the fact that the exterior differential d coincides with $d_{X/S}$ on functions (Theorem 11.39), and due to the fact that the differential $d_{X/S}$ commutes (by definition) with pull-back on functions. Therefore

$$(\phi_* \mathbf{v})_d = (\phi^{-1})^* \circ \mathbf{v}_d \circ \phi^*,$$

where $(\phi^{-1})^*$ and ϕ^* both denote the pull-back of functions. Suppressing the index d , we obtain

$$\begin{aligned} [\phi_* \mathbf{v}, \phi_* \mathbf{w}](f) &= \phi_* \mathbf{v}(\phi_* \mathbf{w}(f)) - \phi_* \mathbf{w}(\phi_* \mathbf{v}(f)) \\ &= \phi_* \mathbf{v}((\phi^{-1})^*(\mathbf{w}(\phi^* f))) - \phi_* \mathbf{w}((\phi^{-1})^*(\mathbf{v}(\phi^* f))) \\ &= (\phi^{-1})^* \mathbf{v}(\phi^*((\phi^{-1})^*(\mathbf{w}(\phi^* f)))) - (\phi^{-1})^* \mathbf{w}(\phi^*((\phi^{-1})^*(\mathbf{v}(\phi^* f)))) \\ &= (\phi^{-1})^*((\mathbf{v}\mathbf{w} - \mathbf{w}\mathbf{v})(\phi^* f)) = (\phi_* [\mathbf{v}, \mathbf{w}])(f). \end{aligned}$$

in terms of derivations. Therefore $[\phi_* \mathbf{v}, \phi_* \mathbf{w}] = \phi_* [\mathbf{v}, \mathbf{w}]$ is also true for vector fields. Now

we conclude as follows:

$$\begin{aligned}
& (\phi^{-1})^* ((\phi^*(d\alpha))(\mathbf{v}_0, \dots, \mathbf{v}_k)) \\
&= (d\alpha)(\phi_* \mathbf{v}_0, \dots, \phi_* \mathbf{v}_k) \\
&= \sum_{i=0}^k (-1)^i \underbrace{(\phi_* \mathbf{v}_i)}_{=(\phi^{-1})^* \mathbf{v}_i \phi^*} (\alpha(\phi_* \mathbf{v}_0, \dots, \phi_* \check{\mathbf{v}}_i, \dots, \phi_* \mathbf{v}_k)) \\
&+ \sum_{0 \leq i < j \leq k} (-1)^{i+j} \underbrace{\alpha([\phi_* \mathbf{v}_i, \phi_* \mathbf{v}_j], \phi_* \mathbf{v}_0, \dots, \phi_* \check{\mathbf{v}}_i, \dots, \phi_* \check{\mathbf{v}}_j, \dots, \phi_* \mathbf{v}_k)}_{=\phi_* [\mathbf{v}_i, \mathbf{v}_j]} \\
&= \sum_{i=0}^k (-1)^i (\phi^{-1})^* \mathbf{v}_i ((\phi^* \alpha)(\mathbf{v}_0, \dots, \check{\mathbf{v}}_i, \dots, \mathbf{v}_k)) \\
&+ \sum_{0 \leq i < j \leq k} (-1)^{i+j} (\phi^{-1})^* (\phi^* \alpha)([\mathbf{v}_i, \mathbf{v}_j], \mathbf{v}_0, \dots, \check{\mathbf{v}}_i, \dots, \check{\mathbf{v}}_j, \dots, \mathbf{v}_k) \\
&= (\phi^{-1})^* ((d(\phi^* \alpha))(\mathbf{v}_0, \dots, \mathbf{v}_k)). \quad \square
\end{aligned}$$

8.22 Proposition $\psi_a^* \Omega = \text{Ad}(a^{-1}) \Omega$ for all S -valued points $a \in G(S)$.

PROOF The translation by a is an isomorphism. Recalling the notion of pull-back of differential forms of higher degree from Definition 7.20, and using the fact that $(\psi_a)_*$ commutes with hor (see Proposition 8.18 c)), we obtain:

$$\begin{aligned}
(\psi_a^{-1})^* ((\psi_a^* \Omega)(\mathbf{v}, \mathbf{w})) &= \Omega((\psi_a)_* \mathbf{v}, (\psi_a)_* \mathbf{w}) = (d\omega)(\text{hor}(\psi_a)_* \mathbf{v}, \text{hor}(\psi_a)_* \mathbf{w}) \\
&= (d\omega)((\psi_a)_* \text{hor} \mathbf{v}, (\psi_a)_* \text{hor} \mathbf{w})
\end{aligned}$$

Therefore

$$\begin{aligned}
(\psi_a^* \Omega)(\mathbf{v}, \mathbf{w}) &= ((\psi_a)^* d\omega)(\text{hor} \mathbf{v}, \text{hor} \mathbf{w}) \\
&= (d(\psi_a)^* \omega)(\text{hor} \mathbf{v}, \text{hor} \mathbf{w}) \quad \text{by Lemma 8.21} \\
&= \text{Ad}(a^{-1})(d\omega)(\text{hor} \mathbf{v}, \text{hor} \mathbf{w}) \\
&= \text{Ad}(a^{-1}) \Omega(\mathbf{v}, \mathbf{w}). \quad \square
\end{aligned}$$

8.23 structure-equation $\Omega(\mathbf{v}, \mathbf{w}) = d\omega(\mathbf{v}, \mathbf{w}) + [\omega(\mathbf{v}), \omega(\mathbf{w})]$, where $[\cdot, \cdot]$ denotes the Lie-bracket of $\mathcal{O}_P \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$ (see Remark 8.13).

PROOF First, let us consider the Lie-algebra valued vector field $[\omega, \omega]$ which is defined by $([\omega, \omega])(\mathbf{v}, \mathbf{w}) := [\omega(\mathbf{v}), \omega(\mathbf{w})] - [\omega(\mathbf{w}), \omega(\mathbf{v})] = 2[\omega(\mathbf{v}), \omega(\mathbf{w})]$. Then we proposition may be stated as the following equation of differential forms:

$$\Omega = d\omega + \frac{1}{2}[\omega, \omega].$$

Due to the $\mathcal{O}_P$ -multi-linearity of differential forms, it therefore suffices to consider the following three cases:

- (i) $\mathfrak{v}$ and $\mathfrak{w}$ are vertical: If we set $\omega(\mathfrak{v}) =: r \otimes \mathfrak{t}_e$, then we may write $\mathfrak{v} = \sigma(\omega(\mathfrak{v})) = \sigma(r \otimes \mathfrak{t}_e) = r \cdot \sigma(1 \otimes \mathfrak{t}_e)$. Then, again due to $\mathcal{O}_P$ -multi-linearity, it suffices to make the proof in the case $r = 1$; i.e. we may assume that $\mathfrak{v} = \sigma(1 \otimes \mathfrak{t}_e)$. For the same reason let $\mathfrak{w} = \sigma(1 \otimes \mathfrak{u}_e)$. Thus, by definition of the exterior differential d (see Corollary 11.40), we conclude:

$$(d\omega)(\mathfrak{v}, \mathfrak{w}) = \mathfrak{v}(\omega(\mathfrak{w})) - \mathfrak{w}(\omega(\mathfrak{v})) - \omega([\mathfrak{v}, \mathfrak{w}]) = \mathfrak{v}(1 \otimes \mathfrak{u}_e) - \mathfrak{w}(1 \otimes \mathfrak{t}_e) - \omega([\mathfrak{v}, \mathfrak{w}]).$$

Now, $\mathfrak{v}(1 \otimes \mathfrak{u}_e) = 0 = \mathfrak{w}(1 \otimes \mathfrak{t}_e)$, because the derivations $\mathfrak{v}$ and $\mathfrak{w}$ act on constant Lie-algebra valued functions. Thus we are done, because ω is a homomorphism of Lie-algebras.

- (ii) $\mathfrak{v}$ and $\mathfrak{w}$ are horizontal: In this special case, the statement of the proposition reduces to the definition of Ω .
- (iii) $\mathfrak{v}$ is vertical and $\mathfrak{w}$ horizontal: We get

$$(d\omega)(\mathfrak{v}, \mathfrak{w}) = \mathfrak{v}(\omega(\mathfrak{w})) - \mathfrak{w}(\omega(\mathfrak{v})) - \omega([\mathfrak{v}, \mathfrak{w}]).$$

Let us explain, why each of the summands is zero. The first one is zero, because $\omega(\mathfrak{w}) = 0$ due to horizontality of $\mathfrak{w}$. Therefore, $\omega([\mathfrak{v}, \mathfrak{w}]) = [\omega(\mathfrak{v}), \omega(\mathfrak{w})]$ vanishes, too. The second summand is zero due to the arguments already given in (i). $\square$

8.24 Bianchi-identity $D\Omega = 0$.

PROOF Notice that

$$\begin{aligned} (d[\omega, \omega])(\mathfrak{u}, \mathfrak{v}, \mathfrak{w}) &= \mathfrak{u}([\omega, \omega](\mathfrak{v}, \mathfrak{w})) - \mathfrak{v}([\omega, \omega](\mathfrak{w}, \mathfrak{u})) + \mathfrak{w}([\omega, \omega](\mathfrak{u}, \mathfrak{v})) \\ &\quad - [\omega, \omega](\mathfrak{u}, \mathfrak{v}, \mathfrak{w}) + [\omega, \omega](\mathfrak{u}, \mathfrak{w}, \mathfrak{v}) - [\omega, \omega](\mathfrak{v}, \mathfrak{w}, \mathfrak{u}) \\ &= 0 \end{aligned}$$

if $\mathfrak{u}, \mathfrak{v}$ and $\mathfrak{w}$ are horizontal vector fields. Using the structure-equation 8.23 and $d \circ d = 0$ (see Theorem 11.39), we finally obtain

$$\begin{aligned} (D\Omega)(\mathfrak{u}, \mathfrak{v}, \mathfrak{w}) &= (d\Omega)(\text{hor } \mathfrak{u}, \text{hor } \mathfrak{v}, \text{hor } \mathfrak{w}) \\ &= (dd\omega)(\text{hor } \mathfrak{u}, \text{hor } \mathfrak{v}, \text{hor } \mathfrak{w}) + (d[\omega, \omega])(\text{hor } \mathfrak{u}, \text{hor } \mathfrak{v}, \text{hor } \mathfrak{w}) = 0. \quad \square \end{aligned}$$

8.25 Definition $\Omega = D\omega$ is called the *curvature form* of the connection ω , if $\psi^*\Omega$ is horizontal in the following sense: On stalks we can write

$$\psi^*\Omega = \sum_i (\nu_i, 0) \wedge (\nu'_i, 0)$$

where $(\nu_i, 0), (\nu'_i, 0) \in \pi_1^*\Omega_{P/S}^1 \oplus \pi_2^*\Omega_{G/S}^1 \cong \Omega_{P \times_S G/S}^1$ for all i .

Remark. In the situation of Definition 8.25, the condition $\psi^*\Omega = \sum_i (\nu_i, 0) \wedge (\nu'_i, 0)$ will guarantee that the local field strengths $\mathcal{F}_i$ corresponding to Ω (see Definition 9.4), actually glue to a global field strength $\mathcal{F}$ which lives on the whole universe.

In particular, the condition $\psi^*\Omega = \sum_i (\nu_i, 0) \wedge (\nu'_i, 0)$ may be omitted, if one considers a gauge theory which is induced by a globally trivial X -torsor P under G . In this case, every connection ω gives rise to a curvature form $\Omega := D\omega$.

9. Yang-Mills equation

In this chapter we will finally establish the Yang-Mills theory over commutative rings R . Pulling-back the connection form ω and the curvature form Ω (introduced in Definition 8.16 and Definition 8.25) to space-time, we are led to the gauge potential $\mathcal{A}$ and the field strength $\mathcal{F}$. The main result of this chapter is that the gauge potential and the field strength, which are a priori only defined locally (with respect to étale topology) on space-time, actually glue to global objects of space-time. This proof will be done by means of Grothendieck's theory of faithfully flat descent. Then the Yang-Mills action and the Yang-Mills equation may be established over the base scheme $S := \operatorname{Spec} R$, generalizing the known classical theory over $R := \mathbb{R}$.

More precisely, in our physical applications, R will be a one-dimensional Dedekind ring which is also a principal ideal domain. So, we assume from now on that R is of this type. As a consequence, the sheaves of relative differential forms over $S = \operatorname{Spec} R$ will be free sheaves of modules (Theorem 11.61). In particular, the Lie-algebra $\mathfrak{g}$ of a group scheme over S will be a free $\mathcal{O}_S(S)$ -module and therefore flat. As usual, let $X \rightarrow S$ be a smooth S -scheme, let $p : G \rightarrow S$ be a smooth S -group scheme with unit section $\varepsilon : S \hookrightarrow G$, and let $G_X := G \times_S X$. Let P be a X -torsor under G_X with group action ψ . Furthermore, we assume for physical reasons that X and G are connected.

9.1. The Maurer-Cartan form

Due to the connectedness of G , the constant presheaves

$$\begin{aligned} U &\rightsquigarrow \mathcal{O}_S(S) \\ U &\rightsquigarrow \mathfrak{g} := (\varepsilon^* \Omega_{G/S}^1)(S) \end{aligned}$$

are sheaves, and due to flatness,

$$U \rightsquigarrow \mathcal{O}_G(U) \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$$

is a sheaf, too, which we denote by $\mathcal{O}_G \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$. In particular, the direct image of this sheaf under p is given by $p_* (\mathcal{O}_G \otimes_{\mathcal{O}_S(S)} \mathfrak{g}) = p_* \mathcal{O}_G \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$. We want to find a left-invariant differential form Θ which assigns to every vector field $\mathfrak{v} \in \mathcal{T}_{G/S}(G)$ the constant Lie-algebra valued function $1 \otimes \mathfrak{v}_e \in \mathcal{O}_G(G) \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$ where $\mathfrak{v}_e$ denotes the value of $\mathfrak{v}$ at the unit element e of G .¹ Let us prove that such a differential form exists, and that it

¹More precisely, $\mathfrak{v}_e$ denotes the pull-back of $\mathfrak{v}$ with respect to the unit section ε and is then considered as an element of the Lie-algebra $\mathfrak{g}$ (recall Proposition 11.32).

is uniquely determined. This is due to the following sequence of canonical isomorphism which will be explained below.

$$\begin{aligned}
\mathrm{Hom}_{\mathcal{O}_S(S)}(\mathfrak{g}, \mathcal{O}_G(G) \otimes_{\mathcal{O}_S(S)} \mathfrak{g}) &= \mathrm{Hom}_{\mathcal{O}_S}(\varepsilon^* \Omega_{G/S}^1, p_* \mathcal{O}_G \otimes_{\mathcal{O}_S(S)} \mathfrak{g}) \\
&= \mathrm{Hom}_{\mathcal{O}_S}(\varepsilon^* \Omega_{G/S}^1, p_* (\mathcal{O}_G \otimes_{\mathcal{O}_S(S)} \mathfrak{g})) \\
&= \mathrm{Hom}_{\mathcal{O}_G}(p^* \varepsilon^* \Omega_{G/S}^1, \mathcal{O}_G \otimes_{\mathcal{O}_S(S)} \mathfrak{g}) \\
&= \mathrm{Hom}_{\mathcal{O}_G}(\Omega_{G/S}^1, \mathcal{O}_G \otimes_{\mathcal{O}_S(S)} \mathfrak{g})
\end{aligned}$$

The first isomorphism is ([Liu], Ex. 5.1.5), the second one is due to the remark in the introduction above. The third states the adjointness of the functors p^* and p_* (Proposition 7.15), and the last one originates from extending elements of the Lie-algebra to left-invariant sections (Proposition 11.58). Also recalling Proposition 11.32, this bijection induces a bijection

$$\mathrm{Hom}_{\mathcal{O}_S(S)}((\varepsilon^* \mathcal{T}_{G/S})(S), \mathcal{O}_G(G) \otimes_{\mathcal{O}_S(S)} \mathfrak{g}) = \mathrm{Hom}_{\mathcal{O}_G}(\mathcal{T}_{G/S}, \mathcal{O}_G \otimes_{\mathcal{O}_S(S)} \mathfrak{g})$$

9.1 Definition The uniquely determined Lie-algebra valued differential form $\Theta \in \mathrm{Hom}_{\mathcal{O}_G}(\mathcal{T}_{G/S}, \mathcal{O}_G \otimes_{\mathcal{O}_S(S)} \mathfrak{g})$ corresponding to the $\mathcal{O}_S(S)$ -linear homomorphism

$$\begin{aligned}
(\varepsilon^* \mathcal{T}_{G/S})(S) &\longrightarrow \mathcal{O}_G(G) \otimes_{\mathcal{O}_S(S)} \mathfrak{g} \\
\mathfrak{v}_e &\mapsto 1 \otimes \mathfrak{v}_e
\end{aligned}$$

is called the *Maurer-Cartan form*. Thereby $1 \in \mathcal{O}_G(G)$ denotes the unit element.

In particular, one obtains

$$\Theta(\mathfrak{v}) = 1 \otimes \varepsilon^* \mathfrak{v}$$

for all left-invariant vector fields $\mathfrak{v} \in \Gamma(T_{G/S}/G)$.

9.2. The field strength of a gauge potential

In classical gauge theory, gauge fields are described by a gauge potential. The physical information is encoded in the associated antisymmetric field strength tensor: For example in the case of electromagnetism, the components of the field strength tensor are exactly the components of the electric and the magnetic field. Therefore the field strength has to be a global object living upon the whole space-time manifold, if the theory claims to be physically sensible.

Let us prove that the field strength is also a global object in our algebraic geometric setting of a space-time $X(S)$ consisting of “adelic” space-time points (isomorphic to the base scheme S).

Let P, X, G and S be as stated at the beginning of this chapter. Let $\hat{\theta}$ be a gauge transformation (Definition 8.11), and let $u := u_{\hat{\theta}}$ be the corresponding equivariant morphism (Proposition 8.12). Let ω be a connection form (Definition 8.16). Also recall the notion of translation by S -valued points by means of the group action from 8.10 above.

9.2.1. Gauge transformations of the curvature form

9.2 Proposition *Let $p \in P$ be a physical point. Then the stalk of the curvature form Ω at p transforms as follows under gauge transformations.*

$$(\hat{\theta}^*\Omega)_p = \text{Ad}(u(p)^{-1})\Omega_p$$

PROOF By definition of u (see proof of Proposition 8.12) we have got $\hat{\theta} = \psi \circ (\text{id}, u)$. Let $p : P \rightarrow P$ resp. $u(p) : P \rightarrow G$ be the constant maps in the sense of Proposition 7.23 mapping P to the S -valued point containing p resp. $u(p)$. Writing $\hat{p} := \hat{\theta}(p)$, we know from Proposition 7.24 that at the stalk at p

$$(\hat{\theta}^*\Omega)_p = (\text{id}, u)^*\psi^*\Omega_{\hat{p}} = (p, u)^*\psi^*\Omega_{\hat{p}} + (\text{id}, u(p))^*\psi^*\Omega_{\hat{p}}.$$

Thus we conclude

$$(\hat{\theta}^*\Omega)_p = \psi_{u(p)}^*\Omega_{\hat{p}} + (p, u)^*\psi^*\Omega_{\hat{p}}.$$

By Proposition 8.22, $\psi_{u(p)}^*\Omega_{\hat{p}} = \text{Ad}(u(p)^{-1})\Omega_p$. Thus it suffices to show that the second summand vanishes. As $(p, u) = (p \times \text{id}) \circ (\text{id}, u)$, it suffices to show that $(p \times \text{id})^*\psi^*\Omega_{\hat{p}} = 0$. But Ω is a curvature form (see Definition 8.25) and therefore we may write on stalks

$$\psi^*\Omega = \sum_i (\nu_i, 0) \wedge (\nu'_i, 0)$$

where $(\nu_i, 0), (\nu'_i, 0) \in \pi_1^*\Omega_{P/S}^1 \oplus \pi_2^*\Omega_{G/S}^1 \cong \Omega_{P \times_S G/S}^1$ for all i . Therefore

$$(p \times \text{id})^*\psi^*\Omega_{\hat{p}} = \sum_i (p \times \text{id})^*(\nu_i, 0) \wedge (p \times \text{id})^*(\nu'_i, 0) = \sum_i (p^*\nu_i, 0) \wedge (p^*\nu'_i, 0) = 0,$$

because the pull-back of differential forms under constant morphisms is zero by Proposition 7.21. $\square$

9.2.2. The curvature form in a local trivialisation

Let $\pi : P \rightarrow X$ be the canonical projection. Locally in étale topology on X , there exists a section $\bar{s} : U \hookrightarrow P \times_X U =: \pi^{-1}(U)$ of π over each point $x \in X$ (Proposition 1.19 and Proposition 1.18). We may assume that the torsor P is trivial over U , i.e. $\pi^{-1}(U) = U \times_S G =: G_U$. If $p : G \rightarrow S$ is the canonical morphism and if $\varepsilon : S \hookrightarrow G$ is the unit section, then $\bar{s}$ induces a section

$$s : U \xrightarrow{\bar{s}} \pi^{-1}(U) = U \times_S G \xrightarrow{\text{id} \times p} U \times_S S \xrightarrow{\text{id} \times \varepsilon} U \times_S G = \pi^{-1}(U).$$

This section s is called the *canonical section*.

Furthermore we will make use of the following morphisms:

- a) $\chi : \pi^{-1} \rightarrow U \times_S G$ denotes the trivialisation.
- b) $p_1 : U \times_S G \rightarrow U$ and $p_2 : U \times_S G \rightarrow G$ denote the canonical projections.
- c) Let $\kappa := p_2 \circ \chi$. As $\pi = p_1 \circ \chi$ and as χ is an isomorphism, we get in particular the cartesian diagram

$$\begin{array}{ccc} \pi^{-1}(U) & \xrightarrow{\pi} & U \\ \kappa \downarrow & & \downarrow \\ G & \longrightarrow & S \end{array}$$

- d) Let $m : G \times_S G \rightarrow G$ be the group multiplication. Recall from the definition of torsors that the trivialisation respects the group actions; i.e.

$$\kappa \circ \chi = m \circ (\kappa \times \text{id})$$

9.3 Lemma *Let s be the canonical section, and let $f := \psi \circ (s \circ \pi, \kappa) : P \rightarrow P$. Then $f = \text{id}$.*

PROOF There is a commutative diagram

$$\begin{array}{ccc} P \times_X G & \xrightarrow{\psi} & P \\ \pi_1 \downarrow & & \downarrow \pi \\ P & \xrightarrow{\pi} & X \end{array}$$

Therefore we conclude

$$\pi \circ f = \pi \circ \psi \circ (s \circ \pi, \kappa) = \pi \circ \pi_1 \circ (s \circ \pi, \kappa) = \pi \circ s \circ \pi = \pi.$$

Furthermore,

$$\begin{aligned} \kappa \circ f &= m \circ (\kappa \times \text{id}) \circ (s \circ \pi, \kappa) = m(\kappa \circ s \circ \pi, \kappa) \\ &= m(p_2 \circ \chi \circ \chi^{-1} \circ (\text{id} \times \varepsilon) \circ (\text{id} \times p) \circ \chi \circ \bar{s} \circ \pi, \kappa) \\ &= m(\varepsilon \circ p \circ p_2 \circ \chi \circ \bar{s} \circ \pi, \kappa) = m \circ ((\varepsilon \circ p) \times \text{id}) \circ (\kappa \circ \bar{s} \circ \pi, \kappa) \\ &= p_2(\kappa \circ \bar{s} \circ \pi, \kappa) = \kappa. \end{aligned}$$

Thus (by c)) we know that f is the uniquely determined morphism making the diagram

$$\begin{array}{ccc} & & U \\ & \nearrow \pi & \uparrow \pi \\ \pi^{-1}(U) & \xrightarrow{f} & \pi^{-1}(U) \\ & \searrow \kappa & \downarrow \kappa \\ & & G \end{array}$$

commutative; i.e. $f = \text{id}$. □

9.4 Definition Let $s : U \hookrightarrow \pi^{-1}(U)$ be the canonical section, let ω be a connection form, and let Ω be a curvature form. Then

$$\begin{aligned}\mathcal{A} &:= s^*\omega && \text{is called the local } \textit{gauge potential}, \text{ and} \\ \mathcal{F} &:= s^*\Omega && \text{is called the local } \textit{field strength}.\end{aligned}$$

These are differential forms on U with values in the Lie-algebra $\mathfrak{g}$.

9.5 Proposition *Let $p \in P$ be a physical point. Then on stalks at p we get the identity*

$$\Omega_p = \text{Ad}(\kappa(p)^{-1})(\pi^*\mathcal{F})_p$$

PROOF Let $(s \circ \pi)(p) : P \rightarrow P$ resp. $\kappa(p) : P \rightarrow G$ be the constant maps in the sense of Proposition 7.23 mapping P to the S -valued point containing $(s \circ \pi)(p)$ resp. $\kappa(p)$. By Lemma 9.3, we know that $\psi \circ (s \circ \pi, \kappa) = \text{id}$. Due to Proposition 7.24 we therefore get at the stalk at p

$$\Omega_p = (s \circ \pi, \kappa)^*\psi^*\Omega_p = ((s \circ \pi)(p), \kappa)^*\psi^*\Omega_p + (s \circ \pi, \kappa(p))^*\psi^*\Omega_p.$$

Using the fact that $(s \circ \pi, \kappa(p)) = (\text{id}, \kappa(p)) \circ s \circ \pi$, we conclude

$$\Omega_p = \pi^*s^*\psi_{\kappa(p)}^*\Omega_p + ((s \circ \pi)(p), \kappa)^*\psi^*\Omega_p.$$

Exactly the same argument as in the proof of Proposition 9.2 shows that the second summand $((s \circ \pi)(p), \kappa)^*\psi^*\Omega_p$ vanishes. For the first summand we obtain with Proposition 8.22

$$\pi^*s^*\psi_{\kappa(p)}^*\Omega_p = \pi^*s^*\text{Ad}(\kappa(p)^{-1})\Omega_{p \cdot \kappa(p)^{-1}} = \text{Ad}(\kappa(p)^{-1})\pi^*(s^*\Omega)_{\pi(p)} = \text{Ad}(\kappa(p)^{-1})(\pi^*\mathcal{F})_p,$$

and we are done. $\square$

Let us now choose an étale covering $(U_i \rightarrow X)_i$ which trivializes P . For each étale open subset $U_i \rightarrow X$ let χ_i be the respective trivialisation, and let $s_i : U_i \hookrightarrow \pi^{-1}(U_i) := P \times_X U_i$ be the canonical section of $\kappa_i := p_2 \circ \chi_i : \pi^{-1}(U_i) \rightarrow U_i$. Let $\pi_i : \pi^{-1}(U_i) \rightarrow U_i$ be the canonical projection.

Then we get local gauge potentials and local field strengths

$$\begin{aligned}\mathcal{A}_i &:= s_i^*\omega, \\ \mathcal{F}_i &:= s_i^*\Omega\end{aligned}$$

and by Proposition 9.5 we have got

$$\Omega_p = \text{Ad}(\kappa_i(p)^{-1})(\pi_i^*\mathcal{F}_i)_p \quad \text{for all } i.$$

Pulling back this identity with the canonical projection $U_{ij} := U_i \times_X U_j \rightarrow U_i$ (which is an étale morphism), and using the fact that the notion of physical points is stable under étale base change (Proposition 7.10), it follows that

$$\text{Ad}(\kappa_j(p)^{-1})(\pi_j^*\mathcal{F}_j)_p = \Omega_p = \text{Ad}(\kappa_i(p)^{-1})(\pi_i^*\mathcal{F}_i)_p$$

for all physical points $p \in \pi^{-1}(U_{ij}) := P \times_X U_{ij}$, if we omit restriction signs. As π_i and π_j are obtained from the global morphism π by base change with $U_i \rightarrow X$ resp. $U_j \rightarrow X$, π_i and π_j coincide on U_{ij} , and their restrictions to U_{ij} may both be denoted by π_{ij} . Choosing a section $\hat{s}$ of π_{ij} (which exists, as there are already sections over U_i or U_j), and applying this section to the equation above, we obtain for all physical points $p \in \pi^{-1}(U_{ij})$ over $x \in U_{ij}$

$$\mathrm{Ad}(\kappa_j(p)^{-1})(\mathcal{F}_j)_x = \mathrm{Ad}(\kappa_i(p)^{-1})(\mathcal{F}_i)_x.$$

Recalling from Proposition 11.49 that Ad is compatible with the group law on G , i.e. $\mathrm{Ad}(\kappa_i(p)) \circ \mathrm{Ad}(\kappa_j(p)^{-1}) = \mathrm{Ad}(\kappa_i(p) \cdot \kappa_j(p)^{-1})$, we get

9.6 Proposition *Let $x \in U_{ij}$ be a physical point. Then (omitting restriction signs), one has got*

$$(\mathcal{F}_j)_x = \mathrm{Ad}(\rho_{ij}^{-1}(x))(\mathcal{F}_i)_x,$$

where the family of all $\rho_{ij} : U_{ij} \rightarrow G$ is a 1-cocycle in the sense of Proposition 8.9.

PROOF It remains to prove the statement on the morphism ρ_{ij} . For this purpose it suffices to show that there exists a morphism $\rho_{ij} : U_{ij} \rightarrow G$ such that

$$\rho_{ij} \circ \pi = m \circ (\kappa_i, \iota \circ \kappa_j)$$

where $\pi : \pi^{-1}(U_{ij}) = U_{ij} \times_S G \rightarrow U_{ij}$ is the canonical projection, where m is the group law on G , and where ι is the formation of the inverse in G . In order to define ρ_{ij} choose an arbitrary section of $G \rightarrow S$. By base change, this induces a section ϵ of $\pi : \pi^{-1}(U_{ij}) = U_{ij} \times_S G \rightarrow U_{ij}$. Then we set

$$\rho_{ij} := m \circ (\kappa_i, \iota \circ \kappa_j) \circ \epsilon.$$

This is a 1-cocycle and it remains to prove that $\rho_{ij} \circ \pi = m \circ (\kappa_i, \iota \circ \kappa_j)$. By the Yoneda lemma (Proposition 11.43), we may check this on T -valued points, where $T \rightarrow S$ is an arbitrary S -scheme. It suffices to show that $(m \circ (\kappa_i, \iota \circ \kappa_j))(p \cdot g) = (m \circ (\kappa_i, \iota \circ \kappa_j))(p)$ for all T -valued points p of P and g of G . Using the fact that the trivialization isomorphisms of torsors respect the group action, the following computation indeed shows that

$$\begin{aligned} (m \circ (\kappa_i, \iota \circ \kappa_j))(p \cdot g) &= (\kappa_i)(p \cdot g) \cdot ((\kappa_j)(p \cdot g))^{-1} \\ &= (\kappa_i)(p) \cdot g \cdot g^{-1} \cdot (\kappa_j)(p)^{-1} = (\kappa_i)(p) \cdot (\kappa_j)(p)^{-1} \\ &= (m \circ (\kappa_i, \iota \circ \kappa_j))(p). \end{aligned} \quad \square$$

9.2.3. The global field strength

Let us finally prove that the collection of local field strengths $\mathcal{F}_i$ (considered in the previous subsection 9.2.2) glue to a global field strength $\mathcal{F}$ living on all of $X(S)$.

The starting point is the family of étale-local sections $(\mathcal{F}_i)_i$, where $\mathcal{F}_i = s_i^* \Omega \in \Omega_{U_i/S}^2(U_i) \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$. We already know from Proposition 9.6 that

$$(\mathcal{F}_j)_x = \text{Ad}(\rho_{ij}^{-1}(x))(\mathcal{F}_i)_x$$

upon U_{ij} for all physical points $x \in U_{ij}$. Using the identification of sections of a locally free sheaf and sections of an associated vector bundle, we may interpret $\mathcal{F}_i$ as a morphism $f_i : U_i \rightarrow \mathbb{V}(\mathcal{T}_{U_i/S}^{\otimes 2} \otimes_{\mathcal{O}_S(S)} \mathfrak{g})$. Therefore $\text{Ad}(\rho_{ij}^{-1}(x))$ induces an automorphism of $\mathbb{V}(\mathcal{T}_{U_i/S}^{\otimes 2} \otimes_{\mathcal{O}_S(S)} \mathfrak{g})$, because $\text{Ad} \circ \rho_{ij}^{-1} : U_{ij} \rightarrow \text{Aut}_{\mathcal{O}_S\text{-lin}}(\text{Lie}(G/S))$ by the expositions in subsection 11.4.3. Due to the above equation, the restriction of the morphisms f_j and $(\text{Ad} \circ \rho_{ij}^{-1})f_i$ to U_{ij} coincide on physical points. Let us denote by X' the X -scheme given by the disjoint union

$$\coprod_{\text{physical points } x \in X} V(\mathcal{J}_x)$$

where $\mathcal{J}_x$ is the sheaf of ideal corresponding to the closed point x . The étale covering $(U_i \rightarrow X)$ induces an étale covering $(U'_i \rightarrow X')$ of X' . By construction, the morphisms f_j and $(\text{Ad} \circ \rho_{ij}^{-1})f_i$ coincide on U'_{ij} . As ρ_{ij} is a 1-cocycle, the morphisms $(f'_i := f_i|_{U'_i})$ are morphisms of schemes with descent datum. Therefore, by Theorem 11.69, the family of morphism (f'_i) descends to a morphism $f : X' \rightarrow \mathbb{V}(\iota^* \mathcal{T}_{X/S}^{\otimes 2} \otimes_{\mathcal{O}_S(S)} \mathfrak{g})$, where $\iota : X' \hookrightarrow X$ is the canonical injection. f corresponds to a global section of a locally free sheaf $\mathcal{F}$ over X' and is already the searched global field strength, because set theoretically $X' = X(S)$.

9.7 Remark As the schemes X, G and P are Néron (lft)-models, the above calculations may be performed as well directly on S -valued point instead of physical points. All results of section 9.2 remain true if the word “physical point” is replaced by “ S -valued point”.

This way it is possible to construct the global field strength on $X(S)$, where $X(S)$ is this time endowed with the following structure of a X -scheme:

$$X(S) = \coprod_{S\text{-valued points } \alpha \in X(S)} V(\mathcal{J}_\alpha).$$

Thereby, $\mathcal{J}_\alpha$ is the sheaf of ideals realizing the image of α as a closed subscheme of X .

9.3. Yang-Mills equation

Within this section, let us assume that the torsor $\pi : P \rightarrow X$ is trivial. Then the field strength $\mathcal{F}$ exists as a global section of $\Omega_{X/S}^2 \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$; i.e. $\mathcal{F} \in \Omega_{X/S}^2(X) \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$ due to our assumption on S . Let s be the canonical section of $\pi : P \rightarrow X$ (introduced in the

beginning of subsection 9.2.2). Then we may pull back the covariant derivation D (see Definition 8.19) with respect to s to X :

$$D_X(s^*\alpha) := s^*(D\alpha)$$

for all $\alpha \in \Omega_{P/S}^k(P) \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$. Let us remark that (due to Proposition 1.7) there is a decomposition $s^*\Omega_{P/S}^1 \cong s^*p_1^*\Omega_{X/S}^1 \oplus s^*p_2^*\Omega_{G/S}^1 \cong \Omega_{X/S}^1 \oplus p_X^* \left(\varepsilon^*\Omega_{G/S}^1 \right)$, where $\varepsilon : S \hookrightarrow G$ is the unit section and where $p_X : X \rightarrow S$ is the canonical morphism. In particular $\Omega_{X/S}^1(X) \hookrightarrow (s^*\Omega_{P/S}^1)(X)$, i.e. each differential form $\nu \in \Omega_{X/S}^1(X)$ may be written as pull-back $s^*\alpha$ in our setting. Thus it makes sense to define

9.8 Yang-Mills equation $*D_X*\mathcal{F} = 0$,

where $*$ is the Hodge-star operator (Definition 11.42). Furthermore we deduce from the Bianchi-identity 8.24 the following proposition.

9.9 Proposition $D_X\mathcal{F} = 0$.

PROOF $D_X\mathcal{F} = D_X(s^*\Omega) = s^*(D\Omega) = 0$. □

9.10 Proposition $\mathcal{F}_x(\mathfrak{v}, \mathfrak{w}) = (d\mathcal{A})_x(\mathfrak{v}, \mathfrak{w}) + [\mathcal{A}_x(\mathfrak{v}), \mathcal{A}_x(\mathfrak{w})]$ for all $x \in X$.

PROOF Recalling the definition of the stalkwise push-forward of vector fields by means of closed immersions from Remark 7.18, we know that the identity $s_*\mathfrak{v} = \pi^* \circ \mathfrak{v} \circ s^*$ holds on stalks. Let us first prove that stalkwise s^* commutes with the exterior differential (recall that this is true globally in the case of diffeomorphisms). One finds on stalks

$$\begin{aligned} [s_*\mathfrak{v}, s_*\mathfrak{w}] &= \pi^* \circ \mathfrak{v} \circ s^* \circ \pi^* \circ \mathfrak{w} \circ s^* - \pi^* \circ \mathfrak{w} \circ s^* \circ \pi^* \circ \mathfrak{v} \circ s^* \\ &= \pi^* \circ \mathfrak{v} \circ \mathfrak{w} \circ s^* - \pi^* \circ \mathfrak{w} \circ \mathfrak{v} \circ s^* \\ &= s_*[\mathfrak{v}, \mathfrak{w}]. \end{aligned}$$

Now we conclude as follows:

$$\begin{aligned} &\pi^*((s^*(d\alpha))(\mathfrak{v}_0, \dots, \mathfrak{v}_k)) \\ &= (d\alpha)(s_*\mathfrak{v}_0, \dots, s_*\mathfrak{v}_k) \\ &= \sum_{i=0}^k (-1)^i \underbrace{(s_*\mathfrak{v}_i)}_{=\pi^*\mathfrak{v}_i s^*} (\alpha(s_*\mathfrak{v}_0, \dots, s_*\check{\mathfrak{v}}_i, \dots, s_*\mathfrak{v}_k)) \\ &+ \sum_{0 \leq i < j \leq k} (-1)^{i+j} \alpha(\underbrace{[s_*\mathfrak{v}_i, s_*\mathfrak{v}_j]}_{=s_*[\mathfrak{v}_i, \mathfrak{v}_j]}, s_*\mathfrak{v}_0, \dots, s_*\check{\mathfrak{v}}_i, \dots, s_*\check{\mathfrak{v}}_j, \dots, s_*\mathfrak{v}_k) \\ &= \sum_{i=0}^k (-1)^i \pi^*\mathfrak{v}_i ((s^*\alpha)(\mathfrak{v}_0, \dots, \check{\mathfrak{v}}_i, \dots, \mathfrak{v}_k)) \\ &+ \sum_{0 \leq i < j \leq k} (-1)^{i+j} \pi^*((s^*\alpha)([\mathfrak{v}_i, \mathfrak{v}_j], \mathfrak{v}_0, \dots, \check{\mathfrak{v}}_i, \dots, \check{\mathfrak{v}}_j, \dots, \mathfrak{v}_k)) \\ &= \pi^*((d(s^*\alpha))(\mathfrak{v}_0, \dots, \mathfrak{v}_k)). \end{aligned}$$

As the ring-homomorphism π^* is injective, we obtain the desired commutativity of s^* with d . Also recalling the structure-equation 8.23, it follows

$$\begin{aligned}
 \pi^*(\mathcal{F}(\mathfrak{v}, \mathfrak{w})) &= \pi^*((s^*\Omega)(\mathfrak{v}, \mathfrak{w})) = \Omega(s_*\mathfrak{v}, s_*\mathfrak{w}) = d\omega(s_*\mathfrak{v}, s_*\mathfrak{w}) + [\omega(s_*\mathfrak{v}), \omega(s_*\mathfrak{w})] \\
 &= \pi^*\left(\underbrace{(s^*d\omega)(\mathfrak{v}, \mathfrak{w})}_{=ds^*\omega}\right) + [\pi^*((s^*\omega)(\mathfrak{v})), \pi^*((s^*\omega)(\mathfrak{w}))] \\
 &= \pi^*((d\mathcal{A})(\mathfrak{v}, \mathfrak{w})) + [\pi^*(\mathcal{A}(\mathfrak{v})), \pi^*(\mathcal{A}(\mathfrak{w}))] \\
 &= \pi^*((d\mathcal{A})(\mathfrak{v}, \mathfrak{w})) + \pi^*([\mathcal{A}(\mathfrak{v}), \mathcal{A}(\mathfrak{w})]).
 \end{aligned}$$

Once again making use of the injectivity of the ring-homomorphism π^* , we are done. $\square$

10. Yang-Mills theory in local coordinates

In the following, let us use the same notations as in chapter 9. In particular, we consider a datum of an X -torsor P under $G_X := G \times_S X$ underlying the universe. In order to determine the field strength $\mathcal{F} \in \Omega_{X/S}^2(X) \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$ of a gauge field, it suffices to determine all stalks $\mathcal{F}_x$ for all $x \in X$. Thus we may assume that the torsor P is trivial.

Within this chapter, we will express the global equations governing the gauge field of section 9.3 on stalks. Thus we obtain the Yang-Mills theory in local coordinates.

For clarity, let us fix some notations: Let the smooth S -scheme $X \rightarrow S$ underlying space-time be of relative dimension n , and let us denote the relative dimension of the gauge group $G \rightarrow S$ by N . Then let

$$\begin{aligned} \{dx^\mu\}_{\mu=1}^n & \text{ be a base of } \Omega_{X/S}^1(X), \\ \{\frac{\partial}{\partial x^\mu}\}_{\mu=1}^n & \text{ be a base of } \mathcal{T}_{X/S}(X), \text{ which is dual to } \{dx^\mu\}_{\mu=1}^n, \\ \{b_i\}_{i=1}^N & \text{ be a base of } \mathfrak{g}. \end{aligned}$$

Then we may write the stalk of the gauge potential and the field strength at $x \in X$ in the form

$$\begin{aligned} \mathcal{A}_x &= \sum_{\mu=1}^n \sum_{i=1}^N \mathcal{A}_{\mu,x}^i dx^\mu \otimes b_i \\ \mathcal{F}_x &= \sum_{\mu,\nu=1}^n \sum_{i=1}^N \frac{1}{2} \mathcal{F}_{\mu\nu,x}^i dx^\mu \wedge dx^\nu \otimes b_i \end{aligned}$$

with $\mathcal{A}_{\mu,x}^i, \mathcal{F}_{\mu\nu,x}^i \in \mathcal{O}_{X,x}$. Recall that we embed $\mathcal{O}_{X,x}$ into a ring of formal power series if x is a physical point (see Proposition 1.36). Therefore, $\frac{\partial}{\partial x^\mu} \mathcal{A}_{\mu,x} = \frac{\partial \mathcal{A}_{\mu,x}}{\partial x^\mu}$ for all physical points $x = (x_1, \dots, x_n)$ of X , where $\frac{\partial \mathcal{A}_{\mu,x}}{\partial x^\mu}$ denotes the ordinary partial derivative of the power series $\mathcal{A}_{\mu,x}$ with respect to the variable x^μ . The analogous statement is true for the components $\mathcal{F}_{\mu\nu,x}^i$ of the field strength. In order to simplify the notation let us write $\mathcal{A}_\mu^i$ (resp. $\mathcal{F}_{\mu\nu}^i$) instead of $\mathcal{A}_{\mu,x}^i$ (resp. $\mathcal{F}_{\mu\nu,x}^i$) whenever no confusion is possible.

Furthermore let us introduce the structure coefficients c_{ij}^k for the chosen base of the Lie-algebra $\mathfrak{g}$ which are defined as follows: $[b_i, b_j] =: \sum_{k=1}^N c_{ij}^k b_k$.

10.1. Relations between gauge potential and field strength

10.1 Proposition *Let $x \in X$ be a physical point. Then the following equalities hold at the stalk at x .*

$$\begin{aligned} a) \quad \mathcal{F}_{\nu_1 \nu_2}^k &= \frac{\partial \mathcal{A}_{\nu_2}^k}{\partial x^{\nu_1}} - \frac{\partial \mathcal{A}_{\nu_1}^k}{\partial x^{\nu_2}} + \sum_{i,j=1}^N c_{ij}^k \mathcal{A}_{\nu_1}^i \mathcal{A}_{\nu_2}^j = -\mathcal{F}_{\nu_2 \nu_1}^k . \\ b) \quad \frac{\partial \mathcal{F}_{\mu \nu}}{\partial x^\rho} + \frac{\partial \mathcal{F}_{\rho \mu}}{\partial x^\nu} + \frac{\partial \mathcal{F}_{\nu \rho}}{\partial x^\mu} + [\mathcal{A}_\rho, \mathcal{F}_{\mu \nu}] + [\mathcal{A}_\nu, \mathcal{F}_{\rho \mu}] + [\mathcal{A}_\mu, \mathcal{F}_{\nu \rho}] &= 0 . \end{aligned}$$

PROOF Item a) may be derived as follows:

$$\begin{aligned} & \sum_{i=1}^N \frac{1}{2} (\mathcal{F}_{\nu_1 \nu_2}^i - \mathcal{F}_{\nu_2 \nu_1}^i) b_i \\ &= \sum_{\mu_1, \mu_2=1}^n \sum_{i=1}^N \frac{1}{2} \mathcal{F}_{\mu_1 \mu_2}^i \left\{ dx^{\mu_1} \left(\frac{\partial}{\partial x^{\nu_1}} \right) dx^{\mu_2} \left(\frac{\partial}{\partial x^{\nu_2}} \right) - dx^{\mu_1} \left(\frac{\partial}{\partial x^{\nu_2}} \right) dx^{\mu_2} \left(\frac{\partial}{\partial x^{\nu_1}} \right) \right\} b_i \\ &= \sum_{\mu_1, \mu_2=1}^n \sum_{i=1}^N \frac{1}{2} \mathcal{F}_{\mu_1 \mu_2}^i (dx^{\mu_1} \wedge dx^{\mu_2}) \left(\frac{\partial}{\partial x^{\nu_1}}, \frac{\partial}{\partial x^{\nu_2}} \right) b_i \\ &= \mathcal{F}_x \left(\frac{\partial}{\partial x^{\nu_1}}, \frac{\partial}{\partial x^{\nu_2}} \right) \\ &= (d\mathcal{A})_x \left(\frac{\partial}{\partial x^{\nu_1}}, \frac{\partial}{\partial x^{\nu_2}} \right) + \left[\mathcal{A}_x \left(\frac{\partial}{\partial x^{\nu_1}} \right), \mathcal{A}_x \left(\frac{\partial}{\partial x^{\nu_2}} \right) \right] \quad \text{by Proposition 9.10} \\ &= \frac{\partial}{\partial x^{\nu_1}} \mathcal{A}_x \left(\frac{\partial}{\partial x^{\nu_2}} \right) - \frac{\partial}{\partial x^{\nu_2}} \mathcal{A}_x \left(\frac{\partial}{\partial x^{\nu_1}} \right) - \mathcal{A}_x \left(\left[\frac{\partial}{\partial x^{\nu_1}}, \frac{\partial}{\partial x^{\nu_2}} \right] \right) + \left[\mathcal{A}_x \left(\frac{\partial}{\partial x^{\nu_1}} \right), \mathcal{A}_x \left(\frac{\partial}{\partial x^{\nu_2}} \right) \right] \\ &= \sum_{i=1}^N \frac{\partial \mathcal{A}_{\nu_2}^i}{\partial x^{\nu_1}} b_i - \frac{\partial \mathcal{A}_{\nu_1}^i}{\partial x^{\nu_2}} b_i + \sum_{i,j=1}^N [\mathcal{A}_{\nu_1}^i b_i, \mathcal{A}_{\nu_2}^j b_j] . \end{aligned}$$

This is already the desired equation if we make use of the structure coefficients c_{ij}^k . Item b) may be seen as follows. By Proposition 10.4 we get the following equation on stalks:

$$\begin{aligned} 0 &= (D_X \mathcal{F})(\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3) \\ &= d\mathcal{F}(\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3) \\ &\quad + [\mathcal{A}(\mathbf{v}_1), \mathcal{F}(\mathbf{v}_2, \mathbf{v}_3)] + [\mathcal{A}(\mathbf{v}_2), \mathcal{F}(\mathbf{v}_3, \mathbf{v}_1)] + [\mathcal{A}(\mathbf{v}_3), \mathcal{F}(\mathbf{v}_1, \mathbf{v}_2)] . \end{aligned}$$

Choosing $\mathbf{v}_\mu = \frac{\partial}{\partial x^\mu}$ we are done. $\square$

In section 9.3 we introduced a canonical notion of covariant derivation on X by pulling back a covariant derivation on the X -torsor P under G_X by means of the canonical section s . Alternatively we could have used the following less transparent but more explicit definition of covariant derivation on X .

10.2 Definition Let Y be a smooth S -scheme. Let $\eta \in \Omega_{Y/S}^1(Y) \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$ be a $\mathfrak{g}$ -valued differential 1-form, and let $\vartheta \in \Omega_{Y/S}^k(Y) \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$ be a $\mathfrak{g}$ -valued differential k -form. Then we denote by $[\eta, \vartheta]$ the following $\mathfrak{g}$ -valued differential $(k+1)$ -form:

$$[\eta, \vartheta](\mathbf{v}_1, \dots, \mathbf{v}_{k+1}) := \sum_{i=1}^{k+1} (-1)^{i+1} [\eta(\mathbf{v}_i), \vartheta(\mathbf{v}_1, \dots, \check{\mathbf{v}}_i, \dots, \mathbf{v}_{k+1})].$$

10.3 Definition Let α be a differential k -form on X with values in $\mathfrak{g}$. Then we define the covariant derivation $D_X \alpha$ of α at $x \in X$ in the following way:

$$D_X \alpha := d\alpha + [\mathcal{A}, \alpha].$$

Let us use this notion of covariant derivation on X in this section about Yang-Mills theory in local coordinates. Then Yang-Mills equation may be written as well with respect to the covariant derivation D_X of Definition 10.3, and furthermore one proves:

10.4 Proposition *Let D_X be as in Definition 10.3. Then $D_X \mathcal{F} = 0$.*

PROOF It suffices to show the statement of the proposition on stalks, i.e.: $(D_X \mathcal{F})_x = 0$ for all $x \in X$. Therefore let us perform the following computations in the stalk at x , but let us suppress the index x in order to simplify the notation.

Let ω be a connection form, and let Ω be the corresponding curvature form such that $\mathcal{A} = s^* \omega$ and $\mathcal{F} = s^* \Omega$. Then

$$D_X \mathcal{F} = ds^* \Omega + [s^* \omega, s^* \Omega].$$

By Lemma 10.5, we know that $[s^* \omega, s^* \Omega] = s^* [\omega, \Omega]$, and furthermore s^* commutes with the exterior differential d (the latter was shown in the proof of Proposition 9.10). It follows that

$$D_X \mathcal{F} = s^* (d\Omega + [\omega, \Omega]).$$

Then the statement of the proposition follows from Proposition 10.6. $\square$

10.5 Lemma *Let $s : X \hookrightarrow P$ be a section of the smooth and separated X -torsor $\pi : P \rightarrow X$ under $G_X := G \times_S X$. Let $\eta \in \Omega_{P/S}^1(P) \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$ be a $\mathfrak{g}$ -valued differential 1-form, and let $\vartheta \in \Omega_{P/S}^k(P) \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$ be a $\mathfrak{g}$ -valued differential k -form. Consider the $\mathfrak{g}$ -valued differential $(k+1)$ -form*

$$[\eta, \vartheta](\mathbf{v}_1, \dots, \mathbf{v}_{k+1}) := \sum_{i=1}^{k+1} (-1)^{i+1} [\eta(\mathbf{v}_i), \vartheta(\mathbf{v}_1, \dots, \check{\mathbf{v}}_i, \dots, \mathbf{v}_{k+1})].$$

Then on stalks the following identity holds:

$$[s^* \eta, s^* \vartheta] = s^* [\eta, \vartheta].$$

PROOF

$$\begin{aligned}
\pi^* \left((s^*[\eta, \vartheta])(\mathbf{v}_1, \dots, \mathbf{v}_{k+1}) \right) &= [\eta, \vartheta](s_*\mathbf{v}_1, \dots, s_*\mathbf{v}_{k+1}) \\
&= \sum_{i=1}^{k+1} (-1)^{i+1} [\eta(s_*\mathbf{v}_i), \vartheta(s_*\mathbf{v}_1, \dots, s_*\check{\mathbf{v}}_i, \dots, s_*\mathbf{v}_{k+1})] \\
&= \sum_{i=1}^{k+1} (-1)^{i+1} [\pi^* \left(\underbrace{(s^*\eta)(\mathbf{v}_i)}_{=: \sum_j r_{ij} \otimes g_{ij}} \right), \pi^* \left(\underbrace{(s^*\vartheta)(\mathbf{v}_1, \dots, \check{\mathbf{v}}_i, \dots, \mathbf{v}_{k+1})}_{=: \sum_l r'_{il} \otimes g'_{il}} \right)] \\
&= \sum_{i=1}^{k+1} (-1)^{i+1} \sum_{j,l} [(\pi^* \otimes \text{id})(r_{ij} \otimes g_{ij}), (\pi^* \otimes \text{id})(r'_{il} \otimes g'_{il})] \\
&= \sum_{i=1}^{k+1} (-1)^{i+1} \sum_{j,l} [(\pi^* r_{ij} \otimes g_{ij}), (\pi^* r'_{il} \otimes g'_{il})] = \sum_{i=1}^{k+1} (-1)^{i+1} \sum_{j,l} \left(\underbrace{\pi^* r_{ij} \cdot \pi^* r'_{il}}_{=: \pi^*(r_{ij} \cdot r'_{il})} \otimes [g_{ij}, g'_{il}] \right) \\
&\quad (\text{where we used the Lie-algebra structure on } \mathcal{O}_P \otimes \mathfrak{g} \text{ introduced in Remark 8.13}) \\
&= \sum_{i=1}^{k+1} (-1)^{i+1} \sum_{j,l} (\pi^* \otimes \text{id})(r_{ij} \cdot r'_{il} \otimes [g_{ij}, g'_{il}]) \\
&= \sum_{i=1}^{k+1} (-1)^{i+1} (\pi^* \otimes \text{id}) \left(\left[\sum_j r_{ij} \otimes g_{ij}, \sum_l r'_{il} \otimes g'_{il} \right] \right) \\
&= \sum_{i=1}^{k+1} (-1)^{i+1} \pi^* \left([(s^*\eta)(\mathbf{v}_i), (s^*\vartheta)(\mathbf{v}_1, \dots, \check{\mathbf{v}}_i, \dots, \mathbf{v}_{k+1})] \right) \\
&= \pi^* \left([(s^*\eta), (s^*\vartheta)](\mathbf{v}_1, \dots, \mathbf{v}_{k+1}) \right)
\end{aligned}$$

As the ring-homomorphism π^* is injective, we are done. $\square$

10.6 Proposition $d\Omega + [\omega, \Omega] = 0$.

PROOF The structure-equation 8.23 states that $\Omega = d\omega + \frac{1}{2}[\omega, \omega]$. Therefore

$$d\Omega + [\omega, \Omega] = dd\omega + \frac{1}{2}d[\omega, \omega] + [\omega, d\omega] + \frac{1}{2}[\omega, [\omega, \omega]].$$

The first summand is zero, because $d \circ d = 0$. Let us show that the second and the third summand add to zero. Due to the definition of the exterior differential (Corollary 11.40), the second summand reads as follows:

$$\begin{aligned}
\left(\frac{1}{2}d[\omega, \omega] \right) (\mathbf{u}, \mathbf{v}, \mathbf{w}) &= \frac{1}{2}\mathbf{u}([\omega, \omega](\mathbf{v}, \mathbf{w})) - \frac{1}{2}\mathbf{v}([\omega, \omega](\mathbf{w}, \mathbf{u})) + \frac{1}{2}\mathbf{w}([\omega, \omega](\mathbf{u}, \mathbf{v})) \\
&\quad - \frac{1}{2}[\omega, \omega](\mathbf{u}, \mathbf{v}, \mathbf{w}) + \frac{1}{2}[\omega, \omega](\mathbf{u}, \mathbf{w}, \mathbf{v}) - \frac{1}{2}[\omega, \omega](\mathbf{v}, \mathbf{w}, \mathbf{u}) \\
&= \mathbf{u}([\omega(\mathbf{v}), \omega(\mathbf{w})]) - \mathbf{v}([\omega(\mathbf{w}), \omega(\mathbf{u})]) + \mathbf{w}([\omega(\mathbf{u}), \omega(\mathbf{v})]) \\
&\quad - [\omega(\mathbf{u}, \mathbf{v}), \omega(\mathbf{w})] + [\omega(\mathbf{u}, \mathbf{w}), \omega(\mathbf{v})] - [\omega(\mathbf{v}, \mathbf{w}), \omega(\mathbf{u})]
\end{aligned}$$

For the third summand we obtain

$$\begin{aligned}
[\omega, d\omega](\mathbf{u}, \mathbf{v}, \mathbf{w}) &= [\omega(\mathbf{u}), (d\omega)(\mathbf{v}, \mathbf{w})] - [\omega(\mathbf{v}), (d\omega)(\mathbf{u}, \mathbf{w})] + [\omega(\mathbf{w}), (d\omega)(\mathbf{u}, \mathbf{v})] \\
&= [\omega(\mathbf{u}), (d\omega)(\mathbf{v}, \mathbf{w})] + [\omega(\mathbf{v}), (d\omega)(\mathbf{w}, \mathbf{u})] + [\omega(\mathbf{w}), (d\omega)(\mathbf{u}, \mathbf{v})] \\
&= [\omega(\mathbf{u}), \mathbf{v}(\omega(\mathbf{w})) - \mathbf{w}(\omega(\mathbf{v})) - \omega([\mathbf{v}, \mathbf{w}])] \\
&\quad + [\omega(\mathbf{v}), \mathbf{w}(\omega(\mathbf{u})) - \mathbf{u}(\omega(\mathbf{w})) - \omega([\mathbf{w}, \mathbf{u}])] \\
&\quad + [\omega(\mathbf{w}), \mathbf{u}(\omega(\mathbf{v})) - \mathbf{v}(\omega(\mathbf{u})) - \omega([\mathbf{u}, \mathbf{v}])] \\
&= [\omega(\mathbf{u}), \mathbf{v}(\omega(\mathbf{w}))] - [\omega(\mathbf{u}), \mathbf{w}(\omega(\mathbf{v}))] - [\omega(\mathbf{u}), \omega([\mathbf{v}, \mathbf{w}])] \\
&\quad + [\omega(\mathbf{v}), \mathbf{w}(\omega(\mathbf{u}))] - [\omega(\mathbf{v}), \mathbf{u}(\omega(\mathbf{w}))] - [\omega(\mathbf{v}), \omega([\mathbf{w}, \mathbf{u}])] \\
&\quad + [\omega(\mathbf{w}), \mathbf{u}(\omega(\mathbf{v}))] - [\omega(\mathbf{w}), \mathbf{v}(\omega(\mathbf{u}))] - [\omega(\mathbf{w}), \omega([\mathbf{u}, \mathbf{v}])] \\
&= \left([\omega(\mathbf{u}), \mathbf{v}(\omega(\mathbf{w}))] + [\mathbf{v}(\omega(\mathbf{u})), \omega(\mathbf{w})] \right) - [\omega(\mathbf{u}), \omega([\mathbf{v}, \mathbf{w}])] \\
&\quad + \left([\omega(\mathbf{v}), \mathbf{w}(\omega(\mathbf{u}))] + [\mathbf{w}(\omega(\mathbf{v})), \omega(\mathbf{u})] \right) + [\omega(\mathbf{v}), \omega([\mathbf{w}, \mathbf{u}])] \\
&\quad + \left([\omega(\mathbf{w}), \mathbf{u}(\omega(\mathbf{v}))] + [\mathbf{u}(\omega(\mathbf{w})), \omega(\mathbf{v})] \right) - [\omega(\mathbf{w}), \omega([\mathbf{u}, \mathbf{v}])]
\end{aligned}$$

The derivation $\mathbf{u}$ on $\mathcal{O}_P \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$ is by definition of the form $\mathbf{u} = \mathbf{t}_u \otimes \text{id}$, where $\mathbf{t}_u$ is a derivation on $\mathcal{O}_P$. An analogous statement is of course true for $\mathbf{v}$ and $\mathbf{w}$. Recalling the Lie-algebra structure of $\mathcal{O}_P \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$ from Remark 8.13, we see that the equation

$$\mathbf{u}([a, b]) = [\mathbf{u}(a), b] + [a, \mathbf{u}(b)] \quad (*)$$

holds for all $a, b \in \mathcal{O}_P \otimes_{\mathcal{O}_S(S)} \mathfrak{g}$. Thus we find indeed $\frac{1}{2}d[\omega, \omega] + [\omega, d\omega] = 0$, and it only remains to prove the relation (*). Writing $a = \sum_i f_i \otimes r_i$ and $b = \sum_j g_j \otimes s_j$ this may be seen as follows:

$$\begin{aligned}
\mathbf{u}([a, b]) &= \mathbf{u}\left(\left[\sum_i f_i \otimes r_i, \sum_j g_j \otimes s_j\right]\right) = \sum_{i,j} (\mathbf{t}_u \otimes \text{id}) \underbrace{([f_i \otimes r_i, g_j \otimes s_j])}_{=f_i \cdot g_j \otimes [r_i, s_j]} \\
&= \sum_{i,j} \mathbf{t}_u(f_i \cdot g_j) \otimes [r_i, s_j] = \sum_{i,j} \left(\mathbf{t}_u(f_i) \cdot g_j + f_i \cdot \mathbf{t}_u(g_j) \right) \otimes [r_i, s_j] \\
&= \sum_{i,j} \mathbf{t}_u(f_i) \cdot g_j \otimes [r_i, s_j] + \sum_{i,j} f_i \cdot \mathbf{t}_u(g_j) \otimes [r_i, s_j] \\
&= \sum_{i,j} \left[\mathbf{t}_u(f_i) \otimes r_i, g_j \otimes s_j \right] + \sum_{i,j} \left[f_i \otimes r_i, \mathbf{t}_u(g_j) \otimes s_j \right] \\
&= \left[\sum_i \mathbf{u}(f_i \otimes r_i), \sum_j g_j \otimes s_j \right] + \left[\sum_i f_i \otimes r_i, \sum_j \mathbf{u}(g_j \otimes s_j) \right] \\
&= [\mathbf{u}(a), b] + [a, \mathbf{u}(b)].
\end{aligned}$$

Let us finally prove that also $[\omega, [\omega, \omega]] = 0$. This follows from the Jacobi-identity of

Lie-algebras:

$$\begin{aligned}
[\omega, [\omega, \omega]](\mathbf{u}, \mathbf{v}, \mathbf{w}) &= [\omega(\mathbf{u}), [\omega, \omega](\mathbf{v}, \mathbf{w})] - [\omega(\mathbf{v}), [\omega, \omega](\mathbf{u}, \mathbf{w})] + [\omega(\mathbf{w}), [\omega, \omega](\mathbf{u}, \mathbf{v})] \\
&= 2[\omega(\mathbf{u}), [\omega(\mathbf{v}), \omega(\mathbf{w})]] - 2[\omega(\mathbf{v}), [\omega(\mathbf{u}), \omega(\mathbf{w})]] + 2[\omega(\mathbf{w}), [\omega(\mathbf{u}), \omega(\mathbf{v})]] \\
&= 2\left([\omega(\mathbf{u}), [\omega(\mathbf{v}), \omega(\mathbf{w})]] + [\omega(\mathbf{v}), [\omega(\mathbf{w}), \omega(\mathbf{u})]] + [\omega(\mathbf{w}), [\omega(\mathbf{u}), \omega(\mathbf{v})]]\right) \\
&= 0.
\end{aligned}$$

□

10.7 Remark Let α be a differential k -form on the X -torsor P under G_X , and let ω be a connection form. Then the results of this section motivate the following alternative definition of covariant derivation on P . We define the covariant derivation $D\alpha$ of α with respect to ω by the formula

$$D\alpha := d\alpha + [\omega, \alpha].$$

In particular, Lemma 10.5 and the commutativity of the exterior derivation d with pull-backs under closed immersions (see proof of Proposition 9.10) show that this definition of the covariant derivation D on P yields the identity

$$D_X(s^*\alpha) = s^*(D\alpha),$$

where $s : X \hookrightarrow P$ is a section of $\pi : P \rightarrow X$, and where D_X is the covariant derivation on X in the sense of Definition 10.3. Thus the covariant derivation D_X on X may again be interpreted as pull-back of the covariant derivation D on P .

10.2. Yang-Mills equation in local coordinates

Let $g : T_{X/S} \times_X T_{X/S} \rightarrow \mathbb{A}_X^1$ be the metric which may be considered as well as a $\mathcal{O}_X(X)$ -bilinear homomorphism $g : \mathcal{T}_{X/S}(X) \times \mathcal{T}_{X/S}(X) \rightarrow \mathcal{O}_X(X)$ due to our expositions at the beginning of section 11.3. In particular, we obtain a family of $\mathcal{O}_{X,x}$ -bilinear morphisms

$$g_x : \mathcal{T}_{X/S,x} \times \mathcal{T}_{X/S,x} \rightarrow \mathcal{O}_{X,x}$$

on stalks for all $x \in X$ which may also be considered as a family of $\mathcal{O}_{X,x}$ -linear isomorphisms $g_x : \mathcal{T}_{X/S,x} \xrightarrow{\sim} \Omega_{X/S,x}^1$. One has got $g_{\mu\nu,x} := g_x\left(\frac{\partial}{\partial x^\mu}\right)\left(\frac{\partial}{\partial x^\nu}\right) = \sum_{\rho=1}^n g_{\mu\rho,x} dx^\rho \left(\frac{\partial}{\partial x^\nu}\right)$, i.e.

$$g_x\left(\frac{\partial}{\partial x^\mu}\right) = \sum_{\rho=1}^n g_{\mu\rho,x} dx^\rho.$$

Then the inverse map $g_x^{-1} : \Omega_{X/S,x}^1 \rightarrow \mathcal{T}_{X/S,x}$ is given by $\frac{\partial}{\partial x^\mu} = \sum_{\rho=1}^n g_{\mu\rho,x} g_x^{-1}(dx^\rho)$. Let $g_x^{\mu\nu}$ be the components of the inverse matrix of $(g_{\mu\nu,x})_{\mu\nu}$, i.e. $\sum_{\rho=1}^n g_{\mu\rho} g^{\rho\nu} = \delta_\mu^\nu$. Then

$$g_x^{-1}(dx^\mu) = \sum_{\rho=1}^n g_x^{\mu\rho} \frac{\partial}{\partial x^\rho}.$$

Again, let us write $g_{\mu\nu}$ (resp. $g^{\mu\nu}$) instead of $g_{\mu\nu,x}$ (resp. $g_x^{\mu\nu}$) whenever no confusion is possible. Now we are prepared to derive the Hodge-star operator in local coordinates. Let $\alpha \in \Omega_{X/S}^k(X)$ be a differential k -form on X . Then by Definition 11.42 we have got

$$*\alpha := (\Lambda^k g^{-1})(\alpha)]v_g.$$

Writing

$$\alpha_x = \frac{1}{k!} \sum_{\mu_1, \dots, \mu_k=1}^n \alpha_{\mu_1 \mu_2 \dots \mu_k} dx^{\mu_1} \wedge dx^{\mu_2} \wedge \dots \wedge dx^{\mu_k},$$

where the coefficients $\alpha_{\mu_1 \mu_2 \dots \mu_k}$ are totally antisymmetric, we obtain the first part of the Hodge-star operator

$$(\Lambda^k g^{-1} \alpha)_x = \frac{1}{k!} \sum_{\mu_1, \dots, \mu_k, \nu_1, \dots, \nu_k=1}^n g^{\mu_1 \nu_1} \dots g^{\mu_k \nu_k} \alpha_{\mu_1 \dots \mu_k} \frac{\partial}{\partial x^{\nu_1}} \wedge \dots \wedge \frac{\partial}{\partial x^{\nu_k}}.$$

Let us now choose a volume form (which exists due to Theorem 11.61). As we are working with respect to étale topology, the local rings $\mathcal{O}_{X,x}$ are strictly henselian. Therefore $\sqrt{\det g_x} \in \mathcal{O}_{X,x}$, where $\det g_x$ denotes the determinant of the matrix $(g_{\mu\nu,x})_{\mu\nu}$. Then

$$v_{g,x} := \sqrt{\det g_x} dx^1 \wedge \dots \wedge dx^n$$

is a volume form which is independent of the choice of the base $\{dx^\mu\}_{\mu=1}^n$ of $\Omega_{X/S,x}^1$. We obtain

$$\begin{aligned} (*\alpha)_x \left(\frac{\partial}{\partial x^{\nu_{k+1}}}, \dots, \frac{\partial}{\partial x^{\nu_n}} \right) &= \left(\Lambda^k g^{-1}(\alpha)]v_g \right)_x \left(\frac{\partial}{\partial x^{\nu_{k+1}}}, \dots, \frac{\partial}{\partial x^{\nu_n}} \right) \\ &= \frac{1}{k!} \sum_{\mu_1, \dots, \mu_k, \nu_1, \dots, \nu_k=1}^n \sqrt{\det g_x} g^{\mu_1 \nu_1} \dots g^{\mu_k \nu_k} \alpha_{\mu_1 \dots \mu_k} (dx^1 \wedge \dots \wedge dx^n) \left(\frac{\partial}{\partial x^{\nu_1}}, \dots, \frac{\partial}{\partial x^{\nu_n}} \right) \\ &= \frac{1}{k!} \sum_{\mu_1, \dots, \mu_k, \nu_1, \dots, \nu_k=1}^n \sqrt{\det g_x} g^{\mu_1 \nu_1} \dots g^{\mu_k \nu_k} \alpha_{\mu_1 \dots \mu_k} \underbrace{\sum_{\pi \in S_n} \sigma_\pi \delta_{\nu_1}^{\pi(1)} \dots \delta_{\nu_n}^{\pi(n)}}_{=: \epsilon_{\nu_1 \dots \nu_n}} \\ &= \frac{1}{k!(n-k)!} \sum_{\mu_1, \dots, \mu_k, \rho_1, \dots, \rho_n=1}^n \sqrt{\det g_x} g^{\mu_1 \rho_1} \dots g^{\mu_k \rho_k} \epsilon_{\rho_1 \dots \rho_n} \alpha_{\mu_1 \dots \mu_k} \\ &\quad \cdot (dx^{\rho_{k+1}} \wedge \dots \wedge dx^{\rho_n}) \left(\frac{\partial}{\partial x^{\nu_{k+1}}}, \dots, \frac{\partial}{\partial x^{\nu_n}} \right) \end{aligned}$$

and we may summarize as follows.

10.8 Proposition *Let $x \in X$ be a physical point, let $\alpha \in \Omega_{X/S}^k(X)$ be a differential k -form, and let $*$ be the Hodge-star operator. Then the following equation holds at x :*

$$\begin{aligned} (*\alpha)_x &= \frac{1}{k!(n-k)!} \sum_{\mu_1, \dots, \mu_k, \rho_1, \dots, \rho_n=1}^n \sqrt{\det g_x} g^{\mu_1 \rho_1} \dots g^{\mu_k \rho_k} \epsilon_{\rho_1 \dots \rho_n} \alpha_{\mu_1 \dots \mu_k} dx^{\rho_{k+1}} \wedge \dots \wedge dx^{\rho_n} \\ &= \frac{1}{k!(n-k)!} \sum_{\rho_1, \dots, \rho_n=1}^n \sqrt{\det g_x} \epsilon_{\rho_1 \dots \rho_n} \alpha^{\rho_1 \dots \rho_k} dx^{\rho_{k+1}} \wedge \dots \wedge dx^{\rho_n}. \end{aligned}$$

10.9 Yang-Mills equation *Let $x \in X$ be a physical point. Then the Yang-Mills equation at x reads as follows:*

$$0 = \sum_{\mu, \rho, \sigma=1}^n \left(\frac{1}{\sqrt{\det g_x}} \frac{\partial}{\partial x^\sigma} \left(\sqrt{\det g_x} g^{\mu\nu} g^{\rho\sigma} \mathcal{F}_{\mu\rho}^k \right) + \sum_{i,j=1}^N c_{ij}^k g^{\mu\nu} g^{\rho\sigma} \mathcal{A}_\sigma^i \mathcal{F}_{\mu\rho}^j \right)$$

(where ν and k are non-contracted indices).

PROOF Due to Proposition 10.8, the Hodge-star of the field strength at x is

$$(*\mathcal{F})_x = \frac{1}{2!(n-2)!} \sum_{\mu_1, \mu_2, \rho_1, \dots, \rho_n=1}^n \sqrt{\det g_x} g^{\mu_1 \rho_1} g^{\mu_2 \rho_2} \epsilon_{\rho_1 \dots \rho_n} \mathcal{F}_{\mu_1 \mu_2} dx^{\rho_3} \wedge \dots \wedge dx^{\rho_n}.$$

By Definition 10.3 one has got

$$\begin{aligned} (D_X(*\mathcal{F}))_x \left(\frac{\partial}{\partial x^{\nu_2}}, \dots, \frac{\partial}{\partial x^{\nu_n}} \right) &= (d(*\mathcal{F}))_x \left(\frac{\partial}{\partial x^{\nu_2}}, \dots, \frac{\partial}{\partial x^{\nu_n}} \right) \\ &\quad + \sum_{i=2}^n (-1)^i \left[\mathcal{A}_x \left(\frac{\partial}{\partial x^{\nu_i}} \right), (*\mathcal{F})_x \left(\frac{\partial}{\partial x^{\nu_2}}, \dots, \frac{\partial}{\partial x^{\nu_i}}, \dots, \frac{\partial}{\partial x^{\nu_n}} \right) \right]. \end{aligned}$$

The first summand on the right hand side is

$$\begin{aligned} &(d(*\mathcal{F}))_x \\ &= \frac{1}{2!(n-2)!} \sum_{\mu_1, \mu_2, \mu_3, \rho_1, \dots, \rho_n=1}^n \frac{\partial}{\partial x^{\mu_3}} \left(\sqrt{\det g_x} g^{\mu_1 \rho_1} g^{\mu_2 \rho_2} \epsilon_{\rho_1 \dots \rho_n} \mathcal{F}_{\mu_1 \mu_2} \right) dx^{\mu_3} \wedge dx^{\rho_3} \wedge \dots \wedge dx^{\rho_n}, \end{aligned}$$

and the second one may be written as follows:

$$\begin{aligned} &\sum_{\iota=2}^n (-1)^\iota \left[\mathcal{A}_x \left(\frac{\partial}{\partial x^{\nu_\iota}} \right), (*\mathcal{F})_x \left(\frac{\partial}{\partial x^{\nu_2}}, \dots, \frac{\partial}{\partial x^{\nu_\iota}}, \dots, \frac{\partial}{\partial x^{\nu_n}} \right) \right] \\ &= \sum_{i,j,k=1}^N c_{ij}^k b_k \sum_{\iota=2}^n (-1)^\iota \mathcal{A}_x^i \left(\frac{\partial}{\partial x^{\nu_\iota}} \right) (*\mathcal{F}^j)_x \left(\frac{\partial}{\partial x^{\nu_2}}, \dots, \frac{\partial}{\partial x^{\nu_\iota}}, \dots, \frac{\partial}{\partial x^{\nu_n}} \right) \\ &= \sum_{i,j,k=1}^N c_{ij}^k b_k (\mathcal{A}^i \wedge (*\mathcal{F}^j))_x \left(\frac{\partial}{\partial x^{\nu_2}}, \dots, \frac{\partial}{\partial x^{\nu_n}} \right). \end{aligned}$$

Thereby

$$\begin{aligned} &(\mathcal{A}^i \wedge (*\mathcal{F}^j))_x \\ &= \frac{1}{2!(n-2)!} \sum_{\mu_1, \mu_2, \mu_3, \rho_1, \dots, \rho_n=1}^n \sqrt{\det g_x} g^{\mu_1 \rho_1} g^{\mu_2 \rho_2} \epsilon_{\rho_1 \dots \rho_n} \mathcal{A}_{\mu_3}^i \mathcal{F}_{\mu_1 \mu_2}^j dx^{\mu_3} \wedge dx^{\rho_3} \wedge \dots \wedge dx^{\rho_n} \end{aligned}$$

due to the above expression for $(*\mathcal{F})_x$. After substitution of these relations into the above formula for $(D_X(*\mathcal{F}))_x$, we finally have to apply the Hodge-star operator, in order to

obtain the differential form occurring in Yang-Mills equation. $(D_X(*\mathcal{F}))_x$ is a differential $(n-1)$ -form, and thus the Hodge-star of this form is the following differential 1-form:

$$\begin{aligned}
(*D_X * \mathcal{F})_x &= \sum_{k=1}^N b_k * \left(d(*\mathcal{F}_k) + \sum_{i,j=1}^N c_{ij}^k \mathcal{A}^i \wedge (*\mathcal{F}^j) \right) \\
&= \frac{1}{2!(n-2)!} \sum_{k=1}^N b_k \sum_{\mu_1, \mu_2, \sigma_2, \rho_1, \dots, \rho_n, \nu_1, \dots, \nu_n} \sqrt{\det g_x} g^{\sigma_2 \nu_2} g^{\rho_3 \nu_3} \dots g^{\rho_n \nu_n} \epsilon_{\nu_1 \dots \nu_n} \epsilon_{\rho_1 \dots \rho_n} \\
&\quad \cdot \left(\frac{\partial}{\partial x^{\sigma_2}} \left(\sqrt{\det g_x} g^{\mu_1 \rho_1} g^{\mu_2 \rho_2} \mathcal{F}_{\mu_1 \mu_2}^k \right) + \sum_{i,j=1}^N c_{ij}^k \sqrt{\det g_x} g^{\mu_1 \rho_1} g^{\mu_2 \rho_2} \mathcal{A}_{\sigma_2}^i \mathcal{F}_{\mu_1 \mu_2}^j \right) dx^{\nu_1} \\
&= \frac{1}{2!(n-2)!} \sum_{k=1}^N b_k \sum_{\mu_1, \mu_2, \sigma_1, \sigma_2, \sigma_3, \rho_1, \dots, \rho_n} \sqrt{\det g_x} \underbrace{\sum_{\nu_1, \dots, \nu_n} g^{\sigma_1 \nu_1} g^{\sigma_2 \nu_2} g^{\rho_3 \nu_3} \dots g^{\rho_n \nu_n} \epsilon_{\nu_1 \dots \nu_n} \epsilon_{\rho_1 \dots \rho_n}}_{=(\det g_x^{-1}) \cdot \epsilon_{\sigma_1 \sigma_2 \rho_3 \dots \rho_n}} \\
&\quad \cdot \left(\frac{\partial}{\partial x^{\sigma_2}} \left(\sqrt{\det g_x} g^{\mu_1 \rho_1} g^{\mu_2 \rho_2} \mathcal{F}_{\mu_1 \mu_2}^k \right) + \sum_{i,j=1}^N c_{ij}^k \sqrt{\det g_x} g^{\mu_1 \rho_1} g^{\mu_2 \rho_2} \mathcal{A}_{\sigma_2}^i \mathcal{F}_{\mu_1 \mu_2}^j \right) g_{\sigma_1 \sigma_3} dx^{\sigma_3} \\
&\quad \text{(where we also made use of the relation } g^{\sigma_1 \nu_1} g_{\sigma_1 \sigma_3} dx^{\sigma_3} = \delta_{\sigma_3}^{\nu_1} dx^{\sigma_3} = dx^{\nu_1} \text{)} \\
&= \frac{1}{2!(n-2)!} \sum_{k=1}^N b_k \sum_{\mu_1, \mu_2, \sigma_1, \sigma_2, \sigma_3, \rho_1, \rho_2} \frac{1}{\sqrt{\det g_x}} g_{\sigma_1 \sigma_3} \underbrace{\sum_{\rho_3, \dots, \rho_n} \epsilon_{\sigma_1 \sigma_2 \rho_3 \dots \rho_n} \epsilon_{\rho_1 \rho_2 \rho_3 \dots \rho_n}}_{=\delta_{\rho_1}^{\sigma_1} \delta_{\rho_2}^{\sigma_2} - \delta_{\rho_2}^{\sigma_1} \delta_{\rho_1}^{\sigma_2}} \\
&\quad \cdot \left(\frac{\partial}{\partial x^{\sigma_2}} \left(\sqrt{\det g_x} g^{\mu_1 \rho_1} g^{\mu_2 \rho_2} \mathcal{F}_{\mu_1 \mu_2}^k \right) + \sum_{i,j=1}^N c_{ij}^k \sqrt{\det g_x} g^{\mu_1 \rho_1} g^{\mu_2 \rho_2} \mathcal{A}_{\sigma_2}^i \mathcal{F}_{\mu_1 \mu_2}^j \right) dx^{\sigma_3} \\
&= \frac{1}{2!(n-2)!} \sum_{k=1}^N b_k \sum_{\mu_1, \mu_2, \sigma_1, \sigma_2, \sigma_3, \rho_1, \rho_2} \frac{1}{\sqrt{\det g_x}} g_{\sigma_1 \sigma_3} (\delta_{\rho_1}^{\sigma_1} \delta_{\rho_2}^{\sigma_2} - \delta_{\rho_2}^{\sigma_1} \delta_{\rho_1}^{\sigma_2}) \\
&\quad \cdot \left(\frac{\partial}{\partial x^{\sigma_2}} \left(\sqrt{\det g_x} g^{\mu_1 \rho_1} g^{\mu_2 \rho_2} \mathcal{F}_{\mu_1 \mu_2}^k \right) + \sum_{i,j=1}^N c_{ij}^k \sqrt{\det g_x} g^{\mu_1 \rho_1} g^{\mu_2 \rho_2} \mathcal{A}_{\sigma_2}^i \mathcal{F}_{\mu_1 \mu_2}^j \right) dx^{\sigma_3} \\
&= \frac{1}{(n-2)!} \sum_{k=1}^N b_k \sum_{\mu_1, \mu_2, \sigma_1, \sigma_2, \sigma_3} \frac{1}{\sqrt{\det g_x}} g_{\sigma_1 \sigma_3} \\
&\quad \cdot \left(\frac{\partial}{\partial x^{\sigma_2}} \left(\sqrt{\det g_x} g^{\mu_1 \sigma_1} g^{\mu_2 \sigma_2} \mathcal{F}_{\mu_1 \mu_2}^k \right) + \sum_{i,j=1}^N c_{ij}^k \sqrt{\det g_x} g^{\mu_1 \sigma_1} g^{\mu_2 \sigma_2} \mathcal{A}_{\sigma_2}^i \mathcal{F}_{\mu_1 \mu_2}^j \right) dx^{\sigma_3},
\end{aligned}$$

because $\delta_{\rho_1}^{\sigma_1} \delta_{\rho_2}^{\sigma_2} - \delta_{\rho_2}^{\sigma_1} \delta_{\rho_1}^{\sigma_2}$ yields a factor 2 due to the antisymmetry $\mathcal{F}_{\mu\nu}^i = -\mathcal{F}_{\nu\mu}^i$ of the field strength. As $\{dx^\mu\}_{\mu=1}^n$ forms a base of $\Omega_{X/S}^1(X)$, as $\{b_i\}_{i=1}^N$ forms a base of $\mathfrak{g}$, and due to the fact that the metric $(g_{\mu\nu})_{\mu\nu}$ is an isomorphism, we see that the Yang-Mills equation reads as claimed. $\square$

11. Appendix II

In ordinary (i.e. $\mathbb{R}$ -valued) Yang-Mills theory, gauge fields are described by co-vector-fields. If we want to generalize the $\mathbb{R}$ -valued, differential geometric Yang-Mills theory to arbitrary commutative rings R (or even base schemes S), we therefore have to supply the notion of the tangent bundle in the realm of algebraic geometry. In this chapter 11 we are going to recall some fundamental results which we will make use of later. First we remember of the more special notion of Zariski tangent-vectors, and then we will recall the general notion as it is found in [SGA 3].

As the gauge group in classical Yang-Mills theory is given by a Lie-group, we have to introduce the notion of group functors. In regard to Yang-Mills theory it is particularly important that the maps Ad and ad have algebraic geometric analogues.

11.1. The Zariski tangent space

11.1 Definition a) Let X be a scheme and $x \in X$. Let $\mathfrak{m}_x$ be the maximal ideal of $\mathcal{O}_{X,x}$ and $k(x) = \mathcal{O}_{X,x}/\mathfrak{m}_x$ be the residue field. Then $\mathfrak{m}_x/\mathfrak{m}_x^2 = \mathfrak{m}_x \otimes_{\mathcal{O}_{X,x}} k(x)$ is in a natural way a $k(x)$ -vector space. Its dual $(\mathfrak{m}_x/\mathfrak{m}_x^2)^\vee$ is called the *Zariski tangent space* to X at x . We denote it by $T_{X,x}$.

b) Let $f : X \rightarrow Y$ be a morphism of schemes, let $x \in X$, $y = f(x)$. Then $f_x^\# : \mathcal{O}_{Y,y} \rightarrow \mathcal{O}_{X,x}$ canonically induces a $k(x)$ -linear map

$$T_{f,x} : T_{X,x} \rightarrow T_{Y,y} \otimes_{k(y)} k(x)$$

the *tangent map* of f at x .

If $X \rightarrow S$ is a smooth morphism, then X is regular, i.e. $\dim \mathcal{O}_{X,x} = \dim_{k(x)} T_{X,x}$ for all $x \in X$. Let us see that we are reduced to the ordinary differential geometric notions in the standard situation of classical physics.

physical interpretation: Let $S = \text{Spec } \mathbb{R}$ and let $X = V(f) \hookrightarrow \mathbb{A}_{\mathbb{R}}^n$ be smooth over S (e.g. $f = \sum_{i=1}^n X_i^2 - 1$, i.e. $X(\mathbb{R}) = (n-1)$ -sphere). Let $x \in X(\mathbb{R})$ be a closed point. Then each irreducible component of X has the same dimension ([Liu], 2.5.26). So we may assume that X is irreducible. Then we get:

$$\dim \mathcal{O}_{X,x} = \dim X = \dim \mathbb{A}_{\mathbb{R}}^n - 1 = \underbrace{\dim \text{Spec } \mathbb{R}}_{=0} + n - 1 = n - 1.$$

The first equality is due to [Liu], Cor. 2.5.24, the second one is due to [Liu], Cor. 2.5.26, and finally the third equality follows from [Liu], Cor. 2.5.17. Therefore, $\dim_{\mathbb{R}} T_{X,x} =$

$n - 1$. Furthermore, if we write the polynomial f in the form $f = f(T_1, \dots, T_n)$, one has due to [Liu], Prop. 4.2.5

$$T_{X,x} = \left\{ (t_1, \dots, t_n) \in \mathbb{R}^n \mid \sum_{i=1}^n \frac{\partial f}{\partial T_i}(x) t_i = 0 \right\}.$$

Thus $T_{X,x}$ is really the tangent space in the classical sense.

Important is the following theorem which justifies the notion of physical points (see Definition 6.9). Let us stipulate that we denote the dual of a vector space V by $V^\vee$.

11.2 Theorem *Let $f : X \rightarrow S$ be a smooth morphism, $s \in S$ and $x \in X_s$ closed. Then*

$$(\Omega_{X/S}^1 \otimes_{\mathcal{O}_{X,x}} k(x))^\vee = T_{X_s,x} \oplus (\Omega_{k(x)/k(s)}^1)^\vee.$$

Especially for physical points we get

$$(\Omega_{X/S}^1 \otimes_{\mathcal{O}_{X,x}} k(x))^\vee = T_{X_s,x}.$$

PROOF [Liu], Ex. 6.2.5 □

Remark: By the preceding theorem we establish the desired duality of tangent and co-tangent vectors at physical points.

11.3 Definition a) Let $(A, \mathfrak{m})$ be a regular noetherian local ring of dimension d . Any system of generators of $\mathfrak{m}$ with d elements is coordinate system for A .

b) Especially if $X \rightarrow S$ is smooth, $\mathcal{O}_{X,x}$ is a regular noetherian local ring. A local coordinate system at $x \in X$ is a coordinate system for $\mathcal{O}_{X,x}$. In particular, the residue class of a local coordinate system at $x \in X$ gives rise to a basis of the tangent space $T_{X,x}$ and conversely by Nakayama's lemma.

11.2. The tangent bundle

Let us now summarize the construction of the tangent bundle following [SGA 3]. We will see that, in physical points, this definition reduces to the Zariski tangent space. But it is designed in such a way that it also preserves all classical properties of the tangent bundle (like the duality between tangent and co-tangent vectors one is used to from differential geometry) in non-physical points, too. Therefore we will use this notion for our purposes.

11.4 Definition Let S be a scheme and $\mathcal{M}$ a quasi-coherent $\mathcal{O}_S$ -module. Let $D_{\mathcal{O}_S}(\mathcal{M})$ denote the quasi-coherent $\mathcal{O}_S$ -algebra $\mathcal{O}_S \oplus \mathcal{M}$ (where $\mathcal{M}$ is considered as $\mathcal{O}_S$ -algebra via the definition $\mathcal{M} \cdot \mathcal{M} = 0$). Then

$$I_S(\mathcal{M}) := \underline{\text{Spec}} D_{\mathcal{O}_S}(\mathcal{M}).$$

In particular we define:

$$\begin{aligned} D_{\mathcal{O}_S} &:= D_{\mathcal{O}_S}(\mathcal{O}_S) \\ I_S &:= I_S(\mathcal{O}_S). \end{aligned}$$

11.5 Definition Let S be a scheme. For all $\mathcal{O}_S$ -modules $\mathcal{F}$ we define the functors $V(\mathcal{F})$ and $W(\mathcal{F})$ upon $(Sch)/S$ ¹ by:

$$\begin{aligned} V(\mathcal{F})(S') &:= \text{Hom}_{\mathcal{O}_{S'}}(\mathcal{F} \otimes \mathcal{O}_{S'}, \mathcal{O}_{S'}) \\ W(\mathcal{F})(S') &:= \Gamma(S', \mathcal{F} \otimes \mathcal{O}_{S'}). \end{aligned}$$

11.6 Definition Let S be a scheme and X an S -functor. We say that X verifies (E) relative to S , if for all $S' \rightarrow S$ and for all free $\mathcal{O}_{S'}$ -modules of finite type $\mathcal{M}$ and $\mathcal{N}$ the commutative diagram

$$\begin{array}{ccc} & X(I_{S'}(\mathcal{M} \oplus \mathcal{N})) & \\ \swarrow & & \searrow \\ X(I_{S'}(\mathcal{M})) & & X(I_{S'}(\mathcal{N})) \\ \searrow & & \swarrow \\ & X(S') & \end{array}$$

obtained by applying first $I_{S'}$ and then X to the canonical commutative diagram

$$\begin{array}{ccc} & \mathcal{M} \oplus \mathcal{N} & \\ \swarrow & & \searrow \\ \mathcal{M} & & \mathcal{N} \\ \searrow & & \swarrow \\ & 0 & \end{array}$$

is cartesian. (Recall that $I_{S'}(0) \cong S'$.)

Remark: If X is an S -scheme, it verifies condition (E). In the situation of Definition 11.6 we will sometimes simply say: X/S verifies (E).

11.7 Definition Let $\underline{\mathcal{Q}}$ denote the functor

$$S \rightsquigarrow \Gamma(S, \mathcal{O}_S)$$

provided with its structure of an $\widehat{(Sch)}$ -ring.² In particular $\underline{\mathcal{Q}}$ is represented by the scheme $\text{Spec } \mathbb{Z}[T] = \mathbb{A}_{\mathbb{Z}}^1$. This induces by base change over $\mathbb{Z}$ an affine S -ring which we denote by $\underline{\mathcal{Q}}_S$.

Now we are prepared to introduce the tangent bundle.

¹ (Sch) denotes the category of schemes, and $(Sch)/S$ denotes the category of S -schemes, i.e. the objects are schemes X which come with a unique morphisms $X \rightarrow S$

²If $\mathfrak{C}$ denotes a category, then $\widehat{\mathfrak{C}}$ denotes the category of contravariant functors on $\mathfrak{C}$.

11.2.1. The tangent bundle as a functor.

11.8 Definition Let S be a scheme and $\mathcal{M}$ an $\mathcal{O}_S$ -module of finite type. Let $f : X \rightarrow S$ be a scheme over S . The relative tangent bundle of X over S relative to $\mathcal{M}$ is the S -functor³

$$T_{X/S}(\mathcal{M}) := \underline{\mathrm{Hom}}_S(I_S(\mathcal{M}), X).$$

The relative tangent bundle of X over S is the S -functor

$$T_{X/S} := T_{X/S}(\mathcal{O}_S) = \underline{\mathrm{Hom}}_S(I_S, X).$$

$\mathcal{M} \rightsquigarrow T_{X/S}(\mathcal{M})$ is a covariant functor from the category of free $\mathcal{O}_S$ -modules of finite type to the category of S -functors. In particular, we get two canonical morphisms of S -functors:

- 1.) structure morphism: $T_{X/S}(\mathcal{M}) \rightarrow X$
- 2.) zero section: $X \rightarrow T_{X/S}(\mathcal{M})$

induced by the canonical morphism $0 \rightarrow \mathcal{M}$ and $\mathcal{M} \rightarrow 0$ (recall: $T_{X/S}(0) \cong X$).

11.9 Proposition Let $I_S := I_S(\mathcal{O}_S)$. Then there is an isomorphism of X -functors:

$$\underline{\mathrm{Hom}}_S(I_S, X) \cong V(\Omega_{X/S}^1).$$

PROOF Let $X' \xrightarrow{x'} X$ be an object of $(Sch)/X$. Then

$$\begin{aligned} \underline{\mathrm{Hom}}_S(I_S, X)(X') & \underset{\text{in } (Sch)/X}{=} X(\mathcal{E}_{\mathcal{O}_{X'}})^{-1}(x') \\ & = \mathrm{Hom}_{\mathcal{O}_{X'}}((x')^* \Omega_{X/S}^1, \mathcal{O}_{X'}) \quad \text{by [SGA 3], Expose II, Prop. 2.2} \\ & = V(\Omega_{X/S}^1)(X') \quad \text{by Definition 11.5.} \end{aligned}$$

In particular, $\mathcal{M} \rightsquigarrow T_{X/S}(\mathcal{M})$ is already a covariant functor

$$\begin{aligned} (\text{free } \mathcal{O}_S - \text{modules of finite type}) & \rightarrow (X - \text{functors}) \\ \mathcal{M} & \rightsquigarrow T_{X/S}(\mathcal{M}). \end{aligned}$$

11.10 Definition Let $u \in X(S) = \Gamma(X/S)$. The (relative) tangent space of X over S at u relative to $\mathcal{M}$ is the S -functor

$$L_{X/S}^u(\mathcal{M}),$$

³For all S -functors $\mathcal{F}, \mathcal{G}$ and for all S -morphisms $S' \rightarrow S$ one has got by definition

$$\underline{\mathrm{Hom}}_S(\mathcal{F}, \mathcal{G})(S') := \mathrm{Hom}_{S'}(\mathcal{F}_{S'}, \mathcal{G}_{S'}),$$

where $\mathcal{F}_{S'}$ (resp. $\mathcal{G}_{S'}$) denotes the restriction of $\mathcal{F}$ (resp. $\mathcal{G}$) from $(Sch)/S$ to $(Sch)/S'$.

given by the inverse image of the X -functor $T_{X/S}(\mathcal{M})$ under the morphism $u : S \rightarrow X$; i.e. we have a pull-back diagram

$$\begin{array}{ccc} L_{X/S}^u(\mathcal{M}) & \longrightarrow & T_{X/S}(\mathcal{M}) \\ \downarrow & & \downarrow \\ S & \longrightarrow & X \end{array}$$

The (relative) tangent space of X over S at u is the S -functor

$$L_{X/S}^u := L_{X/S}^u(\mathcal{O}_S).$$

11.11 Proposition *The functors $T_{X/S}(\mathcal{M})$ and $L_{X/S}^u(\mathcal{M})$ are functorial in X : If $f : X \rightarrow X'$ is an S -morphism, then there are commutative diagrams:*

$$\begin{array}{ccc} T_{X/S}(\mathcal{M}) & \xrightarrow{T(f)} & T_{X'/S}(\mathcal{M}) \\ \downarrow & & \downarrow \\ X & \xrightarrow{f} & X' \end{array} \quad \begin{array}{ccc} L_{X/S}^u(\mathcal{M}) & \xrightarrow{T(f)} & L_{X'/S}^{f \circ u}(\mathcal{M}) \\ & \searrow & \swarrow \\ & S & \end{array}$$

PROOF [SGA 3], Expose II, Prop. 3.7 □

11.12 Remark If $f : X \rightarrow X'$ is étale, the above square is cartesian. In general there is a morphism of X -functors:⁴

$$T_{X/S}(\mathcal{M}) \rightarrow (T_{X'/S}(\mathcal{M}))_X.$$

The morphism $T_{X/S}(\mathcal{M}) \rightarrow (T_{X'/S}(\mathcal{M}))_X$ (resp. $L_{X/S}^u(\mathcal{M}) \rightarrow L_{X'/S}^{f \circ u}(\mathcal{M})$) is a morphism of $\mathcal{O}_X$ -modules (resp. $\mathcal{O}_S$ -modules) if X/S and X'/S verify (E).

11.13 Proposition *Let X and Y be two functors above S . Then there are isomorphisms which are functorial in $\mathcal{M}$.*

$$\begin{aligned} T_{X \times_S Y/S}(\mathcal{M}) &\xrightarrow{\sim} T_{X/S}(\mathcal{M}) \times_S T_{Y/S}(\mathcal{M}) \\ L_{X \times_S Y/S}^{(u,v)}(\mathcal{M}) &\xrightarrow{\sim} L_{X/S}^u(\mathcal{M}) \times_S L_{Y/S}^v(\mathcal{M}). \end{aligned}$$

PROOF [SGA 3], Expose II, Prop. 3.8 □

11.14 Proposition *If X/S and Y/S verify (E), then $X \times_S Y/S$ verifies (E), too, and the above isomorphisms are compatible with the module-structures.*

PROOF [SGA 3], Expose II, Prop. 3.8.bis □

⁴Let $X \rightarrow X'$ be a morphism and let $\mathcal{F}$ be a X' -functor. Recall that we denote the restriction of $\mathcal{F}$ to an X -functor by $\mathcal{F}_X$.

As Hom commutes with base change, we get

11.15 Proposition *The formation of $T_{X/S}(\mathcal{M})$ and $L_{X/S}^u(\mathcal{M})$ commutes with base change: Let $S' \rightarrow S$. Then there are isomorphisms functorial in $\mathcal{M}$:*

$$\begin{aligned} T_{X_{S'}/S'}(\mathcal{M} \otimes \mathcal{O}_{S'}) &\xrightarrow{\sim} [T_{X/S}(\mathcal{M})]_{S'} \\ L_{X_{S'}/S'}(\mathcal{M} \otimes \mathcal{O}_{S'}) &\xrightarrow{\sim} [L_{X/S}^u(\mathcal{M})]_{S'}, \quad \text{where } u' := u(s). \end{aligned}$$

PROOF [SGA 3], Expose II, Prop. 3.4 □

11.2.2. The tangent bundle as a scheme.

Let R be a ring, M a R -module. Consider the symmetric algebra $S(M) := S_R(M)$ defined by the following universal property: For all commutative R -algebras A there exist a commutative diagram:

$$\begin{array}{ccc} M & \xrightarrow{T(f)} & S(M) \\ \downarrow & \swarrow \exists! & \\ A & & \end{array}$$

i.e.:

$$\mathrm{Hom}_R(S(M), A) \cong \mathrm{Hom}_{R\text{-lin}}(M, A).$$

Let now $(X, \mathcal{O}_X)$ be a scheme and let $\mathcal{E}$ be a $\mathcal{O}_X$ -module over X . Then an open subset $U \subset X$ induces a $\Gamma(U, \mathcal{O}_X)$ -module $S(\Gamma(U, \mathcal{E}))$, and thus a presheaf of algebras

$$U \rightsquigarrow S(\Gamma(U, \mathcal{E})).$$

Let $S(\mathcal{E}) := S_{\mathcal{O}_X}(\mathcal{E})$ be the sheaf associated to this presheaf. Then $S(\mathcal{E})$ has the following universal property: For all $\mathcal{O}_X$ -algebras $\mathcal{A}$ and for all homomorphisms $\mathcal{E} \rightarrow \mathcal{A}$ $\mathcal{O}_X$ -modules there exists a commutative diagram

$$\begin{array}{ccc} \mathcal{E} & \longrightarrow & S(\mathcal{E}) \\ \downarrow & \swarrow \exists! & \\ \mathcal{A} & & \end{array}$$

S is a functor $\{\mathcal{O}_X\text{-module sheaves over } X\} \rightarrow \{\mathcal{O}_X\text{-algebras over } X\}$. It has the following properties:

- a) $S(\mathcal{E})_x = S(\mathcal{E}_x)$ for all $x \in X$;
- b) $S(\mathcal{E} \oplus \mathcal{F}) = S(\mathcal{E}) \otimes_{\mathcal{O}_X} S(\mathcal{F})$;
- c) $S_{\mathcal{O}_X}(\mathcal{O}_X) = \mathcal{O}_X[T] := \mathcal{O}_X \otimes_{\mathbb{Z}} \mathbb{Z}[T]$;

- d) $S(f^*\mathcal{F}) = f^*S(\mathcal{F})$ for all morphisms $f : (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$;
- e) $S(\tilde{M}) = \widetilde{S(M)}$ if $X = \text{Spec } A$ is affine and if $\mathcal{E} = \tilde{M}$ is associated to an A -module M .

11.16 Definition Let S be a scheme and $\mathcal{E}$ a quasi-coherent $\mathcal{O}_S$ -module. Then

$$\mathbb{V}(\mathcal{E}) := \underline{\text{Spec}} S_{\mathcal{O}_S}(\mathcal{E})$$

is called the *vector bundle* associated to $\mathcal{E}$.

Let now $f : X \rightarrow S$ be a morphism of schemes. Due to the universal property of $S_{\mathcal{O}_S}(\mathcal{E})$ there are canonical bijections:

$$\begin{aligned} \text{Hom}_S(X, \mathbb{V}(\mathcal{E})) &= \underline{\text{Hom}}_{\mathcal{O}_S}(S(\mathcal{E}), f_*\mathcal{O}_X) \\ &= \underline{\text{Hom}}_{\mathcal{O}_S-\text{lin}}(\mathcal{E}, f_*\mathcal{O}_X) \\ &= \underline{\text{Hom}}_{\mathcal{O}_X-\text{lin}}(f^*\mathcal{E}, \mathcal{O}_X), \end{aligned}$$

where the last bijection is due to Proposition 7.15. By means of these equations, we deduce the following theorem.

11.17 Theorem *The sheaf of S -sections of $\mathbb{V}(\mathcal{E})$ is in canonical bijection with the dual $\mathcal{E}^\vee := \underline{\text{Hom}}_{\mathcal{O}_S}(\mathcal{E}, \mathcal{O}_S)$ of $\mathcal{E}$.*

Especially, if $X = \{\xi\}$ is the spectrum of a field K , the structure-morphism $f : X \rightarrow S$ corresponds to a monomorphism $k(s) \hookrightarrow K$, where $s := f(\xi)$. The set of all S -morphisms $\{\xi\} \rightarrow \mathbb{V}(\mathcal{E})$ are the points of $\mathbb{V}(\mathcal{E})$ with values in the field extension K of $k(s)$ (and in particular they are elements of the fibre $\pi^{-1}(s)$, where $\pi : \mathbb{V}(\mathcal{E}) \rightarrow S$ is the structure-morphism). By the above, this set is in bijection with the set of $\mathcal{O}_X$ -module-homomorphisms $f^*\mathcal{E} \rightarrow \mathcal{O}_X = K$. Now,

$$f^*\mathcal{E} = (f^*\mathcal{E})_\xi = \mathcal{E}_s \otimes_{\mathcal{O}_{S,s}} K = \mathcal{E}_s / \mathfrak{m}_s \mathcal{E}_s \otimes_{k(s)} K.$$

Thus we deduce the following one-to-one correspondence between sets:

$$\begin{aligned} \left\{ \begin{array}{ccc} \{\xi\} & \longrightarrow & \mathbb{V}(\mathcal{E}) \\ \downarrow & & \downarrow \\ \{s\} & \hookrightarrow & S \end{array} \right\} &\xleftrightarrow{1:1} \text{Hom}_K(\mathcal{E}_s / \mathfrak{m}_s \mathcal{E}_s \otimes_{k(s)} K, K) \\ &= \text{Hom}_{k(s)}(\mathcal{E}_s / \mathfrak{m}_s \mathcal{E}_s, k(s)) \otimes_{k(s)} K \quad (\text{if } \mathcal{E}_s / \mathfrak{m}_s \mathcal{E}_s \text{ or } K \text{ is} \\ &\hspace{15em} \text{finite-dim. over } k(s)) \\ &= (\mathcal{E}_s / \mathfrak{m}_s \mathcal{E}_s)^\vee \otimes_{k(s)} K. \end{aligned}$$

11.18 Proposition *The association $\mathcal{E} \rightsquigarrow \mathbb{V}(\mathcal{E})$ is a contravariant functor*

$$\mathbb{V} : (\text{quasi-coherent } \mathcal{O}_S\text{-modules}) \rightarrow (\text{affine } S\text{-schemes}).$$

It has got the following properties:

- a) $\mathbb{V}(\mathcal{E})$ is of finite type over S if $\mathcal{E}$ is an $\mathcal{O}_S$ -module of finite type.
- b) $\mathbb{V}(\mathcal{E} \oplus \mathcal{F}) = \mathbb{V}(\mathcal{E}) \times_S \mathbb{V}(\mathcal{F})$.
- c) $\mathbb{V}(g^*\mathcal{E}) = \mathbb{V}(\mathcal{E}) \times_S S'$ for all morphisms $g : S' \rightarrow S$.
- d) $\mathbb{V}(\mathcal{F}) \hookrightarrow \mathbb{V}(\mathcal{E})$ is a closed immersion if $\mathcal{E} \rightarrow \mathcal{F}$ is a surjective morphism of quasi-coherent $\mathcal{O}_X$ -modules.

PROOF [EGA II], Prop. 1.7.11 □

Now we use this general construction in order to define the tangent bundle. We will see that sections of the tangent bundle are indeed linked with derivations and actually provide a generalization of the differential geometric notion of vector-fields.

11.19 Definition Let $f : X \rightarrow S$ be a morphism of schemes. Let $\mathcal{F}$ be a $\mathcal{O}_X$ -module. A homomorphism of sheaves of additive groups

$$\mathcal{D} : \mathcal{O}_X \rightarrow \mathcal{F}$$

is called an S -derivation of $\mathcal{O}_X$ into $\mathcal{F}$ if and only if:

- a) For all open subsets $V \subset X$ and each pair (t_1, t_2) of sections of $\mathcal{O}_X$ above V one has:

$$\mathcal{D}(t_1 \otimes t_2) = t_1 \cdot \mathcal{D}(t_2) + \mathcal{D}(t_1) \cdot t_2.$$

- b) For all open subsets $V \subset X$, each section t of $\mathcal{O}_X$ above V and each section s of $\mathcal{O}_S$ above an open subset $U \subset S$ such that $V \subset f^{-1}(U)$, one has

$$\mathcal{D}(s|_V \cdot t) = s|_V \cdot \mathcal{D}(t).$$

11.20 Remark a) In the situation above $\mathcal{D} : \mathcal{O}_X \rightarrow \mathcal{F}$ is an S -derivation if and only if for all $x \in X$ the homomorphism of additive groups $\mathcal{D}_x : \mathcal{O}_{X,x} \rightarrow \mathcal{F}_x$ is a $\mathcal{O}_{S,f(x)}$ -derivation in the ordinary sense.

- b) The S -derivations of $\mathcal{O}_X$ into $\mathcal{F}$ form a $\Gamma(X, \mathcal{O}_X)$ -module $\underline{\text{Der}}_S(\mathcal{O}_X, \mathcal{F})$.

- c) Let $\mathcal{F} = \mathcal{O}_X$. An S -derivation of $\mathcal{O}_X$ into itself is simply called an S -derivation of $\mathcal{O}_X$.

11.21 Proposition Let $f : X \rightarrow S$ be a morphism of schemes and $d_{X/S} : \mathcal{O}_X \rightarrow \Omega_{X/S}^1$ the canonical S -derivation (see our exposition following Remark 1.2). Then for all $\mathcal{O}_X$ -modules $\mathcal{F}$ there is a canonical isomorphism of $\Gamma(X, \mathcal{O}_X)$ -modules:

$$\begin{aligned} \underline{\text{Hom}}_{\mathcal{O}_X}(\Omega_{X/S}^1, \mathcal{F}) &\xrightarrow{\sim} \underline{\text{Der}}_S(\mathcal{O}_X, \mathcal{F}) \\ u &\mapsto u \circ d_{X/S} \end{aligned}$$

PROOF We may check the statement on stalks, but then the statement reduces to the defining universal property of $\Omega_{X/S,x}^1$, $x \in X$, by Remark 11.20 a). $\square$

Remark: Due to the pointwise characterization of S -derivations, the presheaf

$$U \rightsquigarrow \mathrm{Der}_S(\mathcal{O}_U, \mathcal{F}|_U)$$

is already a sheaf. This $\mathcal{O}_X$ -module

$$\underline{\mathrm{Der}}_S(\mathcal{O}_X, \mathcal{F})$$

is called the sheaf of S -derivations of $\mathcal{O}_X$ into $\mathcal{F}$. Thus we may restate the above proposition as follows:

11.22 Corollary *For all $\mathcal{O}_X$ -modules $\mathcal{F}$ the homomorphism of $\mathcal{O}_X$ -modules*

$$\begin{aligned} \underline{\mathrm{Hom}}_{\mathcal{O}_X}(\Omega_{X/S}^1, \mathcal{F}) &\longrightarrow \underline{\mathrm{Der}}_S(\mathcal{O}_X, \mathcal{F}) \\ u &\mapsto u_d := u \circ d_{X/S} \end{aligned}$$

is bijective.

11.23 Definition The dual $\mathcal{T}_{X/S}$ of the $\mathcal{O}_X$ -module $\Omega_{X/S}^1$ is called the tangent sheaf of X relative to S . By Corollary 11.22 we can write:

$$\mathcal{T}_{X/S} := \underline{\mathrm{Hom}}_{\mathcal{O}_S}(\Omega_{X/S}^1, \mathcal{O}_X) = \underline{\mathrm{Der}}_S(\mathcal{O}_X, \mathcal{O}_X).$$

Therefore we see how the tangent bundle must be defined.

11.24 Definition The tangent bundle $T_{X/S}$ of X relative to S is

$$T_{X/S} := \mathbb{V}(\Omega_{X/S}^1).$$

This is well-defined, because the sheaf of relative differential forms $\Omega_{X/S}^1$ is a quasi-coherent $\mathcal{O}_X$ -module.⁵

11.25 Remark Recalling Theorem 11.17, we see that there is a canonical bijection

$$\Gamma(T_{X/S}/X) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{O}_X}(\Omega_{X/S}^1, \mathcal{O}_X) = \Gamma(X, \mathcal{T}_{X/S}).$$

In this bijection, which is compatible with restrictions, we may replace X by an open subset $U \subset X$. Thus we conclude: There is an isomorphism between the tangent sheaf $\mathcal{T}_{X/S}$ of X relative S and the sheaf of germs of X -sections of the fibre bundle $T_{X/S}$ of X relative S . This means that vector fields are indeed given by sections of the tangent bundle as we are used to from differential geometry.

⁵It is admissible to use here the same label $T_{X/S}$ for the tangent-bundle as in Definition 11.8, because due to Proposition 11.27 both notions coincide if the functor X is representable by a scheme.

But let us finally prove that this notion of vector fields actually reduces to the classical differential geometric notion. We do so by considering the tangent space at physical points (see Definition 6.9). Consider the special case of a commutative diagram

$$\begin{array}{ccc} \{\xi\} & \hookrightarrow & T_{X/S} \\ \downarrow & & \downarrow \\ \{x\} & \hookrightarrow & X \end{array}$$

where $x \in X$ and $\xi \in T_{X/S}$ are points such that $k(x) = k(\xi)$. Let $T_{X/S}(x)$ denote the set of these diagrams; i.e. $T_{X/S}(s)$ is the set of x -valued points of $T_{X/S}$. By our exposition preceding Proposition 11.18, we already know that:

$$T_{X/S}(x) = \mathrm{Hom}_{k(x)}(\Omega_{X/S}^1 \otimes k(x), k(x))$$

On the other hand we saw in Theorem 11.2 that there is also a bijection

$$T_{X_s, x} = \mathrm{Hom}_{k(x)}(\Omega_{X/S}^1 \otimes k(x), k(x))$$

in terms of the Zariski tangent space at x , if $f : X \rightarrow S$ is smooth and if $x \in X$ is a physical point. This yields indeed $T_{X/S}(x) = T_{X_s, x}$ in physical situations.

11.26 Definition $T_{X/S}(x)$ is called the tangent-space of X at x relative to S .

Finally let us remark that this notion of the tangent bundle and of the tangent-space in terms of schemes as given above is a special case of the notion in terms of functors which was introduced in the preceding subsection 11.2.1. Both notions coincide if the considered functors are representable by schemes:

11.27 Proposition *Let X be an S -functor which is representable by an S -scheme $X \rightarrow S$. Let $T_{X/S}$ (resp. $T_{X/S}(\mathcal{M})$) denote the relative tangent bundle in the sense of Definition 11.8. Then $T_{X/S}(\mathcal{M})$ and $L_{X/S}^u(\mathcal{M})$ are representable. In particular, we can write:*

$$\begin{aligned} T_{X/S} &= \mathbb{V}(\Omega_{X/S}^1) \\ L_{X/S}^u &= \mathbb{V}(u^* \Omega_{X/S}^1). \end{aligned}$$

PROOF [SGA 3], Expose II, Prop.3.3 □

11.28 Remark Within the setting of Proposition 11.27 let u be the canonical inclusion $u : \{x\} \hookrightarrow X$ corresponding to a point $x \in X$. Then there is a bijection

$$L_{X/S}^u(\{x\}) = T_{X/S}(x).$$

In particular, it is justified to label $L_{X/S}^u$ as tangent space at u as we did in Definition 11.10.

Using the properties of fibre bundles, which are stated in Proposition 11.18, we conclude as follows.

11.29 Remark a) Let $f : X \rightarrow Y$ be an S -morphism of schemes and let $f^*\Omega_{Y/S}^1 \rightarrow \Omega_{X/S}^1$ be the induced morphism pull-back morphism (see Definition 1.3). Then, due to the identity $\mathbb{V}(f^*\Omega_{Y/S}^1) = \mathbb{V}(\Omega_{X/S}^1) \times_Y X$, the pull-back morphism gives rise to a canonical morphism induced by f :

$$T_{X/S}(f) : T_{X/S} \rightarrow T_{Y/S} \times_Y X.$$

If $g : Y \rightarrow Z$ is a second S -morphism, we get

$$T_{X/S}(g \circ f) = (T_{Y/S}(g) \times \text{id}_X) \circ T_{X/S}(f).$$

b) For all base changes $S' \rightarrow S$ we get an isomorphism:

$$T_{X'/S'} \xrightarrow{\sim} T_{X/S} \times_S S' = T_{X/S} \times_X X',$$

where $X' := X \times_S S'$. If $x' \in X'$ lies over $x \in X$, we get

$$T_{X'/S'}(x') = T_{X/S}(x) \otimes_{k(x)} k(x').$$

c) Let $f : X \rightarrow Y$ be an S -morphism, let $x \in X$ and $y := f(x)$. Then

$$f^*\Omega_{Y/S}^1 \otimes_{\mathcal{O}_{X,x}} k(x) = \Omega_{Y/S}^1 \otimes_{\mathcal{O}_{Y,y}} \mathcal{O}_{X,x} \otimes_{\mathcal{O}_{X,x}} k(x) = (\Omega_{Y/S}^1 \otimes_{\mathcal{O}_{Y,y}} k(y)) \otimes_{k(y)} k(x).$$

If $\Omega_{Y/S}^1$ is a $\mathcal{O}_Y$ -module of finite type (which will be the case in the situations we are going to consider), we have an isomorphism

$$\text{Hom}_{k(x)}(f^*\Omega_{Y/S}^1 \otimes k(x), k(x)) = T_{Y/S}(y) \otimes_{k(y)} k(x),$$

because $k(y) \rightarrow k(x)$ is flat. Thus the pull-back of differential forms $f^*\Omega_{Y/S}^1 \rightarrow \Omega_{X/S}^1$ gives rise to a homomorphism of $k(x)$ -vector spaces

$$T_{f,x} : T_{X/S}(x) \rightarrow T_{Y/S}(y) \otimes_{k(y)} k(x),$$

called the *tangent map* of f at x . At physical points this definition coincides with the former Definition 11.1 b).

As the sheaf Hom occurs in a canonical way if we consider vector bundles (like the relative tangent-bundle), we finish with a collection of some properties of Hom which might be useful.

11.30 Proposition *Let $(X, \mathcal{O}_X)$ be a ringed space and let $\mathcal{E}$ be a locally free $\mathcal{O}_X$ -module of finite rank. We define the dual of $\mathcal{E}$, denoted $\mathcal{E}^\vee$, to be the sheaf of Hom $_{\mathcal{O}_X}(\mathcal{E}, \mathcal{O}_X)$. Then the following assertions are true:*

a) $(\mathcal{E}^\vee)^\vee \cong \mathcal{E}$.

b) For any $\mathcal{O}_X$ -module $\mathcal{F}$

$$\underline{\mathrm{Hom}}_{\mathcal{O}_X}(\mathcal{E}, \mathcal{F}) \cong \mathcal{E}^\vee \otimes \mathcal{F}.$$

c) For any $\mathcal{O}_X$ -modules $\mathcal{F}, \mathcal{G}$

$$\mathrm{Hom}_{\mathcal{O}_X}(\mathcal{E} \otimes \mathcal{F}, \mathcal{G}) \cong \mathrm{Hom}_{\mathcal{O}_X}(\mathcal{F}, \underline{\mathrm{Hom}}_{\mathcal{O}_X}(\mathcal{E}, \mathcal{G})).$$

d) (*Projection Formula*): If $f : (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$ is a morphism of ringed spaces, if $\mathcal{F}$ is an $\mathcal{O}_X$ -module, and if $\mathcal{E}$ is a locally free $\mathcal{O}_Y$ -module of finite rank, then there is a natural isomorphism

$$f_*(\mathcal{F} \otimes_{\mathcal{O}_X} f^*\mathcal{E}) \cong f_*\mathcal{F} \otimes_{\mathcal{O}_Y} \mathcal{E}.$$

e) Let X be a noetherian scheme, let $\mathcal{F}$ be a coherent sheaf on X , let $\mathcal{G}$ be any $\mathcal{O}_X$ -module and let $x \in X$ be a point. Then we have

$$\underline{\mathrm{Ext}}^i(\mathcal{F}, \mathcal{G})_x \cong \mathrm{Ext}_{\mathcal{O}_{X,x}}^i(\mathcal{F}_x, \mathcal{G}_x)$$

for all $i \geq 0$, where the right-hand side is Ext over the local ring $\mathcal{O}_{X,x}$.

PROOF [Har], Chap. II, Ex. 5.1 and [Har], Chap. III, Prop. 6.8 □

11.31 Proposition Let $X = \mathrm{Spec} A$ an affine scheme, and let $\mathcal{F}, \mathcal{G}$ be $\mathcal{O}_X$ -modules. Then the canonical map

$$\mathrm{Hom}_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G}) \rightarrow \mathrm{Hom}_A(\mathcal{F}(X), \mathcal{G}(X))$$

is bijective if $\mathcal{F}$ is quasi-coherent.

PROOF [Liu], Ex. 5.1.5 □

11.32 Proposition Let X be a scheme and let $\mathcal{E}$ be a locally free $\mathcal{O}_X$ -module. Then the sheaf $\underline{\mathrm{Hom}}$ commutes with pull-back, i.e.:

$$\alpha^*\mathcal{E}^\vee \cong (\alpha^*\mathcal{E})^\vee$$

for all morphisms $\alpha : Y \rightarrow X$. Especially, if $p : X \rightarrow S$ is a smooth S -scheme, if $Y = S$, if $\alpha : S \hookrightarrow X$ is a section of p and if $\mathcal{E} = \Omega_{X/S}^1$, this isomorphism is canonical.

PROOF If α is flat, one knows that the canonical morphism

$$\alpha^*\mathcal{E}^\vee \xrightarrow{\sim} (\alpha^*\mathcal{E})^\vee, \quad \varphi \otimes a \mapsto a \cdot \tilde{\varphi},$$

$\tilde{\varphi}(e \otimes a) := \varphi(e) \otimes a$, is an isomorphism (see e.g. [Liu], Ex. 1.2.8). Also if p is not flat, one can construct an explicit inverse of the canonical morphism above. In our physical situation $p : X \rightarrow S$ will be a smooth S -group scheme (see Definition 11.46), $\mathcal{E}$ will be the module of differential forms $\Omega_{X/S}^1$, and $\alpha : S \hookrightarrow X$ will be a section of p . So we will only perform the proof in this case, because then no explicit construction is necessary. We may simply apply the general theory: Recalling that $\alpha = \tau_\alpha \circ \varepsilon$, where ε is the unit section and where τ_α is the isomorphism given by left-translation with α (see subsection 11.4.4), and evoking Proposition 11.58, we get canonical isomorphisms

$$p^*(\alpha^* \Omega_{X/S}^1)^\vee \cong p^*(\varepsilon^* \tau_\alpha^* \Omega_{X/S}^1)^\vee \cong p^*(\varepsilon^* \Omega_{X/S}^1)^\vee \cong (p^* \varepsilon^* \Omega_{X/S}^1)^\vee \cong (\Omega_{X/S}^1)^\vee,$$

because p is flat due to smoothness. Applying the pull-back functor α^* and using the fact that $\alpha^* p^* = (p \circ \alpha)^* = \text{id}^*$ we are done. $\square$

11.3. Differential p -forms and the exterior differential

We already introduced the sheaf $\Omega_{X/S}^1$ of differential 1-forms in the beginning of section 1.1. In the previous section 11.2 on the tangent-bundle, we recognized differential forms as sections of the tangent-bundle: If we denote by $\mathcal{T}_{X/S} := \left(\Omega_{X/S}^1\right)^\vee := \underline{\text{Hom}}_{\mathcal{O}_X}(\Omega_{X/S}^1, \mathcal{O}_X)$ the dual of $\Omega_{X/S}^1$ (i.e. the sheaf of vector fields), then in accordance with Definition 11.24 we define the tangent-bundle $T_{X/S}$ and the cotangent-bundle $T_{X/S}^*$ as follows:

$$T_{X/S} := \mathbb{V}(\Omega_{X/S}^1) \quad T_{X/S}^* := \mathbb{V}(\mathcal{T}_{X/S}).$$

Vector fields and differential forms may be identified with sections of the respective bundles (see Theorem 11.17):

$$\mathcal{T}_{X/S}(X) = \Gamma(T_{X/S}/X) \quad \Omega_{X/S}^1(X) = \Gamma(T_{X/S}^*/X)$$

As already exposed in section 4.3, a global section ω of some tensorpower $\Omega_{X/S}^{\otimes n}$ of $\Omega_{X/S}^1$ may be interpreted as a “multilinear” morphism

$$\omega : T_{X/S} \times_X \dots \times_X T_{X/S} \rightarrow \mathbb{A}_X^1.$$

This morphism is multilinear in two regards:

a) pointwise multilinearity: Let $S = \text{Spec } R$ be an affine scheme, and let X be a smooth and separated S -scheme. Consider an S -valued point $\alpha : S \rightarrow X$ of X . Physically we interpret $\alpha \in X(S)$ as an “adelic” point of space-time. First pulling back ω via α (i.e. performing a base change of ω with α) we get a morphism $\alpha^* \omega$. Recalling the identity $T_{X/S} \times_X S = \mathbb{V}(\alpha^* \Omega_{X/S}^1)$ (Proposition 11.18 c)), then applying the global section functor $\Gamma(\cdot, S)$ to $\alpha^* \omega$, using the universal property $\text{Hom}_X(X', Y \times_X Z) \cong \text{Hom}_X(X', Y) \times \text{Hom}_X(X', Z)$ of fibre products and finally evoking Theorem 11.17

in order to provide the identity $\Gamma(\mathbb{V}(\alpha^*\Omega_{X/S}^1)/S) = (\alpha^*\Omega_{X/S}^1)^\vee(S)$, we arrive at a morphism

$$\omega(\alpha) : (\alpha^*\Omega_{X/S}^1)^\vee(S) \times \dots \times (\alpha^*\Omega_{X/S}^1)^\vee(S) \longrightarrow R.$$

This morphism is R -multilinear. Due to Proposition 11.32 and Proposition 11.31 there are isomorphisms

$$(\alpha^*\mathcal{T}_{X/S})(S) = (\alpha^*\Omega_{X/S}^1)^\vee(S) = \mathrm{Hom}_R((\alpha^*\Omega_{X/S}^1)(S), R)$$

where $(\alpha^*\Omega_{X/S}^1)(S)$ is the cotangent-space, and where $(\alpha^*\mathcal{T}_{X/S})(S)$ is the tangent-space at the “adelic” space-time point α . Thus $\omega(\alpha)$ (i.e. the evaluation of the differential form ω at the “adelic” space-time point α) behaves as one would expect: It is a multilinear form which maps a tuple of tangent-vectors at α to numbers R .

b) global multilinearity: ω does not only induce a multilinear form pointwise, but also on the global sections of the tangent-bundle: In order to see this apply the global section functor $\Gamma(\cdot, X)$ to ω . Recalling the bijection $\mathrm{Hom}_Y(Y, X \times_S Y) = \mathrm{Hom}_S(Y, X)$ for S -schemes X and Y , we obtain in particular $\mathbb{A}_X^1(X) := \mathrm{Hom}_X(X, \mathbb{A}_X^1) = \mathrm{Hom}_{\mathrm{Spec} \mathbb{Z}}(X, \mathbb{A}_{\mathrm{Spec} \mathbb{Z}}^1)$. The latter set equals $\mathrm{Hom}_{\mathbb{Z}}(\mathbb{Z}[T], \mathcal{O}_X(X)) = \mathcal{O}_X(X)$ due to the following lemma.

11.33 Lemma *Let Y be an affine scheme. For any scheme X , the canonical map*

$$\mathrm{Hom}(X, Y) \longrightarrow \mathrm{Hom}(\mathcal{O}_X(Y), \mathcal{O}_X(X))$$

is bijective.

PROOF [Liu], Prop. 2.3.25 □

Finally, making once again use of the universal property $\mathrm{Hom}_X(X', Y \times_X Z) \cong \mathrm{Hom}_X(X', Y) \times \mathrm{Hom}_X(X', Z)$ of fibre products and recalling the identification $\mathcal{T}_{X/S}(X) = \Gamma(\mathcal{T}_{X/S}/X) = \mathrm{Hom}_X(X, \mathcal{T}_{X/S})$ of vector fields with global sections of the tangent-bundle, we see that ω gives rise to a $\mathcal{O}_X(X)$ -multilinear map:

$$\omega : \mathcal{T}_{X/S}(X) \times \dots \times \mathcal{T}_{X/S}(X) \longrightarrow \mathcal{O}_X(X)$$

All in all, ω indeed maps a tuple of vector fields to functions and is linear in each component.

Thus we may think of differential forms in completely the same way as we are used to from differential geometry. Using this interpretation of differential forms as multilinear forms on vector fields, we may state the following definitions in an elegant way.

11.34 Definition a) Let $f : X \rightarrow S$ be a morphism of schemes. The sheaf $\Omega_{X/S}^p$ of *differential p -forms of X relative S* is the p -th exterior product of the $\mathcal{O}_X$ -module $\Omega_{X/S}^1$:

$$\Omega_{X/S}^p := \bigwedge_{i=1}^p (\Omega_{X/S}^1).$$

b) The *exterior product* is a map $\wedge : \Omega_{X/S}^k \oplus \Omega_{X/S}^l \rightarrow \Omega_{X/S}^{k+l}$ defined as follows:

$$(\alpha \wedge \beta)(\mathbf{v}_1, \dots, \mathbf{v}_{k+l}) = \frac{1}{k!l!} \sum_{\pi \in \mathfrak{S}_{k+l}} \sigma_\pi \alpha(\mathbf{v}_{\pi(1)}, \dots, \mathbf{v}_{\pi(k)}) \beta(\mathbf{v}_{\pi(k+1)}, \dots, \mathbf{v}_{\pi(k+l)}) .$$

Thereby $\mathfrak{S}_n$ denotes the group of permutations of the numbers $1, \dots, n$, and σ_π is the sign of the permutation, i.e. the number of transpositions which transfer $1, 2, \dots, n$ into $\pi(1), \pi(2), \dots, \pi(n)$.

The exterior product has the following properties:

- $(f_1\alpha_1 + f_2\alpha_2) \wedge \beta = f_1\alpha_1 \wedge \beta + f_2\alpha_2 \wedge \beta, \quad f_1, f_2 \in \mathcal{O}_X$
- $\alpha \wedge (\beta \wedge \gamma) = (\alpha \wedge \beta) \wedge \gamma$
- $\alpha \wedge \beta = (-1)^{kl} \beta \wedge \alpha, \quad \alpha \in \Omega_{X/S}^k, \beta \in \Omega_{X/S}^l.$

11.35 Remark From these properties we conclude that there is a graded algebra of differential forms in algebraic geometry, too. More precisely, we may state:

- a) $\Omega_{X/S}^0 = \mathcal{O}_X$ and $\Omega_{X/S}^p = 0$ for $p < 0$.
- b) The $\Omega_{X/S}^p$ are the homogeneous components of (the graded) exterior algebra of $\Omega_{X/S}^1$:

$$\Omega_{X/S}^\bullet := \bigwedge (\Omega_{X/S}^1) =: \bigoplus_{p \in \mathbb{Z}} \Omega_{X/S}^p.$$

- c) $\Omega_{X/S}^\bullet$ is a graded, quasi-coherent, anti-commutative $\mathcal{O}_X$ -algebra via the exterior product. For open affine $U \subset X$ one has

$$\Gamma(U, \Omega_{X/S}^\bullet) = \bigwedge \Gamma(U, \Omega_{X/S}^1)$$

where $\Gamma(U, \Omega_{X/S}^1)$ is considered as a $\Gamma(U, \mathcal{O}_X)$ -module.

- d) $\Omega_{X/S}^p = \left(\Omega_{B/R}^p \right)^\sim := \left(\bigwedge_{i=1}^p \Omega_{B/R}^1 \right)^\sim$ if $X = \operatorname{Spec} B$ and $S = \operatorname{Spec} R$ are affine.

11.36 Definition The *interior product* between a vector field $\mathbf{v} \in \mathcal{T}_{X/S}$ and a differential k -Form $\alpha \in \Omega_{X/S}^k$, $k \geq 1$, is a differential $(k-1)$ -form $i_{\mathbf{v}}\alpha \in \Omega_{X/S}^{k-1}$ defined as follows:

$$(i_{\mathbf{v}}\alpha)(\mathbf{v}_1, \dots, \mathbf{v}_{k-1}) := \alpha(\mathbf{v}, \mathbf{v}_1, \dots, \mathbf{v}_{k-1}) ,$$

where $\mathbf{v}_i \in \mathcal{T}_{X/S}$.

Applying the above construction of the exterior product to the sheaf $\mathcal{T}_{X/S}$ of vector fields instead of the sheaf $\Omega_{X/S}^1$ of co-vector fields, we may consider the exterior product

$$\mathcal{T}_{X/S}^k := \bigwedge_{i=1}^k \mathcal{T}_{X/S}.$$

In particular $\mathcal{T}_{X/S}^0 = \mathcal{O}_X$, $\mathcal{T}_{X/S}^1 = \mathcal{T}_{X/S}$ and $\mathcal{T}_{X/S}^k = 0$ if $k < 0$.

11.37 Definition A k -vector field is a global section of $\mathcal{T}_{X/S}^k$.

Then we may define a $\mathcal{O}_X$ -bilinear morphism $\langle \cdot, \cdot \rangle : \mathcal{T}_{X/S}^k \oplus \Omega_{X/S}^k \longrightarrow \mathcal{O}_X$ by means of $\langle \mathbf{v}_1 \wedge \cdots \wedge \mathbf{v}_k, \alpha \rangle := k! \alpha(\mathbf{v}_1, \dots, \mathbf{v}_k)$.

11.38 Definition The *rejuvenation* of a differential $(k+l)$ -form $\omega \in \Omega_{X/S}^{k+l}$ with a k -vector field $\mathbf{v} \in \mathcal{T}_{X/S}^k$ is the $\mathcal{O}_X$ -bilinear morphism

$$\rfloor : \mathcal{T}_{X/S}^k \oplus \Omega_{X/S}^{k+l} \longrightarrow \Omega_{X/S}^l, \quad (\mathbf{v}, \omega) \mapsto \mathbf{v} \rfloor \omega$$

defined as follows: $(\mathbf{v} \rfloor \omega)(\mathbf{v}_1, \dots, \mathbf{v}_l) := \langle \mathbf{v} \wedge \mathbf{v}_1 \wedge \dots \wedge \mathbf{v}_l, \omega \rangle$.

The *rejuvenation* has the following properties:

- $(f_1 X_1 + f_2 X_2) \rfloor \alpha = f_1 (X_1 \rfloor \alpha) + f_2 (X_2 \rfloor \alpha)$,
- $X \rfloor (f_1 \alpha_1 + f_2 \alpha_2) = f_1 (X \rfloor \alpha_1) + f_2 (X \rfloor \alpha_2)$.

A further very important operation is the *exterior differential*.

11.39 Theorem There exists a unique endomorphism $d : \Omega_{X/S}^\bullet \rightarrow \Omega_{X/S}^\bullet$ of sheaves of additive groups, the so called exterior differential, such that:

(i) $d \circ d = 0$.

(ii) For all open $U \subset X$ and for all $f \in \Gamma(U, \mathcal{O}_X)$ one has

$$df = d_{X/S} f.$$

(iii) For all open $U \subset X$, all integers $p, q \in \mathbb{Z}$ and for all pairs of sections $\omega'_p \in \Gamma(U, \Omega_{X/S}^p)$, $\omega''_q \in \Gamma(U, \Omega_{X/S}^q)$ we have:

$$d(\omega'_p \wedge \omega''_q) = (d\omega'_p) \wedge \omega''_q + (-1)^p \omega'_p \wedge (d\omega''_q).$$

PROOF [EGA IV₄], Thm. 16.6.2 □

Especially d is local, because it is a morphism of sheaves; i.e. if $\alpha|_U = \beta|_U$ for some sections α, β of $\mathcal{O}_X$ and an open subset $U \subset X$, then

$$(d\alpha)|_U = d(\alpha|_U) = d(\beta|_U) = (d\beta)|_U.$$

11.40 Corollary Let $\omega \in \Omega_{X/S}^k$ and $\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_k \in \mathcal{T}_{X/S}$. Then the exterior differential has the following form:

$$(d\omega)(\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_k) := \sum_{i=0}^k (-1)^i \mathbf{v}_i(\omega(\mathbf{v}_0, \dots, \check{\mathbf{v}}_i, \dots, \mathbf{v}_k)) \\ + \sum_{0 \leq i < j \leq k} (-1)^{i+j} \omega([\mathbf{v}_i, \mathbf{v}_j], \mathbf{v}_0, \dots, \check{\mathbf{v}}_i, \dots, \check{\mathbf{v}}_j, \dots, \mathbf{v}_k),$$

where $\check{\mathbf{v}}_i$ means that this vector is omitted.⁶ The commutator $[\cdot, \cdot]$ of vector fields is defined in Definition 11.41.

PROOF One has to check that the three conditions of Theorem 11.39 are fulfilled. In order to show (ii) let $f \in \mathcal{O}_X$ and $\mathbf{v} \in \mathcal{T}_{X/S}$. Then by definition $(df)(\mathbf{v}) := \mathbf{v}(f)$, where on the right hand side $\mathbf{v}$ is considered as a derivation by means of Corollary 11.22. Thus $(df)(\mathbf{v}) = \mathbf{v}(d_{X/S}f)$ if $\mathbf{v}$ is now considered as a dual differential form. Finally the canonical isomorphism $\mathcal{E} \rightarrow (\mathcal{E}^\vee)^\vee$ provides an equality $\mathbf{v}(d_{X/S}f) = (d_{X/S}f)(\mathbf{v})$. This proves (ii), because $\mathbf{v}$ is arbitrary. A longer algebraic manipulation (which is known from differential geometry) shows that d fulfills the conditions (i) and (iii) of Definition 11.36, too. $\square$

11.41 Definition Let $X \rightarrow S$ be an S -scheme, let $\mathbf{v}, \mathbf{w} \in \Gamma(\mathcal{T}_{X/S}/X)$ be vector fields, and let $f \in \mathcal{O}_X$. Let us introduce the following abbreviation $\mathbf{v}_d := \mathbf{v} \circ d$. From the algebraic properties of the derivations $\mathbf{v}_d$ and $\mathbf{w}_d$ it follows that

$$[\mathbf{v}, \mathbf{w}](f) := (\mathbf{v}_d)((\mathbf{w}_d)(f)) - (\mathbf{w}_d)((\mathbf{v}_d)(f))$$

is a derivation, too.⁷ The corresponding vector field $[\mathbf{v}, \mathbf{w}] \in \Gamma(\mathcal{T}_{X/S}/X)$ is called the *commutator of $\mathbf{v}$ and $\mathbf{w}$* .

Let us conclude with the definition of the Hodge-star operator. For this purpose recall that the metric yields an isomorphism $g : \mathcal{T}_{X/S} \rightarrow (\mathcal{T}_{X/S})^\vee = \Omega_{X/S}^1$ (Definition 4.20). In particular we get isomorphisms

$$\Lambda^k g^{-1} : \Omega_{X/S}^k \xrightarrow{\sim} \mathcal{T}_{X/S}^k$$

for all $k \in \mathbb{N}$. By Theorem 11.61 there exists a volume form v_g in physical situations.

⁶ In order to give sense to the expression $\mathbf{v}_i(\omega(\mathbf{v}_0, \dots, \check{\mathbf{v}}_i, \dots, \mathbf{v}_k))$, let us point out that in this situation $\mathbf{v}_i$ is considered as an S -derivation of $\mathcal{O}_X$ by means of Corollary 11.22.

⁷ $[\mathbf{v}, \mathbf{w}](fg) = (\mathbf{v}_d)((\mathbf{w}_d)(fg)) - (\mathbf{w}_d)((\mathbf{v}_d)(fg))$
 $= (\mathbf{v}_d)((\mathbf{w}_d)(f) \cdot g + f \cdot (\mathbf{w}_d)(g)) - (\mathbf{w}_d)((\mathbf{v}_d)(f) \cdot g + f \cdot (\mathbf{v}_d)(g))$
 $= (\mathbf{v}_d)((\mathbf{w}_d)(f)) \cdot g + (\mathbf{w}_d)(f) \cdot (\mathbf{v}_d)(g) + (\mathbf{v}_d)(f)(\mathbf{w}_d)(g) + f \cdot (\mathbf{v}_d)((\mathbf{w}_d)(g))$
 $\quad - (\mathbf{w}_d)((\mathbf{v}_d)(f)) \cdot g - (\mathbf{v}_d)(f) \cdot (\mathbf{w}_d)(g) - (\mathbf{w}_d)(f)(\mathbf{v}_d)(g) - f \cdot (\mathbf{w}_d)((\mathbf{v}_d)(g))$
 $= [\mathbf{v}, \mathbf{w}](f) \cdot g + f \cdot [\mathbf{v}, \mathbf{w}](g) \quad \text{for all } f, g \in \mathcal{O}_X.$

11.42 Definition Let $X \rightarrow S$ be of relative dimension n and assume that there exists a volume form v_g on X . Let $\alpha \in \Omega_{X/S}^k$ be a differential k -form. Then the operator $*$: $\Omega_{X/S}^k \rightarrow \Omega_{X/S}^{n-k}$,

$$* \alpha := (\Lambda^k g^{-1})(\alpha) \lrcorner v_g$$

is called the *Hodge-star operator*.

11.4. Group Schemes

11.4.1. Definition of group schemes

Let $\mathfrak{C}$ be a category; for example, let $\mathfrak{C}$ be the category (Sch/S) of schemes over a fixed scheme S . Each object $X \in \mathfrak{C}$ gives rise to its functor of points

$$h_X : \mathfrak{C} \rightarrow (Sets)$$

which associates to any $T \in \mathfrak{C}$ the set

$$h_X(T) := X(T) := \text{Hom}(T, X)$$

of T -valued points of X . Each morphism $X \rightarrow X'$ in $\mathfrak{C}$ induces a morphism $h_X \rightarrow h_{X'}$ of functors by the composition of morphisms in $\mathfrak{C}$. In this way one gets a covariant functor

$$f : \mathfrak{C} \rightarrow \text{Hom}(\mathfrak{C}^0, (Sets))$$

of $\mathfrak{C}$ to the category of covariant functors from $\mathfrak{C}^0$ (the dual of $\mathfrak{C}$) to the category of sets; the category $\text{Hom}(\mathfrak{C}^0, (Sets))$ is denoted by $\hat{\mathfrak{C}}$; it is called the category of contravariant functors from $\mathfrak{C}$ to $(Sets)$.

11.43 Proposition *The functor $h : \mathfrak{C} \rightarrow \hat{\mathfrak{C}}$ is fully faithful; i.e., for any two objects $X, X' \in \mathfrak{C}$, the canonical map*

$$\text{Hom}_{\mathfrak{C}}(X, X') \rightarrow \text{Hom}_{\hat{\mathfrak{C}}}(h_X, h_{X'})$$

is bijective. More generally, for all objects $X \in \mathfrak{C}$ and $\mathcal{F} \in \hat{\mathfrak{C}}$, there is a canonical bijection

$$\mathcal{F}(X) \xrightarrow{\sim} \text{Hom}_{\hat{\mathfrak{C}}}(h_X, \mathcal{F})$$

mapping $u \in \mathcal{F}(X)$ to the morphism $h_X \rightarrow \mathcal{F}$ which to a T -valued point $g \in h_X(T)$, where T is an object of $\mathfrak{C}$, associates the element $\mathcal{F}(g)(u) \in \mathcal{F}(T)$. The bijection coincides with the above one if $\mathcal{F} = h_{X'}$ and is functorial in X and $\mathcal{F}$ in the sense that $\mathcal{F} \mapsto \text{Hom}_{\hat{\mathfrak{C}}}(h(\cdot), \mathcal{F})$ defines an isomorphism $\hat{\mathfrak{C}} \rightarrow \hat{\mathfrak{C}}$.

In particular, if a functor $\mathcal{F} \in \text{Hom}(\mathfrak{C}^0, (\text{Sets}))$ is isomorphic to a functor h_X , then X is uniquely determined by $\mathcal{F}$ up to an isomorphism in the category $\mathfrak{C}$. In this case, the functor f is said to be *representable*. Thus Proposition 11.43 says that the functor h defines an equivalence between the category $\mathfrak{C}$ and the full subcategory of $\text{Hom}(\mathfrak{C}^0, (\text{Sets}))$ consisting of all representable functors.

In order to define group objects in the category $\mathfrak{C}$, it is necessary to introduce the notion of a law of composition on an object X of $\mathfrak{C}$. By the latter we mean a functorial morphism

$$\gamma : h_X \times h_X \rightarrow h_X.$$

Thus, a law of composition on X consists of a collection of maps

$$\gamma_T : h_X(T) \times h_X(T) \rightarrow h_X(T)$$

(laws of composition on the sets of T -valued points of X) where T varies over the objects in $\mathfrak{C}$. The functoriality of γ means that all maps γ_T are compatible with canonical maps between points of X , i.e., for any morphism $u : T' \rightarrow T$ in $\mathfrak{C}$, the diagram

$$\begin{array}{ccc} h_X(T) \times h_X(T) & \xrightarrow{\gamma_T} & h_X(T) \\ h_X(u) \times h_X(u) \downarrow & & \downarrow h_X(u) \\ h_X(T') \times h_X(T') & \xrightarrow{\gamma_{T'}} & h_X(T') \end{array}$$

is commutative. If the law of composition has the property that $h_X(T)$ is a group under γ_T for all T , then γ defines on h_X the structure of a group functor, i.e., of a contravariant functor from C to the category of groups. In this case, γ is called a group law of X .

11.44 Definition A group object in $\mathfrak{C}$ is an object X together with a law of composition $\gamma : h_X \times h_X \rightarrow h_X$ which is a group law.

It follows that a group object in $\mathfrak{C}$ is equivalent to a group functor which, as a functor to the category of sets, is representable.

When dealing with group objects, it is convenient to know that the category in question contains direct products and a final object, say S . The latter means that, for each object T of $\mathfrak{C}$, there is a unique morphism $T \rightarrow S$. So, in the following, assume that $\mathfrak{C}$ is of this type, and consider a groups object X of $\mathfrak{C}$ with group law γ . Then, since the product $X \times X$ exists in $\mathfrak{C}$ and since the functor $h : \mathfrak{C} \rightarrow \text{Hom}(\mathfrak{C}^0, (\text{Sets}))$ commutes with direct products, the law of composition $\gamma : h_X \times h_X \rightarrow h_X$ corresponds to a morphism $m : X \times X \rightarrow X$, as is seen by using Proposition 11.43. Furthermore, the injection of the unit element into each group $h_X(T)$ yields a natural transformation from h_S to h_X , hence it corresponds to a morphism

$$\varepsilon : S \rightarrow X,$$

called the *unit section of X* which is a section of the unique morphism $X \rightarrow S$. Finally, the formation of the inverse in each $h_X(T)$ defines a natural transformation $h_X \rightarrow h_X$ and hence a morphism

$$\iota : X \rightarrow X,$$

called the *inverse map on X* . The group axioms which are satisfied by the groups $h_X(T)$, and hence by the functor h_X , correspond to certain properties of the maps m, ε and ι . Namely, the following diagrams are commutative:

a) *associativity*

$$\begin{array}{ccc} X \times X \times X & \xrightarrow{m \times id_X} & X \times X \\ id_X \times m \downarrow & & \downarrow m \\ X \times X & \xrightarrow{m} & X \end{array}$$

b) *existence of a left-identity*

$$\begin{array}{ccccc} X & \xrightarrow{(p, id_X)} & S \times X & \xrightarrow{\varepsilon \times id_X} & X \times X \\ & \searrow id_X & & & \downarrow m \\ & & & & X \end{array}$$

where $p : X \rightarrow S$ is the morphism from X to the final object S .

c) *existence of a left-inverse*

$$\begin{array}{ccc} X & \xrightarrow{(\iota, id_X)} & X \times X \\ p \downarrow & & \downarrow m \\ S & \xrightarrow{\varepsilon} & X \end{array}$$

d) *commutativity* (only if all groups $h_X(T)$ are commutative)

$$\begin{array}{ccc} X \times X & \xrightarrow{\tau} & X \times X \\ & \searrow m & \downarrow m \\ & & X \end{array}$$

where τ commutes the factors.

Note that a left-identity is also a right-identity and that a left-inverse is also a right-inverse. It is clear that once we have an object X and morphisms m, ε and ι with the above properties, we can construct a group object in the given category from these data, and furthermore that group objects in $\mathfrak{C}$ and data $(X, m, \varepsilon, \iota)$ correspond bijectively to each other.

11.45 Proposition *The group objects in a category $\mathfrak{C}$ correspond one-to-one to data $(X, m, \varepsilon, \iota)$ where X is an object of $\mathfrak{C}$ and where*

$$m : X \times X \rightarrow X, \quad \varepsilon : S \rightarrow X, \quad \iota : X \rightarrow X$$

are morphisms in $\mathfrak{C}$ such that the diagrams a), b), c) above are commutative. Furthermore, a group object in $\mathfrak{C}$ is commutative if and only if, in addition, the corresponding diagram d) is commutative.

In the following we restrict ourselves to the category (Sch/S) of S -schemes where S is a fixed base scheme. Then the direct product in (Sch/S) is given by the fibred product of schemes over S , and the S -scheme S is a final object in (Sch/S) .

11.46 Definition An S -group scheme is a group object in the category of S -schemes (Sch/S) .

Due to Proposition 11.45, an S -group scheme G can be viewed as an S -scheme X together with appropriate morphisms m , ε and ι . When no confusion about the group structure is possible, we will not mention these morphisms explicitly. In particular, in our notation we will make no difference between the group object G and the associated representing scheme X . Also we want to point out that there exist group functors on (Sch/S) which are not representable and thus do not correspond to S -group schemes.

It follows immediately from Definition 11.46 that the technique of base change can be applied to group schemes. Thus, for any base change $S' \rightarrow S$, one obtains from an S -group scheme G an S' -group scheme $G_{S'} := G \times_S S'$.

Let us look at some examples of S -group schemes. We start with the classical groups $\mathbb{G}_a$ (the additive group), $\mathbb{G}_m$ (the multiplicative group), GL_n (the general linear group), and PGL_n (the projective general linear group). In terms of group functors, these groups are defined as follows. For any S -scheme T set

$$\begin{aligned} \mathbb{G}_a(T) &:= \text{the additive group } \mathcal{O}_T(T) \\ \mathbb{G}_m(T) &:= \text{the group of units in } \mathcal{O}_T(T) \\ GL_n(T) &:= \text{the group of } \mathcal{O}_T(T)\text{-linear automorphisms of } (\mathcal{O}_T(T))^n \\ PGL_n(T) &:= \text{Aut}_T(\mathbb{P}(\mathcal{O}_T^n)). \end{aligned}$$

All these group functors are representable by affine schemes over $\mathbb{Z}$. Working over $S := \text{Spec } \mathbb{Z}$, the additive group is represented by the scheme $X := \text{Spec } \mathbb{Z}[\zeta]$ (ζ is an indeterminate), where the group law $m : X \times X \rightarrow X$ corresponds to the algebra homomorphism

$$\mathbb{Z}[\zeta] \rightarrow \mathbb{Z}[\zeta] \otimes_{\mathbb{Z}} \mathbb{Z}[\zeta], \quad \zeta \mapsto \zeta \otimes 1 + 1 \otimes \zeta.$$

Similarly, for $\mathbb{G}_m$, the representing object is $\text{Spec } \mathbb{Z}[\zeta, \zeta^{-1}]$ with the group law given by $\zeta \mapsto \zeta \otimes \zeta$.

11.4.2. The Lie-algebra as a functor

Let G be a group functor over S and denote by $\varepsilon : S \rightarrow G$ the unit section. Recall from Definition 11.10 that the $\mathcal{O}_S$ -module $L_{G/S}^\varepsilon(\mathcal{M})$ denotes the tangent space of G over S at ε . Then we introduce the following notation:

11.47 Definition The $\mathcal{O}_S$ -module $\mathrm{Lie}(G/S, \mathcal{M}) := L_{G/S}^\varepsilon(\mathcal{M})$ is called the Lie-algebra of G relative $\mathcal{M}$. In particular,

$$\mathfrak{g} := \mathrm{Lie}(G/S) := \mathrm{Lie}(G/S, \mathcal{O}_S)$$

is called the *Lie-algebra* of G .

By Proposition 11.13, $T_{G/S}(\mathcal{M})$ and $\mathrm{Lie}(G/S, \mathcal{M})$ carry a group structure over S which is induced by the group structure on G . Furthermore, by definition of $L_{G/S}^\varepsilon(\mathcal{M})$, there is a cartesian diagram

$$\begin{array}{ccc} \mathrm{Lie}(G/S, \mathcal{M}) & \xrightarrow{i} & T_{G/S}(\mathcal{M}) \\ \downarrow & & \downarrow p \\ S & \xrightarrow{\varepsilon} & G \end{array}$$

and we recognize $\mathrm{Lie}(G/S, \mathcal{M})$ as the kernel of the canonical projection p . In particular, we get morphisms of groups

$$\mathrm{Lie}(G/S, \mathcal{M}) \xrightarrow{i} T_{G/S}(\mathcal{M}) \begin{array}{c} \xrightarrow{p} \\ \xleftarrow{s} \end{array} G,$$

where s is a section of p . Due to the following Proposition 11.48, $T_{G/S}(\mathcal{M})$ is the semi-direct product of G and $\mathrm{Lie}(G/S, \mathcal{M})$.

11.48 Proposition Let $\mathfrak{C} := (\mathrm{Sch}/S)$ be the category of schemes over S . Let $f : W \rightarrow G$ be a morphism of $\hat{\mathfrak{C}}$ -groups, and let $H(S) := \ker f(S)$. Let $u : G \rightarrow W$ be a morphism of $\hat{\mathfrak{C}}$ -groups which is a section of f (in particular a monomorphism). Then W is the semi-direct product of H and G with an operation of G on H given by the interior automorphism

$$(g, h) \mapsto \mathrm{Int}(u(g)) \cdot h := u(g) \cdot h \cdot u(g)^{-1}$$

for all $g \in G(S)$, $h \in H(S)$ and $S \in \mathrm{Ob} \mathfrak{C}$.

PROOF [SGA 3], Expose I, Prop. 2.3.7 □

The corresponding operation of G on $\mathrm{Lie}(G/S, \mathcal{M})$ is denoted by Ad and is called the *adjoint representation* of G . By definition we have got

$$\mathrm{Ad}(x)X = i^{-1}(s(x)i(X)s(x)^{-1}) \quad \text{for all } x \in G(S') \text{ and } X \in \mathrm{Lie}(G/S, \mathcal{M})(S').$$

If G and H are two group functors over S and if $f : G \rightarrow H$ is a group homomorphism, then (by means of functoriality) we arrive at a commutative diagram with exact rows:

$$\begin{array}{ccccccc} 1 & \longrightarrow & \mathrm{Lie}(G/S, \mathcal{M}) & \longrightarrow & T_{G/S}(\mathcal{M}) & \longrightarrow & G \longrightarrow 1 \\ & & \downarrow \mathrm{Lie}(f) & & \downarrow T(f) & & \downarrow f \\ 1 & \longrightarrow & \mathrm{Lie}(H/S, \mathcal{M}) & \longrightarrow & T_{H/S}(\mathcal{M}) & \longrightarrow & H \longrightarrow 1 \end{array}$$

11.49 Proposition *Let $g \in G(S)$. Then the adjoint representation Ad of G can be written as follows:*

$$\mathrm{Ad}(g) = \mathrm{Lie}(\mathrm{Int}(g)).$$

PROOF $\mathrm{Ad}(g)X \stackrel{\mathrm{def}}{=} i^{-1}(\mathrm{Int}(g)i(X)) = \mathrm{Lie}(\mathrm{Int}(g))(X).$ □

Remark: Recalling Definition 11.6 let us assume that G/S verifies (E). This is for example true if G is representable by an S -group scheme. Then $\mathrm{Ad}(g)$ respects the $\mathcal{Q}_S$ -module structure of $\mathrm{Lie}(G/S, \mathcal{M})$, i.e. in this case Ad is a linear representation of G in the $\mathcal{Q}_S$ -module $\mathrm{Lie}(G/S, \mathcal{M})$:

$$\mathrm{Ad} : G \rightarrow \mathrm{Aut}_{\mathcal{Q}_S\text{-lin}}(\mathrm{Lie}(G/S, \mathcal{M})).$$

As one is used to from differential geometry, one may identify the Lie-algebra with left-invariant vector fields.

11.50 Proposition *There is an isomorphism*

$$\begin{array}{ccc} \mathrm{Hom}(G, \mathrm{Lie}(G/S, \mathcal{M})) & \xrightarrow{\sim} & \Gamma(T_{G/S}(\mathcal{M})/G), \\ f & \mapsto & S_f \end{array}$$

where $S_f(g) := i(f(g)) \cdot s(g)$ for all $g \in G(S')$ and for all $S' \rightarrow S$.

PROOF see [SGA 3], page 61 □

If h is an automorphism of functors over S , to each section t of $T_{G/S}(\mathcal{M})$ can be associated the unique section $h(t)$ making the following diagram commutative:

$$\begin{array}{ccc} G & \xrightarrow{t} & T_{G/S}(\mathcal{M}) \\ h \downarrow & & \downarrow T(h) \\ G & \xrightarrow{h(t)} & T_{G/S}(\mathcal{M}) \end{array}$$

In particular, we can choose h to be the right-translation with $x \in G(S)$:

$$h(g) := t_x(g) := g \cdot x \quad g \in G(S') \text{ and } S' \rightarrow S.$$

Then the isomorphism of Proposition 11.50 is compatible with right-translations. In particular those sections, which are invariant under right translations, are mapped to constant morphisms of G to $\mathrm{Lie}(G/S, \mathcal{M})$, i.e. to morphisms which factorize over S .

11.51 Proposition *The map*

$$\begin{aligned} \mathrm{Lie}(G/S, \mathcal{M})(S) &\rightarrow \Gamma(T_{G/S}(\mathcal{M})/G) \\ X &\mapsto (x \mapsto X \cdot x), \end{aligned}$$

is an isomorphism of $\mathrm{Lie}(G/S, \mathcal{M})(S)$ to those sections of $\Gamma(T_{G/S}(\mathcal{M})/G)$ which are invariant under right-translation.

PROOF [SGA 3], Expose II, Prop. 4.1.2 □

11.4.3. The maps Ad and ad

11.52 Definition A $\mathcal{Q}_S$ -module $\mathcal{F}$ is called *good*, if the canonical morphism

$$\mathcal{F} \otimes_{\mathcal{Q}_S} T_{\mathcal{Q}_S/S}(\mathcal{M}) \rightarrow T_{\mathcal{F}/S}(\mathcal{M})$$

is an isomorphism.

11.53 Definition A group functor G over S is called *good*, if it verifies (E) (see Definition 11.6) and if $\mathrm{Lie}(G/S)$ is a good $\mathcal{Q}_S$ -module.

Remark: Let us remark that a group functor G over S which is representable by a scheme G is good. This will be the case in the physical situations we are going to consider. For more details on “good $\mathcal{Q}_S$ -modules” see e.g. [SGA 3], Expose II, Chap. 4. We are just interested in some theorems about good modules which show that many classical, differential geometric results concerning the Lie-algebra of Lie-groups carry over to algebraic geometry.

11.54 Proposition *If $\mathcal{F}$ is a good $\mathcal{Q}_S$ -module, the $\mathcal{Q}_S$ -module structure of $\mathrm{Lie}(\mathcal{F}/S, \mathcal{M})$ is induced by that of $\mathcal{F}$.*

PROOF [SGA 3], Expose II, Cor. of Def. 4.4 □

11.55 Proposition *If $\mathcal{F}$ is a good $\mathcal{Q}_S$ -module, there is a functorial isomorphism*

$$\mathrm{Lie}(\mathrm{Aut}_{\mathcal{Q}_S\text{-lin}}(\mathcal{F})/S, \mathcal{M}) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{Q}_S\text{-lin}}(\mathcal{F}, \mathrm{Lie}(\mathcal{F}/S, \mathcal{M})).$$

In particular:

$$\mathrm{Lie}(\mathrm{Aut}_{\mathcal{Q}_S\text{-lin}}(\mathcal{F})/S) \xrightarrow{\sim} \mathrm{End}_{\mathcal{Q}_S\text{-lin}}(\mathcal{F}).$$

PROOF [SGA 3], Expose II, Prop. 4.6 □

Let now G be a good S -group-scheme. Then $\mathrm{Lie}(G/S)$ is a good $\mathcal{Q}_S$ -module. We already introduced the linear representation Ad of G in the $\mathcal{Q}_S$ -module $\mathrm{Lie}(G/S)$:

$$\begin{aligned} \mathrm{Ad} : G &\longrightarrow \mathrm{Aut}_{\mathcal{Q}_S\text{-lin}}(\mathrm{Lie}(G/S)) \\ g &\mapsto \mathrm{Lie}(\mathrm{Int}(g)). \end{aligned}$$

Applying the functor Lie and using Proposition 11.55, we arrive at the $\mathcal{Q}_S$ -module morphism ad :

$$\text{ad}: \text{Lie}(G/S) \longrightarrow \text{End}_{\mathcal{Q}_S\text{-lin}}(\text{Lie}(G/S))$$

which may also be considered as a bilinear morphism

$$\text{Lie}(G/S) \times_S \text{Lie}(G/S) \longrightarrow \text{Lie}(G/S), \quad (x, y) \mapsto [x, y] := \text{ad}(x)(y)$$

where $x, y \in \text{Lie}(G/S)(S') = \text{Lie}(G_{S'}/S')(S')$.

11.56 Proposition *Let G be a representable group functor over S (or more generally a group functor which is a subfunctor of a representable group over S). Then $\text{Lie}(G/S)$ is a $\mathcal{Q}_S$ -Lie-algebra, i.e.:*

- (i) $[x, x] = 0$;
- (ii) $[x, y] + [y, x] = 0$;
- (iii) $[x, [y, z]] + [y, [z, x]] + [z, [x, y]] = 0$.

PROOF [SGA 3], Expose II, pages 68-72 □

11.4.4. The Lie-algebra and invariant differential forms

In subsection 11.4.2 we stated several results about the Lie-algebra of an S -group scheme G . In analogy to differential geometry, Proposition 11.50 states the one-to-one correspondence of elements of the Lie-Algebra $\text{Lie}(G/S, \mathcal{M})(S)$ and left-invariant global sections of the tangent-bundle $\Gamma(G, T_{G/S}(\mathcal{M}))$.

In this paragraph we will recall these statements from the dual point of view, i.e. we will consider differential forms instead of vector fields. In order to this, we first introduce the notion of translations on an S -group scheme G for a given T -valued point $g: T \rightarrow G$, i.e. an S -morphism from an S -scheme T to G .

Throughout this section, let G be a group scheme over a fixed scheme S . First we want to introduce the notion of translations on G . In order to do this, consider a T -valued point

$$g: T \rightarrow G$$

of G ; i.e., an S -morphism from an S -scheme T to G . Then g gives rise to the T -valued point

$$g_T := (g, \text{id}_T): T \rightarrow G_T := G \times_S T$$

of the T -scheme $G_T := G \times_S T$. If $p_1: G_T \rightarrow G$ denotes the first projection, we have $g = p_1 \circ g_T$. In the special case where $T := G$ and where $g := \text{id}_G$ is the so called universal point of G , the morphism g_T equals the diagonal morphism Δ of G . For any

other T -valued point g of G , the morphism g_T is obtained from Δ by performing the base change $g : T \rightarrow G$.

As usual, let $m : G \times_S G \rightarrow G$ be the group law of G and write m_T for its extension when a base change $T \rightarrow S$ is applied to G . Then, for any T -valued point g of G , we define the left translation by

$$\tau_g : G_T \xrightarrow{\sim} T \times_T G_T \xrightarrow{g_T \times \text{id}} G_T \times_T G_T \xrightarrow{m_T} G_T$$

and the right translation by

$$\tau'_g : G_T \xrightarrow{\sim} G_T \times_T T \xrightarrow{\text{id} \times g_T} G_T \times_T G_T \xrightarrow{m_T} G_T.$$

Both morphisms are isomorphisms. Quite often we will drop the index T and characterize the map τ_g by writing

$$\tau_g : G \rightarrow G, \quad x \mapsto gx;$$

the same procedure will be applied for τ'_g and for similar morphisms. In the special case where $T := G$ and $g := \text{id}_G$ is the universal point, τ_g is the so-called *universal left translation*, namely the morphism

$$\Phi : T \times_S G \rightarrow T \times_S G, \quad (x, y) \mapsto (x, xy).$$

Similarly, τ'_g gives rise to the *universal right translation*

$$\Psi : G \times_S T \rightarrow G \times_S T, \quad (x, y) \mapsto (xy, y).$$

Each left translation by a T -valued point $g : T \rightarrow G$ is obtained from the universal left translation Φ by performing the base change $g : T \rightarrow G$; in a similar way one can proceed with right translations.

Now let us consider the sheaf $\Omega_{G/S}^i$ of relative differential forms of some degree $i \geq 0$ on G ; it is defined as the i -th exterior power of $\Omega_{G/S}^1$ (see Definition 11.34). For any S -scheme T and any T -valued point $g \in G(T)$, the left translation $\tau_g : G_T \rightarrow G_T$ gives rise to an isomorphism⁸

$$\tau_g^* \Omega_{G_T/T}^i \xrightarrow{\sim} \Omega_{G_T/T}^i.$$

A global section ω in $\Omega_{G/S}^i$ is called *left-invariant* if $\tau_g^* \omega_T = \omega_T$ in $\Omega_{G_T/T}^i$ for all $g \in G(T)$ and all T , where ω_T is the pull-back of ω with respect to the projection $p_1 : G_T \rightarrow G$. Using right translations τ'_t , one defines *right-invariant* differential forms in the same way. Since each translation on the group scheme G_T is obtained by base change from the universal translation, it is clear that one has to check the invariance under translations only for the universal translation. Generally, in connection with translations, we will drop the index T and write ω instead of ω_T if no confusion is possible. Within this context, let us state the following two remarks:

⁸This isomorphism is the canonical pull-back morphism induced by τ_g . In the special case of differential forms it was introduced in Definition 1.3; for the general case of forms of higher degree see Definition 7.20.

- a) All morphisms $g_T : T \rightarrow G_T$ for a given $g : T \rightarrow G$ are induced by base change from $\Delta : G \rightarrow G \times_S G$.
- b) Consider two global sections ω and ω' of a sheaf F on G , let F_T denote the pull-back of F to G_T . Then:

$$\omega = \omega' \Leftrightarrow g_T^* \omega_T = g_T^* \omega'_T \in g_T^* F_T.$$

Similarly two sheaves are isomorphic if their restriction to each T -valued point of G are isomorphic.

11.57 Proposition *Let G be an S -group scheme with unit section $\varepsilon : S \rightarrow G$. Then, for each $\omega_0 \in \Gamma(S, \varepsilon^* \Omega_{G/S}^i)$, there exists a unique left-invariant differential form $\omega \in \Gamma(G, \Omega_{G/S}^i)$ such that $\varepsilon^* \omega = \omega_0$ in $\varepsilon^* \Omega_{G/S}^i$. The same is true for right-invariant differential forms.*

PROOF Let us shortly indicate, how the left invariant differential form ω is obtained from ω_0 . For details of the prove, we refer the reader to [BLR], Prop 4.2/1. Due to the uniqueness assertion, the problem is local in S (because we may glue). Furthermore it suffices to consider the case $i = 1$. Thus we may assume that ω_0 lifts to a section ω' of $\Omega_{G/S}^1$ which is defined over a neighborhood U of the unit section. This is due to the fact that $(\varepsilon^* \Omega_{G/S}^1)(S) = (\Omega_{G/S}^1)(U) \otimes_{\mathcal{O}_G(U)} \mathcal{O}_S(S)$ if S and $U \subset G$ are affine, and that $p^* : \mathcal{O}_S(S) \hookrightarrow \mathcal{O}_G(U)$ is injective, because $p : G \rightarrow S$ has the section ε . Then the decomposition $p_1^* \Omega_{G/S}^1 \oplus p_2^* \Omega_{G/S}^1 \xrightarrow{\sim} \Omega_{G \times_S G/S}^1$ of Proposition 1.7 gives a decomposition $m^* \omega' = \omega_1 \oplus \omega_2$ over $m^{-1}(U)$, where $m : G \times_S G \rightarrow G$ is the multiplication. If $\delta : G \rightarrow G \times_S G$, $x \mapsto (x, x^{-1})$, denotes the twisted diagonal morphism, $m^* \omega'$ is defined in a neighborhood of the image of δ so that $\delta^* \omega_2$ gives rise to a global section ω of $\Omega_{G/S}^1$. Then ω is the searched left-invariant differential form with $\varepsilon^* \omega = \omega_0$ in $\varepsilon^* \Omega_{G/S}^1$.

In particular the association $\omega_0 \mapsto \omega$ is $\mathcal{O}_S(S)$ -linear. $\square$

11.58 Proposition *There are canonical isomorphisms*

$$p^* \varepsilon^* \Omega_{G/S}^i \xrightarrow{\sim} \Omega_{G/S}^i \quad \text{for all } i \in \mathbb{N}$$

which are obtained by extending sections in $\varepsilon^ \Omega_{G/S}^i$ to left-invariant sections in $\Omega_{G/S}^i$. Similar isomorphisms are obtained by using right-invariant differential forms.*

PROOF [BLR], Prop 4.2/2 $\square$

As a direct consequence we obtain:

11.59 Corollary *Let G be a smooth S -group scheme of relative dimension n , and assume that $\varepsilon^* \Omega_{G/S}^i$ is free. Then $\Omega_{G/S}^i$ is free $\mathcal{O}_S$ -module generated by $\binom{n}{i}$ left-invariant differential forms of degree i . The same is true for right-invariant differential forms.*

In physical applications, space-time is given by the set of S -sections $X(S)$ of a smooth, separated algebraic space $X \rightarrow S$ with certain further properties (see Definition 0.4 for a complete list of properties of X). In special cases, X is even a group object and thus carries a canonical CPT -operation (see Definition 0.7). Then, all results on group schemes, which are summarized above, are valid for X . The preceding Corollary 11.59 is the basis for the proof that there exists a volume form v in X . This result is essential, because we need to make use of a volume form within our physical purposes of a Yang-Mills theory over rings.

11.60 Definition Let $X \rightarrow S$ be a smooth S -scheme of relative dimension n (in particular $\Omega_{X/S}^n$ is a line-bundle). Let us assume that there exists a global section $v \in \Omega_{X/S}^n(X)$ which generates $\Omega_{X/S}^n$. Then v is called a *volume form* on X .

11.61 Theorem Let $S = \text{Spec } R$ be the spectrum of a Dedekind ring R which is a principal ideal domain. Let $X \rightarrow S$ be a smooth S -group scheme of relative dimension n . Then the sheaves $\Omega_{X/S}^i$ are free $\mathcal{O}_X$ -modules generated by left-invariant differential forms of degree i . In particular there exists a volume form on X .

PROOF By Corollary 11.59 it suffices to show that $\varepsilon^* \Omega_{X/S}^i$ is a free $\mathcal{O}_S$ -module. We already know that $\varepsilon^* \Omega_{X/S}^i$ is a locally free $\mathcal{O}_S$ -module, because it is the pull-back of the locally free $\mathcal{O}_X$ -module $\Omega_{X/S}^i$. Due to the well-known fact that vector bundles over a scheme S and locally free $\mathcal{O}_S$ -modules are in one-to-one correspondence, it suffices to show that (up to isomorphism) there are only trivial vector bundles over S ; i.e. it suffices to show that $K_0(R) \cong \mathbb{Z}$. But the ideal class group $\text{Cl}(R)$ of R is the trivial group $\{1\}$ if and only if R is a principal ideal domain. Thus we are done by the following result which may be found in nearly every book on algebraic K -theory. $\square$

11.62 Proposition Let R be a Dedekind ring and let $\text{Cl}(R)$ denote the ideal class group of R . Then there is an isomorphism of groups:

$$K_0(R) \cong \mathbb{Z} \oplus \text{Cl}(R)$$

11.63 Corollary Let $X \rightarrow S$ be a model of type (SR) (see Definition 0.7). Then there exists a volume form v on X .

As we are finally interested in physical applications, let us from now on assume that G is a smooth S -group scheme of relative dimension n , and that there is a left-invariant differential form $\omega \in \Omega_{G/S}^n(G)$ generating $\Omega_{G/S}^n$ as an $\mathcal{O}_G$ -module. Performing a base change with the canonical morphism $G_K \rightarrow G$ (where K is the field of fractions of S), ω is pulled back to a left-invariant differential form $\omega_K \in \Omega_{G_K/K}^n(G_K)$ generating $\Omega_{G_K/K}^n$. Interpreting the generic fibre as classical limit (as explained in section 6.1), we may interpret ω_K as the classical limit of ω . Let us finish this section with the prove that the volume form ω_K is invariant with respect to translations if G describes space-time itself. By Theorem 3.12 the latter means that the generic fibre G_K of G is given by

an abelian variety. In particular, G_K is bounded by Proposition 3.23, i.e. $G_K(K^{sh})$ is bounded. More precisely, we claim that ω_K is not only invariant with respect to left-translations, but that furthermore $|\tau'_g{}^*\omega_K| = |\omega_K|$ for all right-translations by physical points $g \in G_K(K^{sh})$ and for all valuation $|\cdot|$ (i.e. prime spots $\mathfrak{p}$) corresponding to closed points $s \in S$. Before stating the proof of this translation invariance, let us first recall the definition of the interior automorphism which was already considered in subsection 11.4.2.

11.64 Definition For an arbitrary T -valued point g of G we define the interior automorphism

$$\text{Int}(g) := \tau_g \circ \tau'_{g^{-1}} : G \longrightarrow G, \quad x \mapsto gxg^{-1}$$

induced by g .

Now our claim is a direct consequence of the following two propositions.

11.65 Proposition *There exists a unique group homomorphism $\chi : G \rightarrow \mathbb{G}_m$ (a so called character on G), such that*

$$\text{Int}(g)^*\omega = \tau'_{g^{-1}}{}^*\omega = \chi(g)\omega$$

for each T -valued point g of G .

PROOF [BLR], Prop. 4.2/4 □

11.66 Proposition *Let G_K be a smooth K -group of relative dimension n , and assume that $G_K(K)$ (resp. $G_K(K^{sh})$) is bounded in G_K . Then the character χ considered in Proposition 11.65 satisfies $|\chi(g)| = 1$ for each $g \in G_K(K^{sh})$ (resp. each $g \in G_K(K^{sh})$).*

PROOF [BLR], Prop. 4.2/5 □

11.5. Faithfully flat descent

Let $p : S' \rightarrow S$ be a morphism of schemes and consider the functor $\mathcal{F} \rightarrow p^*\mathcal{F}$ which associates to each quasi-coherent $\mathcal{O}_S$ -module $\mathcal{F}$ its pull-back under p . Then, in its simplest form, the problem of descent relative to $p : S' \rightarrow S$ is to characterize the image of this functor. But descent may also be viewed as a natural generalization of a patching problem. The procedure of solution is as follows. Set $S'' := S' \times_S S'$, and let $p_i : S'' \rightarrow S'$ be the projection onto the i -th factor ($i = 1, 2$). For any quasi-coherent $\mathcal{O}_{S'}$ -module $\mathcal{F}'$, call an S'' -isomorphism $\varphi : p_1^*\mathcal{F}' \rightarrow p_2^*\mathcal{F}'$ a *covering datum* of $\mathcal{F}'$. Then the pairs $(\mathcal{F}', \varphi)$ of quasi-coherent $\mathcal{O}_{S'}$ -modules with covering data form a category in a natural way. A morphism between two such objects $(\mathcal{F}', \varphi)$ and $(\mathcal{G}', \psi)$ consists of an

$\mathcal{O}_{S'}$ -module-homomorphism $f : \mathcal{F}' \rightarrow \mathcal{G}'$ which is compatible with the covering data φ and ψ ; thereby we mean that the diagram

$$\begin{array}{ccc} p_1^* \mathcal{F}' & \xrightarrow{\varphi} & p_2^* \mathcal{F}' \\ p_1^* f \downarrow & & \downarrow p_2^* f \\ p_1^* \mathcal{G}' & \xrightarrow{\psi} & p_2^* \mathcal{G}' \end{array}$$

is commutative.

Starting with a quasi-coherent $\mathcal{O}_S$ -module $\mathcal{F}$, we have a natural covering datum on $p^* \mathcal{F}$ which consists of the canonical isomorphism

$$p_1^*(p^* \mathcal{F}) \cong (p \circ p_1)^* \mathcal{F} = (p \circ p_2)^* \mathcal{F} \cong p_2^*(p^* \mathcal{F}).$$

So we can interpret the functor $\mathcal{F} \rightsquigarrow p^* \mathcal{F}$ as a functor into the category of quasi-coherent $\mathcal{O}_{S'}$ -modules with covering data. It is this functor which will be of interest in the following. It turns out that this functor is fully faithful if $p : S' \rightarrow S$ is faithfully flat and quasi-compact (see e.g. [BLR], Prop. 6.1/1). Furthermore, it is an equivalence of categories if, instead of covering data, we consider descent data. A descent datum is a special covering datum which satisfies a certain cocycle condition. In order to introduce them, set $S''' := S' \times_S S' \times_S S'$, and let $p_{ij} : S''' \rightarrow S''$ be the projection onto the factors with indices i and j for $i < j$; $i, j = 1, 2, 3$. In order that a quasi-coherent $\mathcal{O}_{S'}$ -module $\mathcal{F}'$ with covering datum $\varphi : p_1^* \mathcal{F}' \rightarrow p_2^* \mathcal{F}'$ belongs to the essential image of the functor $\mathcal{F} \rightsquigarrow p^* \mathcal{F}$, it is necessary that the diagram

$$\begin{array}{ccc} p_{12}^* p_1^* \mathcal{F}' & \xrightarrow{p_{12}^* \varphi} & p_{12}^* p_2^* \mathcal{F}' = p_{23}^* p_1^* \mathcal{F}' \xrightarrow{p_{23}^* \varphi} p_{23}^* p_2^* \mathcal{F}' \\ \parallel & & \parallel \\ p_{13}^* p_1^* \mathcal{F}' & \xrightarrow{p_{13}^* \varphi} & p_{13}^* p_2^* \mathcal{F}' \end{array}$$

is commutative; the unspecified identities are the canonical ones. Namely, if $\mathcal{F}'$ is the pull-back under p of a quasi-coherent $\mathcal{O}_S$ -module and if φ is the natural covering datum on $\mathcal{F}'$, then the diagram is commutative, because all occurring isomorphisms are the identical ones. The commutativity of the diagram is referred to as the *cocycle condition* for φ ; in short, we can write it as

$$p_{13}^* \varphi = p_{23}^* \varphi \circ p_{12}^* \varphi.$$

It corresponds to the usual cocycle condition on triple overlaps when a global object is to be constructed by gluing local parts. A covering datum φ on $\mathcal{F}'$ which satisfies the cocycle condition is called a *descent datum* on $\mathcal{F}'$. The descent datum is called *effective* if the pair $(\mathcal{F}', \varphi)$ is isomorphic to the pull-back $p^* \mathcal{F}$ of a quasi-coherent S -module $\mathcal{F}$ where, on $p^* \mathcal{F}$, we consider the canonical descent datum. Also we want to mention that the notions of covering and descent datum are compatible with base change over S . Now we are ready to state the desired result on the descent of quasi-coherent S' -modules due to Grothendieck.

11.67 Theorem *Let $p : S' \rightarrow S$ be faithfully flat and quasi-compact. Then the functor $\mathcal{F} \rightsquigarrow p^*\mathcal{F}$, which goes from quasi-coherent $\mathcal{O}_S$ -modules to quasi-coherent $\mathcal{O}_{S'}$ -modules with descent data, is an equivalence of categories.*

PROOF [BLR], Thm. 6.1/4 □

11.68 Remark (Étale coverings). Consider a quasi-separated scheme S and a finite étale covering $(S_i \rightarrow S)_{i \in I}$ of S (see Definition 7.5). Let $S' := \coprod_{i \in I} S_i$ be the disjoint union of the S_i , and let $p : S' \rightarrow S$ be the canonical projection. Note that p is faithfully flat and quasi-compact. A quasi-coherent $\mathcal{O}_{S'}$ -module $\mathcal{F}'$ may be thought of as a family of $\mathcal{O}_{S_i}$ -modules $\mathcal{F}_i$. Under what conditions does $\mathcal{F}'$ descent to a quasi-coherent $\mathcal{O}_S$ -module $\mathcal{F}$; i.e. under what conditions can one glue the $\mathcal{F}_i$ in order to obtain a quasi-coherent $\mathcal{O}_S$ -module $\mathcal{F}$ from them? By Theorem 11.67 we need a descent datum for $\mathcal{F}'$ with respect to $p : S' \rightarrow S$. Such a datum consists of an isomorphism $\varphi : p_1^*\mathcal{F}' \xrightarrow{\sim} p_2^*\mathcal{F}'$ satisfying the cocycle condition, where p_1 and p_2 are the projections from S'' to S' . In our case, we have

$$S'' = S' \times_S S' = \coprod_{i,j \in I} S_{ij},$$

where $S_{ij} := S_i \times_S S_j$. Thus the isomorphism φ consists of a family of isomorphisms

$$\varphi_{ij} : \mathcal{F}_i|_{S_{ij}} \xrightarrow{\sim} \mathcal{F}_j|_{S_{ij}}$$

satisfying the cocycle condition, namely, the condition that

$$\varphi_{ik}|_{S_{ijk}} = \varphi_{jk}|_{S_{ijk}} \circ \varphi_{ij}|_{S_{ijk}}$$

for all $i, j, k \in I$, where $S_{ijk} := S_i \times_S S_j \times_S S_k$. Thereby, $\mathcal{F}_i|_{S_{ij}}$ denotes the pull-back $p_i^*\mathcal{F}_i$ of $\mathcal{F}_i$ under the canonical projection $p_i : S_i \times_S S_j \rightarrow S_i$, etc.

So the descent datum φ in the form of the above family of isomorphism is the patching datum for the $\mathcal{O}_{S_i}$ -modules $\mathcal{F}_i$ with respect to étale topology on S .

Keeping the morphism $S' \rightarrow S$, one may also study the problem of when an S' -scheme X' descends to an S -scheme X . The general setting will be the same as in the case of quasi-coherent modules, and the definitions we have given can easily be adapted to the new situation. For example, a *descent datum on an S' -scheme X'* is an S'' -isomorphism

$$\phi : p_1^*X' \rightarrow p_2^*X'$$

which satisfies the cocycle condition; p_i^*X' is the scheme obtained from X' by applying the base change $p_i : S'' \rightarrow S'$. Again there is a canonical functor $X \rightsquigarrow p^*X$ from S -schemes to S' -schemes with descent data.

11.69 Theorem *Let $p : S' \rightarrow S$ be faithfully flat and quasi-compact. Then the functor $X \rightsquigarrow p^*X$ from S -schemes to S' -schemes with descent data is fully faithful.*

PROOF [BLR], Thm. 6.1/6 □

Bibliography

- [ADFV] I.Ya. Aref'va, B. Dragovich, P.H. Frampton and I.V. Volovich; Wave function of the universe and p -adic gravity, International Journal of Modern Physics A, Volume 6, Issue 24, pp. 4341-4358, 1991
- [ADV] I.Ya. Aref'va, B. Dragovich and I.V. Volovich; *Phys. Lett.* **B209**, No. 445 (1988), **B212**, No. 283 (1988), **B214**, No. 339 (1988)
- [BF] L. Brekke, Peter G.O. Freund, p -adic numbers in physics, *Phys. Rep.* **233**, No. 1 (1993)
- [BLR] Bosch, Siegfried, Lütkebohmert, Werner and Raynaud, Michel; *Néron models*, Springer-Verlag, Berlin, Heidelberg, New York, 1990
- [Bosch] Bosch, Siegfried; Lectures on Formal and Rigid Geometry, Preprintreihe des SFB 478 - Geometrische Strukturen in der Mathematik, Heft 378, 2005
- [CD] Choquet-Bruhat, Dewitt-Morette; *Analysis, Manifolds and Physics: Part I,II Topology and Physics*, Elsevier science B.V., Amsterdam, 2000
- [DD] G.S. Djordjevic, B. Dragovich; p -adic and adelic path integrals, *Proceedings of the XI Yugoslav Conference on the Nuclear and Particle Physics, Studenica, Yugoslavia* (1998) pp. 312-315
- [DFU] B. Dragovich, P.H. Frampton and C.V. Urosevic; Classical p -adic space-time, *Mod. Phys. Lett.* **A5** (1990) 1521-1528
- [DN1] B. Dragovich, L.D. Nesic; On p -adic models of the universe, *Facta Universitatis: Series "Physics, Chemistry and Technology"* **1** (1997) 223-236
- [DN2] B. Dragovich, L.D. Nesic; Adelic models of the universe, *Proceedings of the Conference BPU-3, Cluj-Napoca (Romania), Balkan Physics Letters* **5 Suppl.** (1997) 104-107
- [DN3] B. Dragovich, L.D. Nesic; Adelic quantum cosmology, Conference 'Gravitation, Cosmology and Relativistic Astrophysics', 8.-11. Nov. 2001, Kharkov, Ukraine. *Collection of Papers (Kharkov University, 2001)* pp. 31-34
- [DN4] B. Dragovich, L.D. Nesic; On p -adic numbers in gravity, *Balkan Physics Letters* **6** (1998) 78-81
- [DDN1] G.S. Djordjevic, B. Dragovich, L.D. Nesic; p -adic and adelic free relativistic particle, *Modern Physics Letters* **7** (1999) 150-154

- [DDN2] G.S. Djordjevic, B. Dragovich, L.D. Netic; Adelic path integrals for quadratic lagrangians, *Infinite Dimensional Analysis, Quantum Probability and Related Topics* **6** (2003) 179-195
- [DDNV] G.S. Djordjevic, B. Dragovich, L.D. Netic, I.V. Volovich; p-adic and adelic minisuperspace quantum cosmology, *International Journal of Modern Physics* **A17** (2002) 1413-1433
- [Dra1] Branko Dragovich; Adelic harmonic oscillator, *Int. J. Mod. Phys.* **A10** (1995) 2349-2365
- [Dra2] Dragovich, Branko; Non-archimedean geometry and physics on adelic spaces, arxiv: math-ph/0306023
- [Dra3] Dragovich, Branko; p-adic and adelic cosmology: p-adic origin of dark energy and dark matter, arXiv: hep-th/0602044
- [Dra4] Dragovich, Branko; Adelic aspects of quantum cosmology, *Proceedings elementary particle physics*, Moskau (1999), pp. 116-122. Ed. by A.I. Studenikin
- [Dra5] Dragovich, Branko; Invariance in adelic quantum mechanics, arXiv: hep-th/0612063
- [Dra6] Dragovich, Branko; Adelic Wave Function of the Universe, in *Proc. the Third A. Friedmann Int. Seminar on Grav. and Cosmology*, St. Petersburg, pp. 311-321 (1995)
- [Dra7] Dragovich, Branko; Adelic Wave Function of the de Sitter Universe, in *Proc. 7th Lomonosov Conf. on Elem. Particle Physics*, Moscow, pp. 137-140 (1995)
- [DNF] Dubrovin, Novikov, Fomenko; *Modern geometry - methods and applications*, Springer-Verlag, Berlin, Heidelberg, 1990
- [GS] Gökeler, Schücker; *Differential Geometry, Gauge Theories and Gravity*, Cambridge University Press, 1987
- [EGA II] Grothendieck, Alexander and Dieudonné Jean; *Etude globale élémentaire de quelques classes de morphismes*, Publ. Math IHES 8 (1961)
- [EGA IV] Grothendieck, Alexander and Dieudonné Jean; *Etude local de schémas et des morphismes de schémas*, Publ. Math IHES 20 (1961), 24 (1965), 28 (1966), 32 (1967)
- [SGA 3] Grothendieck, Alexander; *Schémas en groupes I*, Lect. Notes Math. 151, Springer-Verlag, Berlin Heidelberg New York, 1970
- [Har] Hartshorne, Robin; *Algebraic geometry*, Springer-Verlag, New York, Berlin, Heidelberg, 5. Aufl. edition, 1977
- [Haw] J. Hartle and S. Hawking, *Phys. Rev.* **28**, 2960 (1983)
- [Hea] Heath-Brown, D.R.; Cubic forms in ten variables, *Proceedings of the London Mathematical Society*, 47(3), pp. 225-257, 1983

- [KN] Kobayashi, Nomizu; *Foundations of differential geometry, Vol.1*, John Wiley & Sons, Inc., 1963
- [Liu] Liu, Qing; *Algebraic geometry and arithmetic curves*, Oxford University Press, Oxford, New York, 2002
- [MM] Marathe, Martucci; *The mathematical foundations of gauge theory*, Studies in Mathematical Physics Vol. 5, North-Holland, Amsterdam, 1992
- [Meu] Y. Meurice; *Int. J. Mod. Phys. A***4**, 5133 (1989)
- [Milne] Milne, J.S.; *Etale cohomology*, Princeton Math. Series 33, Princeton University Press, 1980
- [Mor] Mordell, L.J.; A remark on indeterminate equations in several variables, *Journal of the London Mathematical Society*, 12, pp. 127-129, 1937
- [Nab] Naber; *Topology, geometry and gauge fields: Foundations*, Springer-Verlag, New York, 1997
- [Nak] Nakahara; *Geometry, topology and physics*, Taylor & Francis Group, Abington, 2003
- [Neu] Neukirch, Jürgen; *Algebraische Zahlentheorie*, Springer-Verlag, Berlin Heidelberg New York, 1992
- [Schar] Scharlau, Winfried; *Quadratic and Hermitian Forms*, Springer-Verlag, Berlin Heidelberg, 1985
- [Serre1] Serre, Jean Pierre; *A course in arithmetic*, Springer-Verlag, New York, 1973
- [Serre2] Serre, Jean Pierre; *Groupes algebriques et corps de classes*, Hermann, Paris, 1959
- [Vla] V.S. Vladimirov, *Lett. Math. Phys.* **27**, 123 (1993)
- [VV1] V.S. Vladimirov and I.V. Volovich, *Dokl. Akad. Nauk SSSR* **302**, 320 (1988)
- [VV2] V.S. Vladimirov and I.V. Volovich, *Commun. Math. Phys* **123**, 659 (1989)
- [VV3] V.S. Vladimirov and I.V. Volovich, *Phys. Lett. B***217**, 411 (1989)
- [VVZ] V.S. Vladimirov, I.V. Volovich and E.I. Zelenov, *p-adic analysis and mathematical physics*, World Scientific Publishing Co.Pte.Ltd., 1994
- [Vol1] I.V. Volovich; Number theory as the ultimate physical theory, CERN-TH.4781/87
- [Wey] H. Weyl, *The theory of groups and quantum mechanics* (Dover, New York, 1931)
- [Zel1] E.I. Zelenov, *Teor. Mat. Fiz.* **80**, 253 (1989)
- [Zel2] E.I. Zelenov, *Teor. Mat. Fiz.* **86**, 375 (1991)